



# **Extent of Home Based Care Program Support to Hospital Based Care for Children with Chronic and Terminal Illness in Meru County, Kenya**

**P. K. Kubai<sup>1\*</sup>, M. R. Kei<sup>1</sup> and A. M. Mutema<sup>1</sup>**

<sup>1</sup>Meru University of Science and Technology, P.O.Box 972-60200, Meru, Kenya.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author PKK designed the study, performed the statistical analysis, developing the study protocols, and wrote the first draft of the manuscript. Authors AMM and MRK reviewed data analyses and managed the literature searches. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JAMMR/2021/v33i1330951

### Editor(s):

(1) Dr. Ashish Anand, GV Montgomery Veteran Affairs Medical Center, USA and University of Mississippi Medical Center, USA and William Carey School of Osteopathic Medicine, USA.

### Reviewers:

(1) Azam Doustmohammadian, Iran University of Medical Sciences, Iran.

(2) L. Manoj Kumar, St Thomas College of Nursing, Kerala.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/69310>

**Original Research Article**

**Received 28 March 2021**

**Accepted 03 June 2021**

**Published 05 June 2021**

## **ABSTRACT**

**Introduction:** Chronic and Terminal illnesses continue to increase and aggravate the burden of disease and the diminish space in our hospitals and communities, Worldwide, 57 million persons died in 2008, an estimated 40 million were in need of HBCP, 6.6 - 10.8 million Children died, 98% Chronic and Terminal illnesses (CI/TI) are found in low and middle-income Countries. Chronic and Terminal illnesses in Children are on the rise in Sub Saharan Africa. Kenya lags in implementation of Home Based Care to mitigate effects of CI/TI. WHO, 2017 and Ministry of Health-Kenya 2013 shows that Children are affected by these Illnesses. These illnesses have made families' to suffer psychosocial and economic hardships. Evaluation of determinants of Home Based Care Program (EHBCP) services by assessing if effective and quality delivery of HBCP Supports Hospital Based Care is key.

**Aims:** To determine the extent to which Home Based Care Program services Support to Hospital Based Care for Children aged 1-14 years diagnosed with selected Chronic and Terminal illnesses in Meru County Kenya.

**Study Design:** A descriptive Cross Sectional Survey.

\*Corresponding author: E-mail: [patrickkubai27@gmail.com](mailto:patrickkubai27@gmail.com), [pkubai@must.ac.ke](mailto:pkubai@must.ac.ke);

**Place and Duration of Study:** Meru County, Kenya between June 2018 and Dec 2019.

**Methodology:** Methodology: Descriptive Cross Sectional Survey of 245 Caregivers of Children diagnosed with Chronic and Terminal illness were selected by proportionate to size sampling and simple random sampling from Meru County Health Facilities.

**Results:** Delivery of quality and effective Home Based Care Program was positively associated with age, return rate of over 100% was achieved with about 245 respondents being interviewed (132%), experience of Health Care Workers (HCWs) 4.8 [95% CI = 1.06 – 21.68, P = 0.041]. HCWs profession, gender and years of work were positively related to Hospital Based Care services 3.03 [95%CI = 1.64 – 5.59, P<0.001]. Use of minimum HBCP package was found to be useful in support of hospital care, as 50.4% reported that its support was of a good or great extent. The trend was observed in all aspects of the minimum HBCP package studied, with 52.4%, 54.4% and 46.4% having the opinion that use of the package by caregivers, in management and mitigation of complications supported hospital care by a good or great extent respectively.

**Conclusion:** Need for an urgent adoption of holistic approach on health care system strengthening by putting up a well-integrated Home Based Care Program with quality Minimum HBCP essential package to support the mainstream health care system. In addition adopt better management practices, planning strategies, resource allocation and monitoring for effective HBCP programmes as per the current guidelines to bridge the gaps identified in planning, management practices to fill the disjointed policy implementation and resource inadequacies in implementation of HBCP services to support Hospital Based care for ultimate delivery of quality care among the Chronically and Terminally ill Children.

**Keywords:** *Determinants; chronic/terminal illnesses; children; home based care program; health care workers; supports and hospital based care.*

## **DEFINITION OF OPERATIONAL TERMS**

**Activities of Daily Living:** *Refers to the routine activities (ADL) that person Undertakes every day with or without the need of assistance, such as: eating, bathing, dressing, toileting, transferring (walking) and continence control.*

**Caregiver:** *Refers to the guardian or biological parent who has been taking care of a child for the last six months.*

**Children:** *Refers to a person who is equal to one year or less than 14 years of age.*

**Client:** *Refers to any child who resides in a home-based Care centers Majority of Children referred to home based care are the subject of Protective services Intervention.*

**Chronic Illness:** *Refers to Health problems/ conditions that requires ongoing management over a long period or decades as stated in WHO.*

**Effectiveness:** *Refers to the level or degree of achievement or accomplishment of HBCP services by HBCP offering health facilities or centers as set in accordance with WHO set standards.*

**Evaluation:** *Refers to assessment or judgement of various aspects HBCP services for Children with chronic and terminal illness management in terms of its context, inputs, processes and products as presented in CIPP model.*

**Health Care Provider:** *Refers to various medical and allied professionals involved in care for Children diagnosed with chronic and terminal illnesses.*

**Hindrances:** *Refer to infrastructural, utilization of HBCP services, policy issues, and Stakeholder's involvement attributes limiting Home Based Care Program services for Children with selected chronic and terminal illnesses.*

**Home Based Care Program (HBCP):** *Refers to planned set of activities, procedures and processes of care designed for provision of home based care services to Children diagnosed with selected chronic and terminal illnesses that is extended from the health facility to the clients' home through family participation and involvement within available resources according to World Health Organization/UNAIDS set standards.*

**Minimum Essential Package:** *Refers to the WHO defined components Prescribed for provision of Home Based Care Programme*

services for Chronically and Terminally ill Children.

**Program:** Refers to “any set of organized activities supported by a set of resources to achieve a specific and intended result.”

**Selected Chronic and Terminal Illnesses:** Refers to identified conditions such as Tuberculosis, Asthma, Cancer, Heart conditions, congenital abnormalities, Stroke, and among other Chronic and Terminal illnesses.

**Training:** Means deliberate process of educating caregivers/Health Care Workers to develop HBCP services delivery skills and competences that are required in management of Children with selected Chronic and Terminal illness.

**Terminal illness:** Refers to disease (s) or medical conditions with fatal outcome irrespective of management options provided.

**Quality:** Means delivery of real time effective and efficient HBCP services to Children at Home in accordance with WHO/UNAIDS set HBC program standards.

## ABBREVIATIONS

ADL	: Activities of Daily Living
AOR	: Adjusted odds Ratios
CDC	: Center for Disease Control
CI	: Chronic Illness
CoC	: Continuum of Care
CIPP	: Context, Input, Process and Product
CPSG	: Psychosocial Support Groups
CS	: Community Strategy
DOH	: Directory of Health Facilities
EHBCP	: Evaluation of Home Based Care Program
GoK	: Government of Kenya
HBCP	: Home Based Care Program
HBCPs	: Home Based Care Program Services
HICs	: High Income Countries
IHBCDPs	: Institutionalized Home Based Care Delivery Points
KDHS	: Kenya Demographic and Health Survey
MEP	: Minimum Essential Package
MoH	: Ministry of Health
MUST	: Meru University of Science and Technology
NCDs	: Non-Communicable Diseases
OIs	: Opportunistic Infections
TI	: Terminal Illness

UNAIDS : The Joint United Nations Programme on HIV/AIDS

UNGA : United Nations General Assembly

WHO : World Health Organization

## 1. INTRODUCTION

According to WHO, Lopez and Mathers [1,2,3], 80% of chronic disease deaths worldwide occur in Lower and Middle Income Countries (LAMICs). According to Ama de-Graft et al. [4] the burden of non-communicable diseases was assessed in twenty three low and middle income Countries and the data shows that they account for 50% of the total disease burden. However, in these Countries Chronic diseases and Terminal illnesses develop at an earlier age than in high-income Countries, often resulting in prolonged period of disability before death [5,6,7]. The rising health care costs, growing interest in moving more patient care from Hospital Based Care to Home Based Care settings [8,9,10,11]. Efforts to facilitate effective and quality delivery of child health care or home based care services which enables a larger percentage of Chronically and Terminally ill Children to receive ultimate health care in their homes [1,12]. As home health care technologies develop, so does the range of Home Based Care Program services and with growing body of evidence suggesting that the location in which Children receive, palliative care has serious implications. Provision of Home Based Care Program services in home based settings, parental satisfaction appears to be high, subsequent adaptation and improved child health outcomes for parents, relatives, siblings and community drastically improve [1,13,14,15]. A Child who is cared for in a home based setting, suffers minimal stress and anxiety compared to the ones being cared for in hospital, clinic or nursing home environment. Clients being cared for in a familiar environment during their illness are more tolerable as they actively participate in family matters. Family heads continue doing so and can be consulted on various family issues which could have been difficult when admitted in hospital or a clinic to make a decision. When the chronically and terminally ill persons are close to their immediate family members, friends and relatives they have a sense of belonging which is not be the case in hospital where health care providers are strangers [16]. This familiar surroundings makes the sick accept their conditions and illnesses, hence contributing to quick recovery from life threatening or limiting illness [1,17,14,18,19]. Kenya remains behind in implementation of Home Based Care to mitigate

effects of CI/TI. WHO [9,20,21,22] and Ministry of Health-Kenya [23] shows that Children continue suffering the effects of these illnesses. These illnesses have made families' to suffer psychosocial and economic hardships. Evaluation of determinants of Home Based Care Program (EHBCP) services by assessing if effective and quality delivery of minimum HBCP essential package Supports Hospital Based Care is key [9,24,20,22].

## 2. MATERIALS AND METHODS

The study was a descriptive cross-sectional survey, conducted in Meru County between June and September 2019. A sample of 245 respondents comprising of the health workers/caregivers was selected using proportional to size, simple random sampling to select the six health facilities across Meru County. Descriptive analysis, multivariate logistic regression, and Odds ratios were pulled from Statistical Package format (IBM SPSS) version 25. The extent to which HBC program services supports Hospital Based Care was used as the dependent variable. Variables such as Provision of Clinical, Nursing services, Minimum essential Package support Hospital Care, Health education and Health Promotion, age, sex, gender, parent / caregiver / guardian age, determinants of HBCP support to provision Hospital Based Care, delivery of HBC program services, determinants of HBCP services delivery points, preferred health facilities for provision of Children HBCP services, Socio- economic, Cultural factors, skills, care competences acquired by Health care providers, HBCP caregivers, impact and processed skills reinforcing concepts through quality delivery of Home Based Care were used as independent variables.

### 2.1 Sampling Method

The required sample size was computed using Taro Yamane formula [25], proportion to size of the population was used to estimate the minimum required sample for the study. A sample of 241 caregivers of Children diagnosed with selected Chronic and Terminal illnesses under Home Based Care Program was selected.

$$n = \frac{N}{1 + N(e)^2}$$

Where:

n = Minimum required sample;

N = Population size;  
e = Degree of tolerance error; with a significance level put at 95%, the degree of error term was 5% (i.e. 0.05).  
Therefore:

$$n = \frac{611}{1 + 611(0.05)^2}$$

n = 241 hence a Sample size of 241. Adjusting the sample upwards for 10% non-response, the sample is;

$$n_1 = \frac{241}{1 - 0.1}$$

Hence n was 268 respondents. Further, since the target population was less than 10000, the sample size was adjusted downwards using finite population correction factor (f.c.f) as follows;

$$n_{f.c.f} = \frac{n_1}{1 + \frac{n_1}{N}}$$

Where  $n_{f.c.f}$  = the minimum desired sample size.

$$n_{f.c.f} = \frac{268}{1 + \frac{268}{611}}$$

$$n_{f.c.f} = 186$$

To ensure that we have a representative sample size, the study utilized proportional to size sampling method with the aid of the proportionality formula was thus used to allocate participants per site:

$$Q = \frac{A}{N} * n \text{ Where:}$$

Q = the number of the questionnaire to be allocated to each proportion.

A = the proportion of each segment.

N = the total population of all the segments.

n = the estimated sample size used in the study.

### 2.2 Data Collection and Analysis

A structured Likert questionnaire was administered. The questionnaires were pre-tested in neighbouring Tharaka Nithi County at Chuka Level 5 Teaching Referral Hospital which borders Meru County. The instruments were tested for reliability yielding a Cronbach's alpha of 0.75 and therefore considered reliable. The data was entered and analyzed using SPSS Software Version 25. Descriptive analysis, multiple regression, and Odds ratios were pulled with utilization and delivery of HBCP services being the dependent variable. The results were

presented using descriptive and inferential statistics.

### 3. RESULTS

#### 3.1 Socio Demographic Characteristics

##### 3.1.1 Gender and age

Five out of seven demographic factors was positively associated with level of Home Based Care for Children with selected Chronic and Terminal illness,  $P < 0.05$ . Age category of the participants was significantly associated with increased level of utilization of Home Based Care which has direct bearing to Hospital Based Care. The proportion of participants indicating that utilization of HBCP was good/very good/excellent for Children with CI/TI was observed among participants aged 31 – 40 years and 41 – 50 years while participants aged 21 – 30 years were 63% 0.37 [95%CI = 0.20 – 0.68,  $P = 0.001$ ] less likely to indicate that the utilization of HBCP was good/very good/excellent compared to participants aged 21 – 30 years., Likewise, participants aged 41 – 50 years were at 66% 0.34 [95%CI = 0.15 – 0.77,  $P = 0.010$ ] less likely to indicate that utilization of HBCP was good/very good/excellent compared to participants aged 21 – 30 years. The findings shows that there were equally as many female (49.4%) as there were male (50.6%) respondents included in the study. The results agrees with Muthuri et al. [26] who found that 61.3% of healthcare workers in both public and private sector in Meru County were female, although both genders are well represented in employment in the public hospitals.

#### 3.2 Extent to Which Home Based Care Program Supports Hospital based Care

The study aim was to evaluate the extent to which Home Based Care Programs services supports Hospital-Based Care for Children aged between 1-14 years diagnosed with selected Chronic and Terminal illnesses. Consequently, respondents were asked questions to gauge the extent to which Home Based Care Program supports Hospital-Based Care. Likert scale was used to score the extent to which various aspects of Home-based Care Program supports Hospital Based Care, ranging from 1-5, representing not at all, very little, to some extent, to a good extent and to a great extent.

#### 3.3 Extent of Clinical and Minimum Package Support to Hospital Based Care

Participants exhibited varied responses as to what extent Clinical services support hospital-based care, with majority (38.5%) stating that this aspect of Home-based care supported to a very little extent, 21.5%, reported that it supported to some extent, while 29.6% felt that it was supportive to a good and/or great extent. Results presented in Table 1 further indicates that Clinical services offered in home-based care did not sufficiently support Hospital care, with a mean of approximately 3 (mean=2.86) depicting little support. Pooja et al. [27] stated that there are a number of challenges that deter clinicians and other health care workers from participating in home-based care, including home visits and management of patient needs at home. This could hint on why the Home-Based Care

**Table 1. Extent to which clinical services provision at home supports hospital care**

S/No.	Likert Items	1F(%)	2F(%)	3F(%)	4F(%)	5F(%)	Mean
1.	Clinical home visits services	11(8.1)	56(41.5)	27(20)	27(20)	14(10.4)	2.86
2.	Home treatment	14(10.4)	47(34.8)	39(28.9)	18(13.3)	17(12.6)	
3.	Prescription at home	19(14.1)	41(30.4)	31(23)	23(17)	21(15.6)	
4.	Pain management and analgesia services at home	15(11.1)	46(34.1)	34(25.2)	23(17)	17(12.6)	
5.	Physical Assessment	17(12.6)	45(33.3)	25(18.5)	31(23)	17(12.6)	
	Extent to which Clinical services provision at home Supports Hospital Based Care score	14(10.4)	52(38.5)	29(21.5)	23(17)	17(12.6)	

Key: Frequency Number (1F, 2F, 3F, 4F and 5F)

Program seems not to be able to provide support to hospital-based care, especially due to poor staffing and weak referral system, according to Kenya Health workforce report and Sips et al., [14,28].

**3.4 Early Screening and Diagnosis of Chronic and Terminal Illnesses**

Findings on early screening and diagnosis indicate that efforts are being made to conduct screening for chronic and terminal illnesses among Children, as majority (36.8%) agreed that screening was conducted to a good extent. It however needs improvement as close to a third (25.6%) felt that very little screening was being conducted while 9.6% felt that it was not being done at all, as illustrated in Table 2. Further, it is evident that early diagnosis of these illnesses contributed substantially in support of hospital-based care, with 48.0% stating a good and/or great extent of support. Fragala et al., [29,30,31] noted that early screening and diagnosis enables early detection and prompt treatment of chronic diseases, preventing disease progression. As evident in these findings, the County's health sector is making strides towards this, with majority of the interviewed participants finding it supportive towards hospital care.

**3.4.1 The level of home based care for children with selected chronic and terminal illness in relation to extent of HBCP support to hospital based care**

Two out of six factors on extent of Support to Hospital Based Care were significantly associated with the level of Home Based Care for Children with selected chronic and terminal illness, P<0.05.

The extent to which Clinical services are provided at home supports hospital care in table

3 above was significantly associated with level of home based care. High proportion of participants indicating that the level of home based care was good/very good/ excellent was observed among participants who revealed that Clinical services provision at home to support hospital care was somewhat/available/to great extent 54 (78.3%) done compared to participants who revealed that Clinical services provision at home to support Hospital Based Care was not at all/very little 25 (37.9%) done. Participants who revealed that Clinical services provision at home to support Hospital Based Care was somewhat/available/to great extent done were 5.90 [95%ci = 2.77 – 12.60, p < 0.001] times more likely indicate that the level of home based care was good/very good/excellent compared to participants who revealed that it was not at all/very little done.

**3.5 Stepwise Multivariate Logistic Regression to Determine the Level of Home Based Care for Children with selected Chronic and Terminal illness**

Stepwise multivariable logistic regression was fitted on all significant (P<0.05) factors at bivariate analyses to model factors determining the level of Home Based Care for Children with selected chronic and terminal illness. Backward conditional method was specified and variable removal threshold set at P < 0.05. Four factors were retained in the reduced model after nine iterations were run. Participants who had worked in the health facility for a period of 1 – 3 years were 5.29 [95%CI = 1.45 – 19.34, P = 0.012] times more likely to indicate good/very good/ excellent level of Home Based Care compared to participants who had worked for 7 or more years. Participants who revealed that Clinical services for Children with selected CI/TI were occasionally/frequently/very frequently provided were 5.17 [95%CI = 1.81 – 14.82, P = 0.002]

**Table 2. Early screening and diagnosis of chronic and terminal illnesses**

S/No.	Likert Items	1F(%)	2F(%)	3F(%)	4F(%)	5F(%)	Mean
1.	Screening of Children for early detection of chronic and terminal is done	12(9.6)	32(25.6)	27(21.6)	46(36.8)	8 (6.4)	3.14
2.	Early diagnosis of chronic and terminal illness in Children Supports hospital care is done	13(10.4)	20 (16)	32(25.6)	45 (36)	15 (12)	
	Early screening and diagnosis of chronic and terminal illnesses score	9 (7.2)	21(16.8)	32(25.6)	44(35.2)	19(15.)	

Key: Frequency Number (1F, 2F, 3F, 4F and 5F)

**Table 3. Home based care for children with selected chronic and terminal illness in relation to extent of clinical services**

Variables	Good/very good/excellent		Poor/fair		OR	95%CI		P-Value
	N	%	N	%		L	U	
Extent to which Clinical services Provision at Home Supports Hospital care								
Not at all/Very little	25	37.9%	41	62.1%	Ref			
Somewhat/Available/To great extent	54	78.3%	15	21.7%	5.90	2.77	12.60	<0.001

Legend: Ref = Reference, OR =Odds Ratio, CI confidence Interval

**Table 4. Reduced Model of factors Used to Determining the Level of Home Based Care for Children with selected Chronic and Terminal illness**

Variables	AOR	95% C.I		P-Value
		Lower	Upper	
<b>Years have you worked in this Health facility</b>				
1 – 3 years	5.29	1.45	19.34	0.012
4 – 6 years	2.03	0.56	7.31	0.278
7 years or more	Ref			
<b>Provision of Clinical services for Children with selected CI/TI</b>				
Never/Very rarely/Rarely	Ref			
Occasionally/Frequently/Very frequently	5.17	1.81	14.82	0.002
<b>Health education and promotion to Children with selected CI/TI</b>				
Never/Very rarely/Rarely	Ref			
Occasionally/Frequently/Very frequently	5.14	1.81	14.56	0.002
<b>Extent to which Clinical services provision at home Supports Hospital care</b>				
Not at all/Very little	Ref			
Somewhat/Available/To great extent	3.15	1.13	8.75	0.028

Legend: Ref = Reference, AOR =Adjusted Odds Ratio, CI confidence Interval

times more likely to indicate good/very good/ excellent levels of Home Based Care compared to participants who revealed that clinical services for Children with selected CI/TI were never/very rarely/rarely provided. Participants who revealed that health education and promotion to Children with selected CI/TI occasionally/frequently/very frequently existed were 5.14 [95%CI = 1.81 – 14.56, P = 0.002] times more likely to indicate good/very good/ excellent levels of Home Based Care compared to participants who revealed that health education and promotion to Children with selected CI/TI never/very rarely/rarely existed. Participants who revealed that clinical services provision at home to support hospital care was somewhat/available/to great extent done were 3.15 [95%CI = 1.13 – 8.75, P =0.028] times more likely indicate that the level of Home Based Care was good/very good/excellent compared to participants who revealed that clinical services provision at home supports Hospital Based Care was not at all/very little done as shown in Table 4.

#### 4. DISCUSSION

The study involved 245 Health Care Providers and Caregivers of Children diagnosed with selected Chronic and Terminal illnesses and it sought to determine the extent to which Home Based Care Programme Supports Hospital Based Care provide HBCP services for Children aged 1-14 years diagnosed with selected Chronic and or Terminal illnesses by evaluating Essential Home Based Care Program minimum package from selected study Health Care Facilities of Meru County [18].

The 245 Health Care Workers were sampled from one Level 5 Teaching and Referral Hospital and five sub county hospitals with mean age of 34.4± 9.06 (SD). Majority (46.1%) of the study participants were aged between 21 – 30 years while a small percentage (8.2%) were more than 50 years, this agrees with studies conducted elsewhere in the world revealed that Public Health interventions requires skilled health care workers [32,33].

#### **4.1 Extent of Home Based Care Program Services Support to Hospital Based Care**

About 50.4% of the participant's revealed that utilization of minimum HBCP essential package by caregivers in management of CI/TI complications support Hospital Based Care hence reducing the burden of Chronic and Terminal illness, this agrees with study conducted in US [12,34,35,31]. Likewise, about 50.4% of the respondents indicated that efficient outreach for immunization of Children at home settings, horizontal Referral, vertical Referral, linkages and networks of HBCP centers [36], development and linkage of community based HBCP centers were some of the available effective HBCP approaches and strategies supporting Hospital Based Care for Children diagnosed with selected Chronic and Terminal Illness. This agrees with a studies by Brust et al., [37,28], Brust et al. [38], and Goodson et al. [39] who concluded that harmonization and integration of HBCP programme into mainstream health care system reduces the burden of disease in Health Care Facilities. Majority of the participants revealed that provision of clinical services at home supports very little 52(38.5%) of hospital care.

However majority of the participants indicated that early screening and diagnosis of Chronic and Terminal Illnesses is available 44(35.2%); to support hospital based care. Likewise, a large number of participants indicated that HBCP services by caregivers was somewhat 35(28%) available 34(27.2%) supports hospital care which agrees with a study by Gans et al. [12,40].

#### **4.2 Composite Scores and Extent to which HBCP Services Supports Hospital Based Care**

Majorly, respondents rated the extent to which Clinical services, Early screening and diagnosis of Chronic and Terminal Illnesses, HBCP services by caregivers, Minimum HBCP essential package, Effective referral system and linkages and Health education and Health promotion as 5.3% not at all, 19.1% very little, 32.9% somewhat, 33.6% available and 9.2% to great extent supports hospital based care. These agrees with a study conducted by Ministry of Kenya [41,42].

#### **5. CONCLUSION**

There is need for an urgent need for adoption of a holistic approach focusing on health care system strengthening, by putting in place a well-integrated Home Based Care Program minimum HBCP essential package services being adopted into mainstream Health Care System. In addition there is need for adoption of better management practices, planning strategies, resource allocation and monitoring for effective HBCP programmes as per the current guidelines to bridge the gaps identified in planning, management practices, to fill the disjointed policy implementation and resource inadequacies in implementation of HBCP to enhance service utilization to support hospital based care for ultimate delivery of quality care among the chronically and terminally ill Children. The extent to which Clinical services are provided at home supports hospital care in Table 3 above was significantly associated with level of home based care at 5.90 [95%ci = 2.77 – 12.60,  $p < 0.001$ ] times more likely indicate that the level of home based care was good/very good/excellent compared to participants who revealed that it was not at all/very little done.

#### **CONSENT**

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

#### **ETHICAL APPROVAL**

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

#### **ACKNOWLEDGEMENTS**

My sincere gratitude goes to my supervisors Prof. Alfred M. Mutema and Prof. Robert M. Kei for their guidance and invaluable inputs into this study, also more appreciation go to Prof. Eric M. Muchiri Dean School of Health Sciences and Ruth Gibendi University librarian for their support during the initial stages of my research. Further, I wish to acknowledge my colleagues pursuing doctoral programme in Meru University of Science and Technology for their encouragement and motivation even when times were difficulty.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.




## REFERENCES

1. Connelly M, Schanberg LE, Ardoin S, Blakley M, Carrasco R, Chira P, Hayward K, Ibarra M, Kimura Y, Kingsbury DJ, Klein-Gitelman MS, Lawson E, Stinson J. multisite randomized clinical trial evaluating an online self-management program for adolescents with juvenile idiopathic arthritis. *J Pediatr Psychol.* 2019; 44(3):363-374.  
DOI:10.1093/jpepsy/jsy066.PMID: 30204919 Free PMC article. *Clinical Trial.*
2. Lopez AD, Mathers CD. Measuring the global burden of disease and epidemiological transitions: 2002-2030. *Annals of Tropical Medicine and Parasitology.* 2006;100(5-6): 481-499.  
Available:https://doi.org/10.1179/136485906X97417
3. Rahman M, Jhohura FT, Mistry SK, Chowdhury TR, Ishaque T, Shah R, Afsana K. Assessing community based improved maternal neonatal child survival (IMNCS) program in rural Bangladesh. *PLoS ONE;* 2015.  
Available:https://doi.org/10.1371/journal.pone.0136898
4. Ama de-Graft A, Nigel U, Agyemang C, Pascale A, Catherine C, Arhinful D. Tackling Africa's chronic disease burden: From the local to the global; 2010.
5. Alwan A. Global status report on non-communicable diseases. Book: Global status report on noncommunicable diseases. 2011;176.
6. Black RE, Morris SS, Bryce J. Where and why are 10 million Children dying every year? *Lancet.* 2013;361:2226-34.
7. Emami SG. Public health law and Non-communicable diseases. London: UK Health Forum; 2013.
8. Chong PH, De Castro Molina JA, Teo K, Tan WS. Paediatric palliative care improves patient outcomes and reduces healthcare costs: evaluation of a home-based program *BMC. Journal, Palliative Care.* 2018;17:11.  
DOI: 10.1186/s12904-017-0267-z
9. Kreicbergs U. Why and where do Children die? First published in *Acta Paediatrica Journal;* 2018.  
Available:https://doi.org/10.1111/apa.14389
10. US. Census Bureau. Home Care Workers: Key Facts. (New York: Paraprofessional Healthcare Institute); 2016.  
Available:https://national.org/sites/phinational.org/files/phi-home-care-workers-key-facts.pdf  
Accessed on 07/02/2018.
11. De-Graft, Aikins, Ama, Unwin, Nigel, Agyemang, Charles, Allotey, Pascale, Campbell, Catherine, Arhinful, Daniel. Tackling Africa's chronic disease burden: from the local to the global. *Journal of Globalization and health.* 2010;6(5). ISSN 1744-8603.
12. Gans D, Hadler MW, Chen X, Wu S, Diamante A, Roby DH, Kominski GF. California pediatric palliative care (PPC) waiver: Evaluation report. Los Angeles, CA: UCLA Center for Health Policy Research; 2014.
13. Kenya Master Health Facility List; 2013.  
Available:http://kmhfl.health.go.ke/#/gis
14. Kenya Health Workforce Report. The Status of Healthcare Professionals in Kenya; 2019.
15. Layzer JI, Goodson BD. Care in the home: A description of family Child care and the experiences of the families and Children who use it. National Study of Child Care for Low-Income Families, Wave I Report. Cambridge, MA: Abt Associates; 2010.
16. Schatz E, Seeley J, Zalwango F. Intergenerational care for and by Children: Examining psychological dimensions of chronic disease in Africa reciprocity through focus group interviews with older adults in rural Uganda. *Demographic Research.* 2018;38:2003-2026.
17. Feudtner C, Feinstein JA, Zhong W, Hall M, Dai D. Pediatric complex chronic conditions classification system version 2: updated for ICD-10 and complex medical technology dependence and transplantation. Philadelphia, PA 10194, USA; 2014.
18. KEPHCA.. 3<sup>rd</sup> Kenya National Palliative Care Conference 2012; 2013.
19. Bainbridge D, Seow H, Sussman J. Common components of efficacious in-home end-of-life care programs: A review of systematic reviews. *Journal of the American Geriatrics Society.* 2016;64(3): 632-639.

20. World Health Organization. Culture matters: Using a cultural contexts of health approach to enhance policy-making. 2017;1.  
Available:[https://www.euro.who.int/data/assets/pdf\\_file/0009/334269/14780\\_World-Health-Organisation-Context-of-Health-TEXT-AW-WEB.pdf](https://www.euro.who.int/data/assets/pdf_file/0009/334269/14780_World-Health-Organisation-Context-of-Health-TEXT-AW-WEB.pdf)
21. WHO. The African health monitor special issue: Universal health coverage. 2015;20.
22. WHO. Standards for improving the quality of care for Children and young adolescents in health facilities; 2018. ISBN 978-92-4-156555-4.  
Available:[http://www.who.int/maternal\\_child\\_adolescent/documents/quality-standards-child-adolescent/en/](http://www.who.int/maternal_child_adolescent/documents/quality-standards-child-adolescent/en/)
23. MoH, Ministry of health republic of Kenya., Trainees Manual Trainees Manual 2013 National Palliative Care Training Curriculum For HIV & AIDS, Cancer and other Life Threatening Illnesses. (N.D.); 2013.  
Available:[Www.Kehpca.Org](http://www.kehpca.org).
24. Kreicbergs U. Paediatric palliative care: A personal perspective April 2021, Acta Paediatrica Journal; 2021.  
DOI: 10.1111/apa.15897, License CC BY-NC 4.0
25. Yamane Taro. Statistics: an introductory analysis. New York: Harper & Row; 1973.
26. Muthuri K, Senkubuge F, Hongoro C. A cross sectional study on the pursuit of happiness among healthcare workers in the context of health systems strengthening: The case of Meru County, Kenya. F1000 research. 2021; 10(163).  
Available:<https://doi.org/10.12688/f1000research.51203.1>
27. Pooja D, Gunukula A, Gupta N, Adams DJ, Kulhari H. Bombesin receptors as potential targets for anticancer drug delivery and imaging. The international journal of biochemistry & cell biology. 2019;114:105567.
28. Sips I, Mazanderani AH, Schneider H, Greeff M, Barten F, Moshabela M. Community care workers, poor referral networks and consumption of personal resources in rural South Africa. PloS one. 2014;9(4).
29. Fragala S, Shiffman D, Birse C. Population health screenings for the prevention of chronic disease progression. American journal of managed care. 2019;25(11):548-553.
30. Forry ND, Iruka I, Kainz K, Tout K, Torquati J, Susman-Stillman A, Bryant D, Starr R, Smith S. Identifying Profiles of Quality in Home-Based Child Care, Issue Brief OPRE 2012-20. Washington, DC: Office of Planning, Research and Evaluation, Administration for Children and Families, U.S. Department of Health and Human Services; 2012.
31. Revill P, Ryan P, McNamara A, Normand A. A cost and outcomes analysis of alternative models of care for young Children with severe disabilities in Ireland Analyse, 1875-0672/\$ – see front matter © 2013 Association ALTER. Published by Elsevier Masson SAS. All rights reserved; 2013.  
Available:<http://dx.doi.org/10.1016/j.alter.2013.04.002>
32. Love and Fox. Home-based parent child therapy for young traumatized children living in poverty: A Randomized Controlled Trial; 2017.  
DOI: ISBN 4142881469
33. MOH. E-health Kenya facilities; Master facility list as at June 2013; 2010.  
Available:<http://ehealth.or.ke/facilitylogin>;  
<http://kmhfl.health.go.ke/#/gis>
34. NIA (National Institute on Aging). Global health and aging. Washington, DC: U.S. Department of Health and Human Services. NQF (National Quality Forum). Palliative and end-of-life care 2015-2016: Technical report. 2016. Washington DC: National Quality Forum; 2011.
35. CBS (Kenya), MoH (Kenya) and ORC Macro, Kenya Demographic Health Survey 2014. Calverton, Maryland: CBS, MoH, and ORC Macro; 2014.
36. Bloom DE, Cafiero-Fonseca ET, Candeias V, Adashi E, Bloom L, Gurfein L, et al. Economics of non-communicable diseases in India: The costs and returns on investment of interventions to promote healthy living and prevent, treat, and manage NCDs. World Economic Forum, Harvard School of Public Health; 2014.
37. Brust JC, Shah NS, Scott M, Chaiyachati K, Lygizos M, van der Merwe TL, Margot B. Integrated, home-based treatment for MDR-TB and HIV in rural South Africa: an alternate model of care [Perspectives]. The international journal of tuberculosis and lung disease. 2012;16(8):998-1004.
38. Brust JCM, Shah NS, Mlisana K, Moodley P, Allana S, Campbell A, Johnson BA, Master I, Thuli Thuli M, Lachman S, Larkan

- L, Ning Y, Malik A, Smith JP, Gandhi NR. Improved survival and cure rates with concurrent treatment for multidrug-resistant tuberculosis–human immunodeficiency virus coinfection in South Africa. *Clinical Infectious Diseases*. 2018; 66(8):1246–1253.  
Available:<https://doi.org/10.1093/cid/cix1125>
39. Gitonga Z, Keiyoro P. Factors influencing the implementation of healthcare projects: The case of Meru County, Kenya. *International academic journal of information sciences and project management*. 2017;2(1):259-280.
40. Young, Camilla H, Amanda M, Gonçalves B, Daniela CQ, Terry JH, Lotty van M, Barbara CS, David J. Home or foster home care versus institutional long-term care for functionally dependent persons. *Cochrane Database of Systematic Reviews*; 2017.
41. MOH. National palliative, care guidelines, MoH Kenya; 2013.
42. Philip RR, Philip S, Tripathy JP, Manima A, Venables E. Twenty years of home-based palliative care in Malappuram, Kerala, India: a descriptive study of patients and their care-givers. *BMC Palliative Care*. 2018;17(1):26.

## APPENDIX 1. Ethical approval MIRERC



# MERU UNIVERSITY OF SCIENCE & TECHNOLOGY

P.O Box, 972-60200 Meru-Kenya

Phone: +254 712524293, 254 725330826, +254 705790660, +254 2092048, +254 706364687 +254 6430320

Email: info@must.ac.ke Website: www.must.ac.ke

### RESEARCH DIRECTORATE INTERNAL MEMO

From: Dr Elijah Walubuka,  
Chairperson, MIRERC Ref: MIRERC/037/2017

Through: Professor Robert G. Kei  
Dean, School of Health Sciences

To: Mr Patrick Kinyua Kubai Date: 4<sup>th</sup> December, 2017  
Principal Investigator

**SUBJECT: MIRERC-037-2017: Evaluation of homebased care programme for children between 1-14 years with selected chronic and terminal illness in Meru County.**

The Meru University of Science and Technology Institutional Research Ethics Review Committee (MIRERC) acknowledges receipt of your protocol hand-delivered by the Dean of the School of Health Sciences on 4<sup>th</sup> December 2017.

MIRERC has thoroughly reviewed the protocol. The Committee is satisfied that you have adequately all the ethical considerations in subsection 3.10; and appended a well formulated Informed Consent Document (ICD) in Appendix 1 (page 78-84). Thus, it recommended to approve the proposal with no amendments.

Therefore, the study is granted approval for implementation effective this day, 5<sup>th</sup> December 2017 for a period of one year. Please note that authorization to conduct this study will automatically expire on 5<sup>th</sup> December 2018. If you plan to continue data collection or analysis beyond this date, please submit an application for continuation approval to MIRERC by 30<sup>th</sup> October 2018.

You are required to submit any proposed changes to this study to MIRERC for review and the changes should not be initiated until written approval is received. Please note that any unanticipated problems resulting from the implementation of this study should be brought to the attention of MIRERC and you should advise MIRERC when the study is completed or discontinued. Please use the relevant MIRERC forms available at the MIRERC Office.

You may now embark on your study.

Yours faithfully,


Dr Elijah Walubuka,

Chairman, MIRERC



MUST IS ISO 9001:2015 CERTIFIED

## APPENDIX 2. MERU County Government approval



© 2021 Kubai et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<http://www.sdiarticle4.com/review-history/69310>