



# Effect of Sub-Lethal Concentrations of *Vernonia amygdalina* (Bitter Leaf) on Testes of *Clarias gariepinus* (African Catfish) Juveniles

Y. Sulaiman <sup>a\*</sup>, Y. M. Sanyinna <sup>b</sup>, T. D. Bala <sup>a</sup>, M. Y. Mohammed <sup>c</sup>  
and S. M. Abdullahi <sup>d</sup>

<sup>a</sup> Department of Zoology, Faculty of Natural and Applied Sciences, Nasarawa State University, Keffi, Nasarawa State, Nigeria.

<sup>b</sup> Department of Animal and Environmental Biology, Faculty of Life Sciences, Kebbi State University of Science and Technology, Aliero, Kebbi State, Nigeria.

<sup>c</sup> Department of Animal Production, Faculty of Agriculture, University of Jos, Plateau State, Nigeria.

<sup>d</sup> Department of Biological Sciences, Faculty of Science, Federal University of Kashere, Gombe State, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJRIZ/2022/v5i230132

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83684>

Received 08 February 2022

Accepted 16 March 2022

Published 28 April 2022

Original Research Article

## ABSTRACT

*Vernonia amygdalina* has recently been linked to insecticidal and pesticidal properties that could replace the harmful agrochemical pesticide usage around the aquatic environment and to aquatic inhibitors such as fish. This study aimed to determine the effect of sub-lethal concentrations of *V. amygdalina* on testes of *Clarias gariepinus* (African catfish) juveniles. The fishes were exposed to 0.00, 0.40, 0.80 and 1.60 g/L graded concentrations of aqueous crude leaves extract of *V. amygdalina* for two months. Oxidative stress biomarkers, histology, histomorphometry, and somatic indices of testes were determined. There was a reduction in activities superoxide dismutase (SOD), malondialdehyde (MDA) and reduced glutathione (GSH) as the concentration of the *V. amygdalina* increased compared with the control. Mild interstitial oedema, mild tubular germ cell, interstitial cell depletions, severe depletion of seminiferous luminal content and sloughing off of the seminiferous tubular boundary tissue were observed testes of catfish exposed to various

\*Corresponding author: Email: [ysulaiman371@gmail.com](mailto:ysulaiman371@gmail.com);

concentrations of *V. amygdalina* compared with the control. There was an increase in testis somatic indices as the plant extract increased compared with the control. Bitter leaf extract was shown to have a toxic effect on *C. gariepinus* juveniles. As such, the pesticide use of the plants near the aquatic environment should be discouraged.

**Keywords:** *Bitter leaf; sub-lethal concentrations; African catfish; oxidative stress biomarkers; histology; somatic indeces.*

## 1. INTRODUCTION

With the world's population estimated to cross 8.2 billion by 2030 and 842 million people estimated to be undernourished in 2011–13, food supply will be an increasing challenge over the next twenty years [1]. With rising income and demographic changes such as family size, population ageing, and urbanisation, as well as consumer trends such as concerns about healthy eating and sustainable production, there will be major changes in demand and significant changes in the composition of demand [1]. This scenario will have an effect on food supply, which will need to enhance and become more effective if it is to grow within the constraints imposed by natural resource availability and existing technology [1]. Fish consumption by people is on the rise around the world, owing to the availability, accessibility, and cost of meats like beef, pork, and poultry [1]. Consequently, some concerns begin to emerge, primarily regarding harvesting fishes from natural water bodies and fish treatment by using piscicides by aquaculturists which may affect the aquatic environment and quality of fish available in the market. The bioactive ingredient could be present in any product of animal origin, causing economic losses and putting human and animal health at risk. According to Bostock et al. [2], aquaculture contributes nearly half of all food of aquatic origin intended for human consumption as a vital part of the global food industry.

The African catfish (*Clarias gariepinus*) is a Clariidae catfish species that is the most widely grown fish in Nigeria. Due to its stress resilience, capacity to survive a wide range of environmental conditions, high stocking density under culture conditions, and relatively fast development, it is the most widely requested freshwater fish on the planet. They live in freshwater lakes, rivers, swamps, and man-made environments such as oxidation ponds and even urban sewage systems throughout Africa and the Middle East. The African catfish can never mentioned when talking about traditional fish capture. The current aquaculture practices, are

similar to tilapia in terms of culture [3]. The *C. gariepinus* is a native of Algeria, Angola, Benin, Botswana, Burkina Faso and an excellent sentinel to study of plant crude extracts. It feeds on detritus. It inhabits the streams, rivers, lakes, estuaries and brackish lagoons, found all year round, and its feeding habits reflect the local contamination simply because it is a bottom dweller.

Botanical extracts have piqued interest in aquaculture for controlling fish parasites, fish fry predators, and unwanted fishes from aquaculture ponds to substitute chemical pesticides and insecticides during the previous several decades. Because the widespread and indiscriminate use of these non-biodegradable synthetic compounds has a negative influence on the aquatic environment and puts non-targeted creatures at danger [4].

Various plants are known for their therapeutic and antibacterial capabilities, including pesticidal, acaricidal, and trypanocidal properties [5, 6]. Depending on the nature and quantities of their bioactive ingredients, some are known as strong arrow and fish poisons [7, 8]. This is due to the fact that plants contain a wide range of biological compounds with different characteristics [9]. Because of their eco-friendliness, ease of availability, high efficiency, rapid biodegradability, and low toxicity to non-targeted animals, plant extracts are regarded potential agents [10]. So far, a large number of plants have been studied in many countries to determine their pesticidal [11, 12] and piscicidal activity [11,12,13,14,15, 16,17,18,19]. However, commercially accessible plant products are still limited, thus new sources of botanical pesticides and piscicides for the constantly expanding pisciculture industry should be sought. Piscicides are plant extracts that have toxicological effects on fish and induce mortality in these aquatic creatures [20]. Plant piscicides are made from a variety of plants from various families and species, which might differ significantly not only in terms of taxonomic differences, but also in terms of plant components used (leaves, barks, fruits, and

seeds), method of application, mode of extraction, and target fish species [21,15]. Botanical materials contain a number of bioactive chemicals that act as piscicides separately or in combination [17]. Any plant extract's toxicity and piscicidal activity can be determined by exposing fish to it and then calculating the median lethal concentration (LC50) [19]. Alteration in haematological, biochemical, and antioxidant parameters of fishes [22, 23], as well as biological and physiological activities of fish, are some of the common toxicological effects of plant extract [22, 23]. In *Clarias gariepinus*, Adeyemo [24] found the haematological and histological impacts of Cassava mill effluent.

Man's activities have been responsible for introducing contaminants into the environment in the last few decades [25]. pH, turbidity, alkalinity, dissolved oxygen, temperature, and conductivity are all environmental parameters that determine the pace at which pollutants enter the water and their fatal effects on aquatic life [26]. Nowadays, contamination of natural ecosystems is on the rise, and it is quickly becoming a major issue as human activities increase. A basic topic in ecotoxicology study is determining toxic substances in aquatic environments and their consequences on aquatic animals. Chemical analysis can reveal the presence of toxic compounds in the environment, but it cannot determine their effects on aquatic animals in aquatic ecosystems. As a result, bioassay experiments are required to assess the environmental impact of harmful substances [27].

Information regarding the adverse effect of sub-lethal concentrations on the reproductive system of *C. gariepinus* is scanty. Healthy testes of fish are important determinants of its breeding potential, and thus any toxicological factor adversely affecting the histopathology of testes will reduce the gross production of fish.

*Vernonia amygdalina* has been reported to be used as growth enhancer, phyto-additive, antibacterial, milt booster and insecticidal by *C. gariepinus* farmers [34]. However recent studies showed that *V. amygdalina* has negative effect on glucose level of *C. gariepinus* [35]. Acute toxicity studies showed that the plant is toxic to catfish [28]. There is scarcity of information on the testis somatic indices, antioxidant biomarkers and histology and histomorphometry of *C. gariepinus* fish exposed to sub-lethal concentrations of *V. amygdalina*. Hence, this study examined the effect of sub-lethal

concentrations of *Vernonia amygdalina* (bitter leaf) on testes of *Clarias gariepinus* (African catfish) juveniles.

## 2. MATERIALS AND METHODS

### 2.1 Description of Study Area

*Vernonia amygdalina* was collected from Keffi Local Government Area of Nasarawa State, Nigeria. This study was carried out at the Department of Zoology Laboratory, Faculty of Natural and Applied Sciences, Nasarawa State University, Keffi.

### 2.2 Procurement and Processing of Experimental Plant (*Vernonia amygdalina*)

The leaves were air-dried at room temperature ( $27 \pm 2^{\circ}\text{C}$ ) in the Laboratory of Zoology Department, Faculty of Natural and Applied Sciences, Nasarawa State University, Keffi. The dried leaves were pounded with laboratory mortar and pestle into powder, sieved with a 30  $\mu\text{m}$  mesh sieve.

### 2.3 Collection, Transportation and Acclimatization of Experimental Animal

*C. gariepinus* was purchased from Rayuwa Fish Farm, Karu Local Government Area, Nasarawa State, Nigeria. Experimental animals were transported in tanks containing water from the pond to the Laboratory Unit of the Zoology Department, Nasarawa State University, Keffi. The fish was transferred into aquaria containing dechlorinated municipal tap water and acclimatized to the laboratory conditions for two weeks. During this period, the fish was fed to satiation at 8:00 a.m. and 6:00 p.m. with commercial fish feed (Vital feed®).

### 2.4 Sub-Lethal Concentrations of Aqueous Crude Leaf Extract of *V. amygdalina* and Test Procedure

Mortality (0, 1, 2, 5, 7 and 10) of *C. gariepinus* juveniles exposed to acute concentrations of 0.0, 0.19, 0.38, 0.75, 1.50 and 3.00 g/L of crude leaf extract of *V. amygdalina* was reported by Audu et al. [36]. The LC<sub>50</sub> was calculated as 0.82 g/L by using the Finney calculator. *V. amygdalina* crude leaves extract administered at sub-lethal concentrations 1/5, 1/10, 1/20 of the LC<sub>50</sub> [37] in a static renewable bioassay system.

The weighed samples were macerated in 1 L of distilled water for 24 hours at room temperature (27.0°C) to obtain the stock solution. The obtained stock solution was filtered through a funnel choked with non-absorbent cotton wool, and each filtrate (1.6, 0.8 and 0.4 g/L) was transferred into tanks A1, B1, and C1, respectively, with D1 as the control. Each of the filtrates was diluted by adding 9 L of dechlorinated municipal tap water. The same procedure was repeated in replicate tanks A2, B2 and C2, while tanks D1 and D2 which were not inoculated with the test materials, served as the controls. The sub-lethal toxicity test lasted for two months (56 days or eight weeks). The test fish was fed 3% body weight at 0800 and 1800hrs with photoperiod being natural.

## 2.5 Experimental Design

The experiments consist of four circular rubber tanks and one hundred and twenty *C. gariepinus* juveniles, mean weight 23.13±2.43 g and mean total length 12.501±0.39 cm in a non-randomized block design. Each of the eight tanks was filled with 9 L of dechlorinated municipal tap water. Six of the filled tanks were inoculated with various concentrations (powder earlier macerated in 1 L of water) of aqueous crude leaves extract of *V. amygdalina* and *C. gariepinus* fingerlings were introduced into each.

## 2.6 Assay for Biomarkers of Oxidative Stress

### 2.6.1 Preparation of the Tissue Samples for the Study

At the end of the experiment, fishes were carefully netted to minimize stress, weighed and sacrificed. Tissues such as liver, testes and gills were carefully removed and weighed. The tissues were washed in chilled phosphate buffer saline. Sample homogenates were made of individual tissues with chilled phosphate buffer (0.1 M, pH 7.4), centrifuged in a centrifuge at 9,000 rpm for 20 min at 4°C [38]. The supernatant obtained was used for further analysis. Samples were transported to Biochemistry Units of National Veterinary Research Institute (NVRI), Vom, Plateau State, Nigeria.

### 2.6.2 Determination of Glutathione Peroxidase

Glutathione peroxidase (GPx) activity was assayed as described by Paglia and Valentine

[39] with modifications according to Lawrence and Burke [40]. The reaction mixture contained 50 mM potassium phosphate buffer (pH 8.3), 1 mM EDTA, 1 mM sodium azide, 0.2 mM nicotinamide adenine dinucleotide phosphate (NADPH), and 1 U/mL glutathione reductase. The reaction was initiated with the addition of 1.5 mM cumene hydroperoxide. The enzyme activity was estimated from the rate of oxidation of NADPH. The reagents were mixed, and the absorbance was measured at 340 nm. Enzyme activity was expressed in mmol/minute/milligram protein.

### 2.6.3 Determination of Superoxide Dismutase Activity

Superoxide dismutase (SOD) activity was assayed as described by Misra and Fridovich [41]. The assay was based on the ability of SOD to inhibit the autoxidation of epinephrine at an alkaline pH. 0.5 ml of tissue homogenate was diluted with 0.5 ml of distilled water, to which 0.25 ml of ice-cold ethanol and 0.15 ml of ice-cold chloroform was added. The mixture was properly mixed using a cyclo mixer for 5 minutes and centrifuged at 2500 rpm. To 0.5 ml of supernatant, 1.5 ml of carbonate buffer (0.05 M, pH 10.2) and 0.5 ml of EDTA solution (0.49 M) were added. The reaction was initiated by the addition of 0.4 ml of epinephrine (3 mM), and the change in optical density/minute was measured at 480 nm against the reagent blank. SOD activity was expressed as units/mg protein. Change in optical density per minute at 50% inhibition of epinephrine to adrenochrome transition by the enzyme was taken as the enzyme unit.

### 2.6.4 Determination of Lipid Peroxidation

The malondialdehyde (MDA) content, a measure of lipid peroxidation, was assayed in the form of thiobarbituric acid reactive substances (TBARS) using the method of Buege and Aust [42]. 1.0 mL of the sample was added to 2 mL of (15% (w/v) trichloroacetic acid, 0.375% (w/v) thiobarbituric acid (TBA), 0.25 M HCl). The solution containing plasma and reagent was heated in a boiling water bath for 15 min and then cooled. Precipitates were removed by centrifuging at 1000 rpm for 10 minutes. The supernatant was removed, and the absorbance read at 535 nm against a blank. MDA was calculated using the molar extinction coefficient  $\epsilon$  for MDA  $1.56 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ . Results were expressed as  $\mu \text{ mol}^{-1}$ .

## 2.7 Extraction of Organs of *C. gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *Vernonia amygdalina*

Following the exposure of *C. gariepinus* juveniles to sub-lethal concentrations of crude leaf extract of *V. amygdalina*, four fishes from treatment and control tanks were removed, sacrificed and dissected to obtain the testis, gill, liver and kidney. Each removed organ was rinsed with distilled water to wash off traces of blood. Each organ was preserved in 10 ml specimen bottle containing 5 ml of formal saline before transportation to NVRI, Vom, Plateau State, Nigeria, for histopathological analyses of the tissues.

### 2.7.1 Determination of Organ Somatic Indices of *C. Gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina*

The calculation of the testes somatic index (TIS) was done by random selection of four individual fishes from each treatment group and total weight of the individuals were noted [30]. The organs of the fishes like testes were removed carefully and weighed in an electronic weight machine, after removing moisture by blotting paper [43].

Organs Somatic Index was calculated using:

$$OSI = \frac{\text{Weight of organ}}{\text{Weight of fish}} \times 100$$

## 2.8 Procedure for Histological Examination of Organs of *C. gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina*

The routine Paraffin wax method and haematoxylin-eosin staining techniques of tissue processing described by Drury *et al.* [44] and Avwioro [45] were adopted to examine the gills, kidneys and liver of *O. niloticus* fingerlings exposed to aqueous crude leaves extract of *B. aegyptiaca*. The harvested gills, kidneys and liver of *C. gariepinus* were fixed in 10% formalin for three days, cut into thin slices of 5 x 2 x 1 mm thick and then processed with the SPIN Tissue Processor (STP) 120 (Thermo Scientific Model). The tissues were buffered in 10% formalin before passing through the following levels of

hydrocarbons for two hours each: 70% Alcohol, 80% Alcohol, 90% Alcohol, 95% Alcohol, Absolute Alcohol I, Absolute Alcohol II, Absolute Alcohol III, Xylene I, Xylene II, Paraffin Wax Oven I, Paraffin Wax Oven II.

Tissues were embedded in molten paraffin wax using embedding moulds. The tissues were embedded using cassettes on a tissue Tek Embedding Centre (SLEE MPS/P2) and cooled rapidly on the cooling component. Tissues were sectioned using a rotary microtome (MICROM HM340E Thermo Scientific) set at four microns, picked on slides and ready for staining.

Haematoxylin and eosin staining technique was used for the staining of the tissues. Tissue sections were dewaxed and hydrated by passing through two changes of xylene and through descending levels of alcohol (100%, 80%, 70%) for three min each and then into the water before staining in Harris' haematoxylin solution for 5 min and washed in running water. They were differentiated in 1% Acid alcohol and then washed thoroughly in water, blued in Scott's tap water substitute for 5 min and rinsed briefly in distilled water. Each tissue was then counterstained in 1% aqueous eosin for 2 min and then washed in water, dehydrated in descending grades of alcohol before clearing in xylene and mounted in Distrene, Plasticizer and Xylene (DPX). Sections were then placed in slide carriers and placed in a 40°C oven to dry overnight. Each tissue was read microscopically. Photographs of the prepared sections were taken using a mounted photo-microscopic camera.

## 2.9 Histomorphometrics of Organs of *Clarias gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *Vernonia amygdalina*

Ten fishes from each experimental group were randomly selected. In addition, hepatocyte histomorphometric measurements such as hepatocyte nuclear diameter and hepatocytes surface area were determined by using Motic image plus 2.0 (Motic Asia, Hong Kong) software [28].

## 2.10 Statistical Analysis Method

Data collected were subjected to analysis of variance (ANOVA) using SPSS version 16. Significant means were set at  $p < 0.05$  and separated using Duncan new multiple range test.

Results were presented as means  $\pm$  standard errors of the mean.

### 3. RESULTS

There is a reduction in SOD activity as the concentration of the toxicant increases. The highest and lowest value of  $19.4462 \pm 0.98461$  and  $18.1615 \pm 1.10000$  were recorded in control (0.00 g/L) and highest concentration (1.60 g/L). There is also a reduction in GSH activity as the concentration of the toxicant increases, the highest and lowest values of  $10.2500 \pm 9.1250$  were recorded in 0.80 g/L and the highest value recorded in 0.40g/L. There is also reduction in GPx activity as the concentration of the toxicant increases, the highest and lowest values of  $2.9746 \pm 1.9046$  were recorded in 0.40g/L and the highest value recorded in 0.80g/L (Table 1).

There is mild interstitial oedema, mild tubular germ cell, interstitial cell depletions, severe depletion of seminiferous luminal content and

sloughing off of the seminiferous tubular boundary tissue observed in testes of the exposed fishes as seen in Plate A-D.

Morphometric parameters such as STD with the highest and lowest value of  $133.79 \pm 9.799$  and  $87.35 \pm 6.211$  shows a significant difference compared to the control recorded in 0.40 g/L and 0.80 g/L. SLD with the highest and lowest value of  $100.45 \pm 7.561$  and  $76.70 \pm 9.241$  shows the significant difference when compared to the control recorded in 0.40 g/L and 0.80 g/L, while SED with the highest and lowest value of  $16.72 \pm 0.740$  and  $9.82 \pm 0.563$  shows significant difference when compared to the control recorded in 0.00 g/L and 1.60 g/L (Table 2).

TSI shows no significant difference compared with the control with the highest and lowest value of  $0.7475 \pm 0.14014$  and  $0.4675 \pm 0.15638$  recorded in the highest concentration (1.60g/L) and (0.80 g/L) (Table 3).

**Table 1. Mean Testes Superoxidase Dismutase, Lipid Peroxidation and Glutathione Peroxidase of *Clarias gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***

Concentrations (g/L)	SOD	GSH	GPx
0.00	$19.4462 \pm 0.98461$	$9.2500 \pm 0.50000$	$2.153 \pm 0.34223$
0.40	$18.2000 \pm 0.41538$	$10.2500 \pm 0.50000$	$1.9046 \pm 0.21081$
0.80	$18.4615 \pm 0.75385$	$9.1250 \pm 0.37500$	$2.9746 \pm 0.64136$
1.60	$18.1615 \pm 1.10000$	$9.5000 \pm 0.75000$	$2.3272 \pm 0.54893$
P-value (0.05)	0.7050	0.5370	0.4940

Key: SOD = Superoxidase Dismutase; GSH = Glutathione Peroxidase; GPx = Lipid Peroxidation

**Table 2. Sub-Lethal Effects of Bitter Leaves Crude Extract on Testes Morphometry Parameters**


Concentrations (g/L)	STD	SLD	SED
0.00	$88.49 \pm 3.174^A$	$84.07 \pm 4.902$	$16.72 \pm 0.740^B$
0.40	$133.79 \pm 9.799^B$	$100.45 \pm 7.561$	$15.05 \pm 0.978^B$
0.08	$87.35 \pm 6.211^A$	$76.70 \pm 9.241$	$15.19 \pm 0.701^B$
1.60	$129.24 \pm 13.068^B$	$99.23 \pm 11.098$	$9.82 \pm 0.563^A$

Key: Values with different letters in the same row are significantly different ( $P < 0.05$ ) compared with the control STD = Seminiferous Tubular Diameter; SLD = Seminiferous Luminal Diameter; SED = Seminiferous Epithelial Diameter

**Table 3. Mean Testes Somatic Index of *Clarias gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***


Concentrations	TSI
1.60	$0.7475 \pm 0.14014$
0.80	$0.4675 \pm 0.15638$
0.40	$0.5100 \pm 0.10141$
0.00	$0.5200 \pm 0.12076$
Total	$0.5613 \pm 0.06515$

Key: TSI = Testes Somatic Index




**Plate A-D: Photomicrographs of the Testes of *Clarias gariepinus* Exposed to Graded Concentrations of *Vernonia amygdalina***


**Key:** A. Control (0.00 g/L): The testes present normal numerous tubular structures housing different spermatogenic cell series (black arrow) and distinct interstitium between the tubules (white arrow). B. 0.4 g/L: There was no visible lesion except for mild interstitial oedema (white arrow). C. 0.8 g/L: Presence of mild tubular germ cell (black arrow) and interstitial cell depletion (white arrow). D. 1.6 g/L: Severe depletion of seminiferous luminal content (star) and sloughing off of the seminiferous tubular boundary tissue (black arrow). Magnification: x400; Stain: Haematoxylin-eosin




**Fig. 1. Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***



**Fig. 2. Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***



**Fig. 3. Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***



**Fig. 4. Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***



#### 4. DISCUSSION

Variation in testes antioxidant biomarkers (MDA, GSH, SOD) indicated oxidative stress in testes which is commensurate with the finding of Burak et al. [30] in the testes of fish from the Lake Van and Karasu River of Van fish *Alburnus tarichi*, who reported that there was an unchanged MDA, GSH and SOD level. This research finding is also in line with a report from Alaa and Rania [31], who opined that there was a significant increase ( $p < 0.05$ ) in activities of serum SOD, glutathione s-transferase (GSH), and malondialdehyde (MDA) when adult *C. gariepinus* were exposed to pure 100 µg/L 4-nonylphenol and quince (*Cydonia oblonga*) to fishes exposed for 15 days when compared to the control, but in this research, CAT was not recorded.

The severe sloughing off of the seminiferous tubular boundary tissue and severe depletion of seminiferous luminal content reported in this research work is in line with findings from Priya and Balu [32] who reported in their work that there was severe deterioration of the testis histology by the 30<sup>th</sup> day of exposure, considerable reduction in spermatozoa in the testis when freshwater fish (*Rasbosa dandia*) was exposed to mercury. However, they reported a congestion of blood vessels and proliferation of interstitial tissues and vacant spaces enlarged which was not seen in this research work. The result shows that there is a significant difference in the testis somatic indices of *C. gariepinus* concerning the graded concentrations of *V. amygdalina* when compared with the control, and this research work is not in line with the finding of Ada et al. [3], who reported that there were no changes in gonad somatic indices in *C. gariepinus* exposed to Atrazine and coconut water after the fishes were exposed for a period of 14 days. This research work is in line with the findings of Claramma and Radhakrishnan [33], which reported that there were histopathological changes observed as a distortion of seminiferous tubules, disorganization of spermatogonia, spermatocytes and spermatids with cytoplasmic vacuolization and nuclear pycnosis after exposing *Clarias batrachus* to sub-lethal concentrations of chromium for 45 days. Kumar et al. [46] also reported distortion of the seminiferous tubules in the testis of freshwater fish *Oreochromis mossambicus* in response to plant nutrients which is in line with the findings of this research work. However, he also recorded an increase in vacuolization, condensation of

spermatocytes besides inflammation and inter-tubular vacuolation, which was very prominent.

There is an increment in the value of testis somatic index as the concentration of the plant increases which is not in line with the findings of Muazu et al. [47] that reported decline in the value of TSI of the fish with increase in the concentrations. Abdel-Hameid [48] also reported that there was a serious reduction in the GSI of blue tilapia juveniles as concentration levels of phenol increases with no significant increase reported for male fish subjected to 20% 96-h LC<sub>50</sub> of phenol, whereas significant reduction was observed in the case of male fish subjected to 40% and 80% 96-h LC<sub>50</sub> of phenol. Findings from this research work are in line with findings from the work of Shalaka and Pragna [29], who reported that there was an increase in the GSI value when freshwater fish *Oreochromis mossambicus* were increasingly fed with plant nutrient.

#### 5. CONCLUSION

Variation in testis antioxidant biomarkers, histological and histomorphometric parameters in testes of the exposed catfish indicated that the bitter leaf is toxic to the fish. Therefore, indiscriminant use of *V. amygdalina* around aquatic environment should be discouraged.

#### 7. DISCLAIMER

The products used for this research are commonly and predominantly used in our research area and country. There is no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for litigation but the advancement of knowledge. Also, the research was not funded by the producing company rather, it was funded by the personal efforts of the authors.

#### 6. RECOMMENDATION

Further toxicity research works on the effect of the plants on the reproductive organs of other commercial fish species should be conducted.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Lem A, Bjorndal T, Lappo A. Economic Analysis of Supply and Demand for Food up to 2030 – Special Focus on Fish and Fishery Products. FAO Fisheries and Aquaculture Circular No. 2014; 1089.
2. Bostock J, Brendan Mc, Randolph R, Kim J, Trevor T, Kai L, David L, Lindsay R, Neil H, Iain G, Richard C. Aquaculture: Global Status and Trends. Philosophical Transactions of the Royal Society Biological Sciences;2010. DOI: 10.1098/rstb.2010.0170
3. Ada F, Ayotunde EO, William K, B. Gonado-hepatosomatic Indexes of *Clarias gariepinus* Sub-adult Exposed to Atrazine, Cocosnucifera Water and Phyllanthusmuelarianus Extract. Journal of Aquaculture Research & Development. 2015;6:11.
4. Kumar A, Lingadurai S, Jain A, Barman NR. *Erythrina variegata* Linn: A Review on Morphology, Phytochemistry, and Pharmacological Aspects. Pharmacognosy Review. 2010; 4(8):147-152.
5. Nascimento GGF, Locatelli J, Freitas PC, Silva GL. Antibacterial Activity of Plant Extracts and Phytochemicals on Antibiotic Resistant Bacteria. Brazilian Journal of Microbiology. 2000;31: 247-256.
6. Adamu HM, Abayeh OJ, Agho MO, Abdullahi AL, Uba A, Dukku HU, Wufem BM. An Ethnobotanical Survey of Bauchi State Herbal Plants and their Antimicrobial Activity. Journal of Ethnopharmacology. 2005;99:1-4.
7. Geidam MA, Dauda E, Hamza HG. Effects of Aqueous Stem-bark Extract of *Momordicabalsamina* Linn on some Serum Enzymes in Normal and Ethanol Fed Rats. Journal of Biological Sciences. 2007;7:397-400.
8. Kamalkishor HN, Kulkarni HN. Fish Stupefying Plants used by the Gond Tribal of Mendha Village of Central India. Indian Journal of Traditional Knowledge. 2009;8:531-534.
9. Istvan U. Semi-Natural Products and Related Substances as Alleged Botanical Pesticides. Journal of Pest Management Sciences. 2000;56:703-705.
10. Yunis JA, Gustavo SC, Paulo FM, Cynthia I, George GO, Silas FE, Claudine CM, Juliet REM, Flavio RM, Joao BKF. Acute toxicity and determination of the acute constituents of aqueous extract of *uncaria tomentosa* bark in *hyphessobrycon eques*. Journal of Toxicology. 2014; 20(14) 56-87.
11. Rahuman AA, Gopalakrishnan G, Venkatesan P, Geetha K. Larvicidal Activity of Some Euphorbiaceae Plant Extracts against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitology Research. 2008;102:867-873.
12. Dubey SP, Manu L, Manu S. Green Synthesis and Characterizations of Silver and Gold Nanoparticles using Leaf Extract of *Rosa rugosa*. International Journal of Colloids and Surfaces A: Physiochemical and Engineering Aspects. 2013;364:34-41.
13. Kulakkattolickal, A. T. (1989). Piscicidal Plants of Nepal: Toxicity to Air Breathing Predatory fish (*Ophicephaluspunctatus*, *Clariasbartrachus* and *Heteropneustesfossilis*) and duration of Risk to Cultivated fish. Aquaculture. 78:285-292.
14. Chiayvareesajja SI, Chiayvareesajja N. Rittibhonbhun I, Wiriyaichitra P. The toxicity of five native Thai Plants to Aquatic Organisms. Asian Fisheries Science. 1997; 9:261-267.
15. Neuwinger HD. Fish Poisoning Plants of Africa. Botanica Acta. 2004; 107:263-270.
16. Tiwari S, Singh A. Control of Common Freshwater Predatory Fish, *Channapunctatus* through *Neriumindicum* leaf Extracts. Chemosphere. 2004;53: 865-875.
17. Obomanu FG, Ogbalu OK, Gabriel UU, Fekarurhobo SGB, Abadi SU. Piscicidal Effects of *Lepidagathisalopecurooides* on Mudskipper, *Periophthalmus papillio* from the Niger Delta. Research Journal Applied Sciences. 2007;2(4):382-387.
18. Murthy KS, Kiran BR, Venkateshwarlu M. A Review on Toxicity of Pesticides in Fish. International Journal of Open Science Research. 2010;1(1):15-36.
19. Njan AA, Bulus A, Amos GA, Dominic B, Silvia David RB. The Analgesic and Antiplasmodial Activities and Toxicity of *Veronia amygdalina*. Journal of Medicinal Food. 2008;11(3):574-581.
20. Burkill HN. The Useful Plants of West Africa (tropical) Ed. 2 Vol. I. Families A.D. Royal Botanical Garden, Kew; 1995.
21. Van F, Andel T. The Diverse Use of Fish Poison Plants in Northwest Guyana.

- Economic Botany. 2000; 54:500–512.
22. Winkaler EU, Santos TRM, Machado-Neto JG, Martinez CBR. Acute Lethal and Sublethal Effects of Neem Leaf Extract on the Neotropical Freshwater Fish *Prochilodus lineatus*. *Comp. Biochem. Physiol. Toxicol., Pharmacol.* 2013;145(2): 236-244.
  23. Kavitha C, Ramesh M, Kumaran SS, Lakshmi SA. Toxicity of Moringa oleifera Seed Extract on Some Hematological and Biochemical Profiles in a Freshwater Fish, *Cyprinus carpio*. *Exp. Toxicol. Pathol.* 2012; 64(7-8): 681-687.
  24. Adeyemo OK. Haematological and Histopathological Effects of Cassava Mill Effluent in *Clarias gariepinus*. *African Journal of Biomedical Research.* 2005;8:179–183.
  25. Awoyinka OA, Omodara TR, Oladele FC, Ajayi DD. In-vitro Protein Digestibility of Selected Under-utilized Local Wild Beans and Bio-availability in Rats Model. *Academia journal Biotechnology.* 2019; 7(4):063-067.
  26. Fagbenro OA. The Dietary Habits of Clarid Catfish *Heterobranchus bidorsalis* (Geoffery st. Hilaria 1809) in Owerri, Southern Nigeria. *Tropical Zoology.* 2002;5:11–17.
  27. Bagheri F. Study of Pessticial Residue (Diazinon, Azinphosmethyl) in Rivers of Golestan Province (Gorganround and Gharehsou). *Open Journal of Marine.* 2007;7(2):1-125.
  28. Audu BS, Omirinde JO, Gosomji IJ, Wazhi PE. Histopathological Changes in the Gills and Liver of *Clarias gariepinus* Exposed to Acute Concentrations of *Vernonia amygdalina*. *Animal Research International.* 2017;14(1):2576-2587.
  29. Shalaka S, Pragna P. Gonadosomatic and Hepatosomatic Indices of Freshwater Fish *Oreochromismossambicus* in Response to a Plant Nutrient. *Journal of Zoology.* 2013;8(1): 110-118.
  30. Burak K, Ertuğrul K, Abdulahad D, Atilla D. Alterations in Histology and Antioxidant Defense System in the Testis of the Lake Van Fish (*AlburnustarichiGüldenstädt*, 1814). *Environment Monitoring Assessment.* 2016;188:474-547.
  31. Alaa EHS, Rania FKI. Endocrine Disruption, Oxidative Stress, and Testicular Damage Induced by 4-nonylphenol in *Clarias gariepinus*: The Protective Role of *Cydonia oblonga*. *Fish Physiology and Biochemistry.* 2016;10:100-117.
  32. Priya R, Balu TK. Mercury Induced Histopathological Changes in the Testis of the Freshwater Fish, (*Rasbora dandia*). *International Journal of Science and Research.* 2013; 4:2319-7064.
  33. Claramma J, Radhakrishnan MV. Histopathological Analysis of Chromium Toxicity to Testis of the Catfish *Clarias batrachus* (Linn). *IOSR Journal of Environmental Science, Toxicology and Food Technology.* 2016;10: 01-04.
  34. Okukpe KM, Yusuf OA, Adeyemo AD, Omotayo OO, Adeyina AO, Alli IO, Adeyemi-Ale OA, Aderibigbe TA, Opowoye IO, Ogunsola FO, Morakinyo OS. Effects of *Vernonia amygdalina* Leaf Powder on Growth Performance of *Clarias gariepinus*. *Nigerian Journal of Animal Science.* 2018;20(3):180-188.
  35. Alagoa KJ, Osakwe J. Blood Glucose Response of the African Cat Fish (*Clarias gariepinus*) to Bitter Leaf (*Vernonia amygdalina*) Incorporated Diet. *International Journal of Research and Innovation in Applied Science (IJRIAS).* 2021;6(1):60-63. ISSN 2454-6194.
  36. Histopathological Alterations in the Gills and Liver of *Clarias gariepinus* Juveniles Exposed to Acute Concentrations of *Anogeissus leiocarpus*. *Jordan Journal of Biological Sciences.* 2021;14(3):537-543. ISSN 1995-6673. Available:<https://doi.org/10.54319/jjbs/140320>.
  37. Abu-Bakar A, Hu H, Lang MA. *Cyp2a5* Promoter-based Gene Reporter Assay: A Novel Design of Cell-based Bioassay for Toxicity Prediction. *Basic & Clinical Pharmacology & Toxicology.* 2018;123:72-80.
  38. Ogunka-Nnoka C, Assor K, Onuoha S, Amadi P. A Study of the Toxicants and Biomarkers of Oxidative Stress in Samples from Ebubu and Elele-Alimini Communities in Rivers State. *Ovidius University Annals of Chemistry.* 2018;29(1):1-7. DOI: 10.2478/auoc-2018-0001.
  39. Paglia DE, Valentine WN. Studies on the Quantitative and Qualitative Characterization of Erythrocyte Glutathione Peroxidase. *J Lab Clin Med.* 1967;70(1): 158-169. PMID: 6066618.

40. Lawrence RA, Burke RF. Glutathione Peroxidase Activity in Se-deficient Rat Liver. *Biochemical and Biophysical Research Communications*. 1978;71:952-958.
41. Misra HP, Fridovich I. The Role of Superoxide Anion in the Auto-Oxidation of Epinephrine and a Simple Assay for Superoxide Dismutase. *Journal of Biological Chemistry*. 1972;247:3170-3175.
42. Buege JA, Aust SD. [30] Microsomal Lipid Peroxidation. *Methods in Enzymology*. 1978;52:302-310.  
Available:[https://doi.org/10.1016/S0076-6879\(78\)52032-6](https://doi.org/10.1016/S0076-6879(78)52032-6).
43. Sinha P, Annamalai SK, Arunachalam KD. A Wee Study on Behavioural, Organ Somatic Index and Histological Alterations of the Fresh Water Fish *Pangasius sutchi* in Response to Protection Studies, Exposed to Gamma Radiation Perceived by Genotoxic Assays. *International Journal of Pharmaceutical Sciences Review and Research*. 2018;50(2):05:18-30.
44. Drury RAB, Wallington EA, Cameron R. Carleton's Histological Techniques. 4<sup>th</sup> Edn., Oxford University Press, NY. USA, pp: 279-280.
45. Avwioro G. Histological Uses of Haematoxylin – A Review. *JPCS*. 2011;1:24-34.  
Available:[www.arpapress.com/Volumes/JPCS/Vol1/JPCS\\_1\\_05.pdf](http://www.arpapress.com/Volumes/JPCS/Vol1/JPCS_1_05.pdf).
46. Kumar N, Prabhu AJ, Pal AK, Remya S, Aklakur M, Rana RS, Gupta S, Raman RP, Jadhao SB. Anti-Oxidative and Immuno-Hematological Status of Tilapia (*Oreochromis mossambicus*) during Acute Toxicity Test of Endosulfan. *Pesticide Biochemistry and Physiology*. 2011;99:45-52.
47. Muazu TA, Sulaiman MH, Hambolu J. Comparative Anatomy of Wild Adult Tilapia (*Oreochromis niloticus*) Testes during Rainy and Dry Periods in Zaria, Nigeria. *Sokoto Journal of Veterinary Sciences*. 2020;18(3):171-174.
48. Abdel-Hameid NH. Physiological and Histopathological Alterations Induced by Phenol Exposure in *Oreochromis aureus* Juveniles. *Turkish Journal of Fisheries and Aquatic Sciences*. 2007;7:131-138.

© 2022 Sulaiman et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/83684>