



Identification of the Main Dishes Made from the Pulp of Tomi (*Tamarindus indica* L) Consumed in the Savannah Region of Côte d'Ivoire

**Konan Gbè Aya Jacqueline ^a, Nimaga Daouda ^b, Kouassi Kouakou Nestor ^{a,c*},
N'dri Yao Denis ^a and Amani N'Guessan Georges ^a**

^a *Laboratory of Food Biochemistry and Technology of Tropical Products, Department of Food Science and Technology, Nangui Abrogoua University, BP 801, Abidjan 02, Côte d'Ivoire.*

^b *Agronomic, Forestry and Environmental Engineering Department, University of Man, P.O.Box 20, Man, Côte d'Ivoire.*

^c *Swiss Center for Scientific Research in Côte d'Ivoire (CSRS-CI), 01 BP 1303, Abidjan 03, Côte d'Ivoire.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors KGAJ, KKN, NYD and ANGG designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors KGAJ, KKN and NYD managed the analyses of the study. Authors KGAJ and KKN managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2022/v44i1030899

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91141>

Original Research Article

**Received 17 June 2022
Accepted 24 August 2022
Published 31 August 2022**

ABSTRACT

Tamarind (*Tamarindus indica* L) is a plant belonging to the Fabaceae family and the subfamily Caesalpinioideae that is often found in arid regions. The various parts of this tree are of great interest to rural African populations. The pulp of its fruit is used for the preparation of cold drinks or in the composition of several other foods. However, no study in Côte d'Ivoire has yet focused on these foods, only the juices or nectars of this wild fruit tree have been studied. Thus, a survey was conducted in the localities of Korhogo, Séguéla and Bouna with 135 people per locality in order to identify these foods, and then to determine their modes, forms, frequency of consumption and their processes. The survey revealed that most of consumers are without level of education (78.51%), Ivorian (84.94%) and married (80.99%). The places of supply of tomi are often the field (65.18%-

*Corresponding author: E-mail: nestorkksi@yahoo.fr;

71.64%) than the market (32.84-41.67%) and in its pod form (65.93-73.13%). People use pods and pulp to make 10 tomi-based foods with a meal/drink ratio of 7:3. This fruit is consumed by rural populations for health needs, for its taste and for the color it gives to dishes. The frequency of consumption varies from 0 to 3 times per month and the Sougé-baca appears as the most known and consumed food with a percentage of knowledge of 74.07 and a frequency of consumption of 3 times per month. Sougé-baca is a meal more consumed at breakfast and at the Muslim jêun break. The production diagram of *Bacacrou* (Sougé-baca, Badégé-baca, Manou-baca and *Sougé ni manou-baca*) vary according the practices of the three localities.

Keywords: Tomi; pulp; foods; consumption practices; food security; Côte d'Ivoire.

1. INTRODUCTION

The Tamarind tree (*Tamarindus indica* L) is a plant belonging to the Fabaceae family and the Caesalpinioideae subfamily, often found in arid regions [1,2,3]. This tree is proving to be of great interest in the process of monitoring rural African populations [4]. The pulp of this fruit is used for the preparation of fresh drinks [5,6] or is used in the composition of several other foods [7]. However, it is mainly in the form of fresh drinks called *Tomidji* that the fruits of this tree are widely encountered and consumed in Côte d'Ivoire.

From a nutritional point of view, tomi pulp is of great importance due to its energetic content and its richness in vitamins (B1, B2, B3 and C), minerals (P, K, Na, Mg and Ca) [3,8] and phenolic compounds (catechins, tannins, polyphenols and flavonoids) [6,9]. However, more than 40% of the Ivorian population suffers from global acute malnutrition and more than 5% from severe acute malnutrition [10-14]. Given the nutritional richness of Tamarind pulp, the valorization of dishes incorporating this pulp in their production could have a beneficial effect on the health of consumers.

Unfortunately, most of the available data on this wild fruit tree in Côte d'Ivoire only concern the ethnobotanical [1,4,5] and nutritional aspect [15]. Only a few recent studies [6,9] have focused on juices or nectars derived from this wild fruit. To date, no study in Côte d'Ivoire has yet looked at these cereal dishes that incorporate tomi pulp in their preparation. With regard to this situation, the optimization of the level of knowledge of these dishes derived from Tomi is necessary. The objective is therefore to identify all the dishes derived from the incorporation of Tomi and their levels of consumption for its valorization.

2. MATERIALS AND METHODS

2.1 Materials

The biological material consists of tomi pulp (*Tamarindus indica* L) commonly consumed in the north of Côte d'Ivoire. Then a questionnaire containing information such as consumption practices (modes, forms and frequencies of consumption, supplemented by the production process) of tomi pulp-based dishes was used.

2.2 Methods

2.2.1 Selection of the departments and villages

According to Kouakou et al. [16] and Ambé [5], the wild fruit *Tamarindus indica* L. is highly disseminated and consumed in the departments of Korhogo, Séguéla and Bouna. Thus, the consumption survey was carried out in 3 villages of each different department after a pre-survey based on the consumption of tomi pulp and the easy access. So, the villages of Nahouokaha, Lataha and Kotchiéri (Korhogo), Sifié, Sélakoro and Bobi (Séguéla), and Niandégué, Bouko and Panzarani (Bouna) were surveyed.

2.2.2 Sampling

The size (n) of the households surveyed was calculated according to formula described by Dagnelie [17] for an independent non-exhaustive sample based on Côte d'Ivoire Population and Housing Census [18].

$$n = t^2 \cdot \frac{p \cdot (1 - p)}{m^2}$$

n = minimum sample size sought;
t = 95% confidence level (standard value of 1.96);
P = proportion of consumers in the study area;

p estimated at 50 % given the lack of knowledge of the number of households consuming wild fruit trees;
 m = margin of error at 5%.

To compensate for probable errors, 20 households were added in each department. Thus, 405 households were surveyed, i.e. 135 per department (Table 1).

Table 1. Number of households surveyed by zone and by village

Departments	Village	Surveyed	Total
Korhogo	Nahouokaha	48	135
	Lataha	51	
	Kohotiéri	36	
Séguéla	Sifié	41	135
	Sélakoro	44	
	Bobo	50	
Bouna	Niandégué	18	135
	Bouko	62	
	Panzarani	55	
Total			405

2.2.3 Questionnaire survey

The survey was conducted to collect quantitative and qualitative on tomi pulp based dishes in rural zones of Korhogo, Séguéla and Bouna. A questionnaire was drawn up using SPHINX Plus² (V5) software (Version 4.5.0.19) in order to determine their modes, forms, frequency of consumption and their processes of obtaining. This questionnaire had three sections, including basic questions about the respondent (socio-demographic characteristics), questions about the consumption and questions about process.

2.2.4 Data treatment

Survey data were analyzed using SPHINX Plus² (V5) software. Recoding was performed on the SPSS 20.0 software to establish the database and then transferred to the EXCEL 2016 spreadsheet. The data were grouped in tabular form and the percentages of the different parameters were calculated. Descriptive statistics and statistical analyses were performed using the XLSTAT version 7.5 statistical software and the EXCEL 2016 spreadsheet. Descriptive statistics was made to translate data into graphs, averages and standard deviations. Excel software was used to graph periods, consumption frequencies and the level of knowledge of food. Comparisons between dependent variables were determined by the Chi squared test and the Z test at the 5% threshold.

Factor component analyses (CFAs) were performed for comparisons of variables with more than three modalities.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Socio-demographic characteristics of surveyed households

The socio-demographic characteristics of tomi processing and consuming households in Korhogo, Séguéla and Bouna are presented in Table 2. The female/male sex ratio of the 405 respondents is 0.05, with the vast majority (78.51%) not attending school. The age of the users varied from 16 to more than 50 years, with 8.39% aged 16 to 20 years and 31.11% aged 21 to 35 years. People aged 36 to 50 years old constitute 32.34% of these wild fruit users and those over 50 years old represent 28.18% of this population. 84.94% of the respondents were Ivoirians and 15.06% were non-Ivoirians. They were 88.99% married, 10.12% single and 8.89% widowed.

3.1.2 Consumption characteristics of Tomi

3.1.2.1 Places and forms of tomi supply

The locations where tomi is procured or collected in the different localities studied are recorded in Table 3. The data reveal that the places of supply of tomi in the three localities studied are the field and the market. However, tomi is more often collected in the field, with proportions of 65.18% in Bouna, 68.94% in Séguéla and 71.64% in Korhogo, compared to 32.84% in Korhogo, 37.78% in Bouna and 41.67% in Séguéla. Consumers in Korhogo, Séguéla and Bouna obtain tomi only in the form of pods and pulp (Table 4). The collection of tomi in its pod form (65.93-73.13%) is about twice as important as in its pulp form (32.09-41.67%) in all localities.

3.1.2.2 Reasons for tomi consumption

The reasons for tomi consumption are presented in Fig. 1. Discriminant factor analysis shows that tomi is consumed more in the three departments for its taste. In addition to taste, tomi is consumed in Bouna for health reasons, while in Séguéla it is consumed not only for health reasons but also for its color.

Table 2. Socio-demographic characteristics of respondents

Features	Bouna	Korhogo	Séguéla	Total
Sex ratio (M/F)	4.65%	8%	1.50%	4.65%
Level of education (%)				
1 ^{er} cycle	5.93	5.18	8.89	6.67
2 nd cycle	2.96	3.70	2.22	2.96
Primary	14.07	6.67	5.93	8.89
Superior	1.48	5.18	0.74	2.47
Koranic	1.48	0	0	0.49
No	74.07	79.26	82.22	78.51
Origin (%)				
Ivorian	65.93	100	88.89	84.94
No Ivorian	34.07	0	11.11	15.06
Age group (Years) (%)				
[16-20]	8.15	8.15	8.89	8.39
[21-35]	40	18.52	34.81	31.11
[36-50]	31.11	37.04	28.89	32.34
>50 years old	20.74	36.30	27.41	28.15
Marital status				
Married	81.48	77.04	84.44	80.99
Widows	8.15	12.59	5.93	8.89
Singles	10.37	10.37	9.63	10.12

Table 3. Tomi collection locations

	Bouna	Séguéla	Korhogo
Market (%)	37.78	41.67	32.84
Field (%)	65.18	68.94	71.64
z	-4.558	-4.505	-6.3592
p	< 0.001	< 0.001	< 0.001

In the same column, data with the same alphabetical letters are not significantly different at the 5% threshold according to the z-test

Table 4. Forms of Tomi collection

	Bouna	Séguéla	Korhogo
Pod (%)	65.93	68.18	73.13
Pulp (%)	37.78	41.67	32.09
z	4.697	4.363	7.247
p	< 0.0001	< 0.001	< 0.001

In the same column, data with the same alphabetical letters are not significantly different at the 5% threshold according to the z-test




Fig. 1. Reasons for tomi consumption by place of residence

3.1.2.3 Identification and Classification of tomi-based dishes

The ten (10) tomi-based dishes identified in Korhogo, Séguéla and Bouna are recorded in Table 5. Factor analysis of these tomi-based foods made it possible to group the foods consumed by locality (Fig. 2). Some foods are

common to all three zones, while others are specific to them. For example, Sougé-baca, Badégé-baca, and Tomidji are consumed in Korhogo, Séguéla, and Bouna, while Sougé ni manou-baca and Zagamin are consumed only in Séguéla and Korhogo. Manou-baca, Kagbèlè and Kachiansim are specialties of Korhogo. Tôh and Zoom-koom are consumed only in Bouna.

Table 5. Description and vernacular names of tomi-based dishes

Foods	Localities	Consumers	Vernacular names
Millet porridge	Bouna	Koulango	<i>Gôdi-baca</i>
	Korhogo	Sénoufo	<i>Sougé-baca</i>
	Séguéla	Kôyaka	<i>Gnon-monnie</i>
Corn porridge	Bouna	Koulango	<i>Brozougô-baca</i>
	Korhogo	Sénoufo	<i>Badégé-baca</i>
	Séguéla	Kôyaka	<i>Caba-monnie</i>
Rice porridge	Korhogo	Sénoufo	<i>Manou-baca</i>
	Séguéla	Kôyaka	<i>Malo-séri</i>
Millet porridge and rice	Korhogo	Sénoufo	<i>Sougé ni manou-baca</i>
	Séguéla	Kôyaka	<i>Gnon malo-monnie</i>
Corn cake	Korhogo	Sénoufo	<i>Kagbèlè</i>
Millet or rice drink	Bouna	Burkinabé	<i>Zoom-koom</i>
Tomi's fermented drink	Korhogo	Sénoufo	<i>Kachiansim</i>
Tomi juice	Bouna	Koulango	<i>Illanoun-</i>
	Korhogo	Sénoufo	<i>Kachian</i>
	Séguéla	Kôyaka	<i>Tomidji</i>
Cabato of millet with tomi	Bouna	Lobi	<i>Djor</i>
Rice with tomi	Korhogo	Sénoufo	<i>Manoukalo</i>
	Séguéla	Kôyaka	<i>Zagamin</i>




Fig. 2. Factor analysis of tomi-based dishes

Sougé-baca = millet granule porridge; *Badégé-baca* = maize granule porridge; *Manou-baca* = rice granule porridge; *Sougé ni manou-baca* = ¼ millet and ¼ rice granule porridge; *Tomidji* = tomi juice; *Zoom-koom* = cereal juice; *Tôh*= cabato; *Kagbèlè* = fritter; *Zahamin* = fat rice

3.1.2.4 Frequency of consumption of tomi-based dishes

The frequencies of consumption of tomi dishes are presented in Fig. 3. Analysis of this figure shows that Tomidji and Sougé-baca are the most consumed foods, with average consumption frequencies of 3 times per month. Badege-baca comes in third place with an average consumption frequency of 2 times per month. Zagamin and Kagbèlè are the least consumed.

3.1.2.5 Consumption periods for tomi-based dishes

Fig. 4 illustrates the consumption periods of tomi-based dishes. It shows that with the exception of Zagamin, which is only eaten at noon, and Tôh, which is eaten at lunch and dinner, all other tomi-based foods are eaten at breakfast and during the Muslim jêun break. However, Sougé-baca and Badégé-baca, Manou-baca and Kachiansim are eaten more at breakfast, while Tomidji is eaten more at the Muslim jêun break.




Fig. 3. Consumption frequencies of Tomi-based dishes




Fig. 4. Consumption periods of Tomi-based dishes

3.1.3 Production process of tomi-based dishes

3.1.3.1 Production of cereal porridges

The production scheme for cereal porridges is shown in Fig. 5. For the production of Bacacrou, one (1) kg of cereal millet (*Sougé-baca*) or rice (*Manou-baca*) or maize (*Badégé-baca*) or $\frac{3}{4}$ of millet + $\frac{1}{4}$ of rice (*Sougé ni manou-baca*) is soaked in 3 L of water for 120 min. The soaking water is then removed and the cereal is washed again in 3 L of water. After draining, the cereal is ground with 100 g of ginger and/or 20 g of cloves (Bouna) or 10 to 20 g of chili pepper (Korhogo) to obtain a flour. The flour obtained is sifted and progressively sprinkled with 100 to 200 mL of water for the formulation of granules. The granules thus obtained are dried in the sun or at room temperature for 4 to 5 hours. For cooking, 7 to 8 L of water is put in a pot and brought to boil. The granules are added to the pot and mixed as they come out to avoid their agglomeration. A macerate obtained from 250 g of tomi pulp in 0.5 L of water is added gently after 10 to 15 min of cooking. The mixture obtained is left to cook for 10 to 15 minutes and then 500 g of sugar and/or


140 g of vanilla sugar (Bouna) or the juice of 30 g of mint leaves (Séguéla) is added directly.

3.1.3.2 Kagbèlè production

The Kagbèlè production process is shown in Fig. 6. In 3 L of water, 1 kg of maize kernels are soaked for 120 minutes, then removed and washed in 3 L of water. The grains obtained are drained and ground with 100 g of chili pepper to obtain a flour. To this flour, we add a macerate obtained with 500 g of tomi pulp in 1.5 L of water and 30 g of salt. The whole is kneaded by hand during 10 to 15 min and the consistent paste obtained is passed to the frying to obtain fritters in the shape of pancake (Kagbèlè).

3.1.3.3 Production of Toh

The production scheme for Toh is shown in Fig. 7. The macerate of 250 g of tomi pulp in 2 L of water is filtered and heated. After boiling, 0.5 kg of millet flour is gradually added and the mixture is homogenized by swirling until it becomes a slurry. After 5 to 10 minutes of cooking the slurry, 1 kg of millet flour is gradually added. The mixture is regularly kneaded for 10 to 15 minutes before being served.



* Practices of Korhogo
 * Practices of Séguéla
 * Practices of Bouna

Fig. 5. Bacacrou production diagram




Fig. 6. *Kagbèlè* production diagram




Fig. 7. *Tôh* production diagram




Fig. 8. Zoom-koom production diagram

3.1.3.4 Production of Zoom-koom

The description of the production of Zoom-koom is shown in Fig. 8. One (1) kg of grain (millet or rice) is soaked in 3 L of water for 120 min before being removed and washed in 3 L of water. After draining, the cereals are ground with 200 g of ginger and 50 g of cloves. The flour obtained is sifted and relayed in 10 to 15 L of water for 30 to 60 min. A macerate of 250 g tomi pulp in 500 mL water is added to the flour solution and the whole is homogenized, filtered before adding 2 kg sugar and 280 g vanilla sugar.

3.1.3.5 Production of Kackiansim and Tomidji

The description of the production process of Kackiansim and Tomidji is detailed in Fig. 9. A 500 g portion of tomi pulp is boiled in 3 L of water for 2 to 3 minutes. The resulting solution is allowed to stand for 15-30 minutes to cool, then

mixed and the juice is collected after filtering. A mass of 500 g of table sugar is directly added to the filtrate to obtain Tomidji. To obtain Kackiansim, the filtrate is left to ferment for 24 hours, then 500 g of table sugar and 200 g of chilli powder are added.

3.1.3.6 Production of Zagamin

The description of zagamin production is shown in Fig. 10. A 1-2 kg mass of meat or fish is fried in 0.25 L of vegetable oil for 5-10 minutes. To this mass 100 g of onion, 50 g of carrot, 100 g of cabbage and 40 g of salt are added before simmering for 5 minutes. Add 30 g of pepper and macerate of 250 g of tomi pulp in 1.5 L of water and leave the mixture on the fire until it boils. When it boils, one (1) kg portion of rice is put in the sauce solution for 15 to 20 minutes of cooking on low heat.




Fig. 9. Kackiansim and Tomidji production Diagram




Fig. 10. Zagamin production diagram

3.2 Discussion

This study consisted in identifying and collecting information on the dishes integrating the pulp of Tomi (*Tamarindus indica* L.) in their formulations in Côte d'Ivoire. The results obtained show the importance of the valorization of local dishes for food security.

From a sociodemographic point of view, the sex ratio of respondents is in favor of women, with a proportion of 95.95%. This high proportion of women could be justified by the fact that they are the ones who generally do the cooking in rural

households and are therefore more skilled in the preparation and knowledge of the dishes consumed. This result is in agreement with Adepoju et al. [19] who identify women as the main pillars of food security in rural households.

The identification of tomi -based foods in the departments of Korhogo, Séguéla, and Bouna revealed 10 tomi-based foods. This diversity of foods produced from this wild fruit could contribute more to the food security of rural and urban populations. This thought is corroborated by the studies of Lockett and Grivetti [20], according to which they stipulate that wild edible

plants are incorporated into foods to improve or diversify them. These authors also argue that the wild edible plant foods have been a mainstay and supplement to the Fulani diet since ancient times. Therefore, the popularization of these foods could help reduce the food insecurity that affects developing countries, particularly in Côte d'Ivoire. According to studies by Sackou et al. [21] on food insecurity in the city of Abidjan, only 3.8% of households were food secure.

From a culinary point of view, this wild fruit tree can be transformed into several food forms. Thus, foods made from tomi pulp are generally consumed in the form of meals and drinks with a meal/drink ratio of 7:3. The high use of these wild fruits in the diet as meals would reflect their importance in the dietary habits of local populations. These different forms of food use as drinks or meals are consistent with those described by several authors [22,23]. Indeed, these authors indicate that tamarind pulp is consumed in non-alcoholic sweetened beverages or mixed with ginger, for the preparation of ice creams. It can also be used in weaning foods. Three (3) of the 10 prepared foods (Sougé-baca, Badégé-baca and Tomidji) are consumed in all localities with some small differences related to production process.

The similarity of the foods consumed in the different study areas could be explained by the interaction between the peoples. Van der Stege et al. [4] corroborate this hypothesis, suggesting that the similarities and differences in tamarind food use are related to knowledge sharing and general interactions between ethnic groups living in close proximity to each other. Kruger and Gericke [24] and Thurber et al. [25] continue this view by stating that it is evident that there is divergence in the foods consumed, as culture is one of the important factors that influence consumer attitudes toward a given food.

For the preparation of all these dishes, tomi is generally collected in the field to the extent of 65.18% to 71.64% depending on the locality. This result could be explained mainly by the fact that the surveys were conducted in rural areas and secondarily by its strong presence in these three different areas of northern Côte d'Ivoire. This is corroborated by the studies of Kouakou et al. [16] and Andon et al. [26] who reported the abundance of this wild fruit tree in these areas.

The results of the study show that the reasons for the uses of tomi in food making vary by

locality ranging from health needs, taste and color of dishes. These reasons are confirmed by [22,27,28,29] who argue that tomi is added to meals for digestion, to enhance taste or to protect food from bacteria. In fact, despite its acidic nature, tomi is described as the most acidic and sweetest fruit at the same time. It is one of the few fruits that retain its tartaric acid content estimated at 98% of the organic acids contained in its pulp during ripening, while the amount of reducing sugars increases to give it a sweeter taste [8]. Tartaric acid is an antimicrobial agent that inhibits the growth of food spoilage microorganisms by lowering the pH of the cell [30,31] and the production of malic acid, which is a key intermediate in the production of glucose in the process of gluconeogenesis, the main fuel of cells [30].

Study data reveal that all foods made by these populations are mostly consumed at breakfast and during the Muslim lent especially porridges and drinks. These results confirm those of Sadiq et al. [7] and N'Guessan et al. [32] who stipulates that porridges is consumed at breakfast, snack time and is highly valued during Ramadan. Indeed, the results show that Sougé-baca porridge is more known and consumed. This preference for millet porridge could be explained by its good acceptability but also by the availability of millet in the study areas. This hypothesis is confirmed by Tou et al. [33] who report the good organoleptic quality of millet porridge compared to maize porridge. This result could also be due to the fact that according to N'guessan et al. [32], millet porridge is the main form of millet processing. Regarding the availability of millet, Koffi et al. [34] reported that millet and sorghum are traditionally grown in the northern region of Côte d'Ivoire. Parry [35], confirms this report by adding that the northern area of Côte d'Ivoire is the most suitable for growing millet. However, the use of tomi in several foods by rural populations clearly indicates that beyond the ignorance of the benefits of these fruits, they are all the same exploited by these populations. But the level of knowledge and information about these foods limits their use.

4. CONCLUSION

The objective of this study was to identify foods made from tomi fruits (*Tamarindus indica* L.) in Côte d'Ivoire. At the end of this study, ten (10) tomi-based foods were identified. The production processes and consumption patterns of these

foods vary according to the localities studied, with a meal/drink ratio of 7:3. This fruit is consumed by rural populations for health reasons, for its taste and for the color it gives to dishes. Millet porridge (Sougé-baca) is the most known and consumed food with a percentage of knowledge of 74.07 and a frequency of consumption of 3 per month. However, a better knowledge of the impact of these different foods on the well-being of these consumers is necessary given the nutritional importance of this fruit. It would be interesting to determine the biochemical composition of these foods for a better popularization. This was made in rural areas. So, we need to extend the survey to urban areas for a broader mapping of tomi-based dishes consumed.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Diallo BO, Joly HI, McKey D, Hossaert-McKey M, Chevallier MH. Population genetic diversity of *Tamarindus indica*: Clues to the origin of its current distribution? *African Journal of Biotechnology*. 2007;6(7):853-860.
2. Morton J. *Tamarind (Tamarindus indica L.)*. In: Morton JF. *Fruits of Warm climates*. FI, USA; 1987.
3. El-Siddig K, Gunasena HPM, Prasad BA, Pushpakumara DKNG, Ramana KVR, Vijayanand P et al. *Tamarind, Tamarindus indica L.* International Centre for Underutilised Crops. University. Southamp U.K.; 2006.
4. Van der stege C, Prehsler S, Hartl A, Vogl CR. *Tamarind (Tamarindus indica L.) in the traditional West African diet: not just a famine food*. *P Sciences All Rights Reserved*. 2011;66: 171-185 .
5. Ambé GA. Les fruits sauvages comestibles des savanes guinéennes de Côte d'Ivoire : état de la connaissance par une population locale, les Malinkés. *Biotechnologie Agronomie Société Environnement*. 2001;5(1):43-58. French.
6. Kouassi KA. Study of the nutritional value of fruit pulp of three edible wild fruit species in Côte d'Ivoire and biochemical and sensory characterization of derived nectars: *Adansonia digitata* (Baobab), *Tamarindus indica* (Tomi) and *Parkia biglobosa* (Néré). PhD thesis in agriculture and tropical forestry, speciality: Biochemistry and Nutrition. Université Jean Lorougnon Guédé de Daloa-Côte D'Ivoire. 2019;164.
7. Sadiq IS, Duruminiya NI, Balogun JB, Kwada D, Izuagie T. Nutritional and anti-nutritional value of tamarind (*Tamarindus indica*) fruit. *International Journal of Applied Research and Technology*. 2016; 5(3):50-56.
8. Grollier C, Debien C, Dornier M, Reynes M. Principales caractéristiques et voies de valorisation du tamarin. *Fruits*. 1998;53:271-280. French
9. Kouassi AK, Kouassi NK, Beugré MAG, N'dri DY, Amani GN, Gnakri D. Glycemic index and glycemic load of wild edible fruit juices (*Adansonia digitata*, *Tamarindus indica* and *Parkia biglobosa*) consumed in Côte d'Ivoire. *Journal of Biosciences and Medicine*. 2018;06(01):63-74. DOI: 10.4236/jbm.2018.61007.
10. FAO. Nutrition education program integrated with community nutrition centers Côte d'Ivoire. FAO Nutrition and Consumer Protection Division (AGN). 2008;2.
11. WHO. Global prevalence of vitamin A deficiency in populations at risk 1995-2005. WHO Global Vitamin A Deficiency Database; 2009.
12. Schwartz H, Grellety Y. Evaluation of the management of acute malnutrition Côte d'Ivoire: Mission report. UNICEF West and Central Africa Regional Office. 2010;29.
13. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F et al. Understanding barriers to fruit and vegetable consumption in the Australian Longitudinal Study of Indigenous Children: A mixed methods approach. *Public Health Nutrition*. 2016;20(5):832-847. DOI:https://doi.org/10.1017/S1368980016003013
14. N'goran P. Sécurité alimentaire et nutritionnelle en Côte d'Ivoire: Etat des lieux, défis. National Nutrition Program. 2014;42. French
15. Cissé I. Caractérisation des propriétés biochimiques et nutritionnelles de la pulpe de baobab des espèces endémiques de Madagascar et d'Afrique continentale en vue de leur valorisation. Thèse, Ecole doctorale: Sciences des Procédés-

- Sciences des Aliments. Montpellier Supargro, France. 2012;6-77. French.
16. Kouakou YB, Malan DF, Kouassi KG, Diop AL, Bakayoko A. Availability of some spontaneous food plants used by Koulango and Lobi populations in the eastern periphery of Comoé National Park, Côte d'Ivoire. *Africa Science*. 2020; 16(3):33-50.
 17. Dagnelie P. Statistiques théoriques et appliquées. Paris, De Boeck & Larcier. 2th ed. Belgique; 1998. French.
 18. RGPH. General census of population and housing. 2014. 5p.
 19. Adepoju AA, Ogunniyi LT, Agbedeyi D. The role of women in household food security in Osun State, Nigeria. *International Journal of Agricultural Policy and Research*. 2015;3(3):104-113.
 20. Lockett C, Calvert C, Grivetti L. Energy and micronutrient composition of wild food and medicinal plants consumed during drought. A study of rural Fulani, northeastern Nigeria. *International Journal of Food Science and Nutrition*. 2000; 51(3):195-208.
 21. Sackou KJ, Lobilé YH, Kouadio KKL. Household food insecurity and association with sociodemographic factors in Abidjan. *European Journal of Public Health*. 2020; 30(5):166-220.
 22. N'Diaye M, Kéita FB, Martin P. Principaux fruits de cueillette consommés et commercialisés en Guinée. *Fruits*. 2003;58:99-116. French.
 23. Nkama I, Iliyas A, Jato A. Studies on the preparation and nutrient composition of kunun gyada, a traditional Nigerian groundnut-cereal-based weaning food. *Food and Nutrition Bulletin*. 1995; 16(3):238-240.
 24. Kruger R, Gericke GJ. A qualitative exploration of feeding and weaning practices, knowledge and attitudes about nutrition in rural areas. *Public Health Nutrition*. 2002;6(2):217-223.
DOI:<https://doi.org/10.1079/PHN2002419>.
 25. Thurber KA, Banwell C, Neeman T, Dobbins T, Pescud M, Lovett R et al. Understanding barriers to fruit and vegetable intake in the Australian Longitudinal Study of Indigenous Children: A mixed-methods approach. *Public Health Nutrition*. 2016;20(5):832–847.
DOI:<https://doi.org/10.1017/S1368980016003013>
 26. Andon GS, Assouman SF, Soro GR. Loss of trees in the agricultural landscape of northern Côte d'Ivoire: the case of the Korhogo department. *Revue de Géographie Tropicale et d'Environnement*. 2018;(2):49-60.
 27. Gustad G, Dhillion SS, Sidibe D. Local use and cultural and economic value of tree products in the parks of the Cinzana commune, Mali. *Economic Botany*. 2004; 58(4):578-587.
 28. Berry L, Ayensu E, Mounkeila G, LeBeau F, Weber F, Corillon C et al. Agroforestry in the West African Sahel, BOSTID, World Environ. Libr., Waikato, Nouvelle-Zélande; 1984.
 29. Etkin NL, Ross PJ. Food as medicine and medicine as food-An adaptive framework for the interpretation of plant utilization among the Hausa of northern Nigeria. *Social Science and Medicine*. 1982; 16:1559-1573.
 30. Shokri H. Evaluation of the inhibitory effects of citric and tartaric acids and their combination on the growth of *Trichophyton mentagrophytes*, *Aspergillus fumigatus*, *Candida albicans* and *Malassezia furfur*. *Comp Clin Cathol*. 2011;20:543-545.
DOI: 10.1007/s00580-011-1195-6.
 31. Jay JM. Modern food microbiology. Aspen Publishers, Gaithersburg, 2000;268.
 32. N'Guessan YD, Bedikou ME, Megnanou RM, Niamke SL. Importance of local cereal flours in the food habit of consumers in the district of Abidjan Côte d'Ivoire. *Journal of the Faculty of Food Engineering, University Ștefancel Mare-Suceava*. 2014;8(2):134-146. French.
 33. Tou ER, Mouquet-Rivier C, Traoré AS, Trèche S, Guyot JP. Nutritional and microbial characteristics of ben-saalga, a fermented millet-based porridge: from Burkina Faso used as a complementary food. *International Congress of Nutrition*. Durban, South Africa; 2006.
 34. Koffi KE, Yao SDM, Diarrassouba N, Siene LAC. État des lieux et gestion des semences des principales céréales cultivées au nord de la Côte d'Ivoire. *Journal Scientifique Européen*. 2017; 13(3):112.

35. Parry G. Le cotonnier et ses produits. Maisonneuve et Larousse et ACCT.
Collection. Technique agricole et 1982;30-404.
production végétale. Paris, France

© 2022 Jacqueline et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91141>