



A Review on Breeding in Ornamental Crops for Abiotic Stress Tolerance

**Heera Lal Atal ^{a*}, Daripalli Srilakshmi ^a, Kunal Debbarma ^a, Lopamudra Jena ^a
and Meikam Ichancha ^b**

^a Department of Floriculture and Landscape Architecture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, West Bengal, India.

^b Department of Floriculture and Landscaping, Bidhan Chandra Krishi Viswavidyalaya, West Bengal, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2031136

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87700>

Review Article

Received 22 March 2022

Accepted 01 June 2022

Published 09 June 2022

ABSTRACT

Abiotic stresses are major concerns in agriculture sector affecting large parts of India. It is caused by Drought, salinity and temperature (high/low) representing a major constraint over crop growth rate and productivity. In ornamental crops meagre efforts have been done towards this way. Now a day's ornamental crops are the prospective way to earn higher income as compared to the other crops and have a significant role in national GDP. How can we grow Ornamental crops in stressed condition is one of the major challenge. To get tolerance power against abiotic stress various ways have been developed but among all methods very few have been proved worthy in case of ornamental crops. Interspecific and intergeneric hybridization is the basic way found hopeful strategy to improve tolerance against drought in some crops through domesticating abiotic stress-tolerant gens from their wild species into cultivated. Combination of both tissue culture and induces mutation approach (*In vitro* mutagenesis) is the one more important way for bringing stress tolerance and improvement in yield and quality of Ornamental crop plants. Genetic engineering is the modern tool which is used for breeding programme. The introduction of innovative characteristics such as new colours, biotic and abiotic stress tolerance in ornamental crops are usually difficult through predictable breeding, but by the use of genetic transformation we can get it easily.

*Corresponding author: E-mail: heera.atal93@gmail.com;

Keywords: Flower breeding; hybridization; mutation; genetic engineering; abiotic stress.

1. INTRODUCTION

Phenotypic findings of a plant are displayed by its genotype, environment and the genotype environment interaction. When some environmental factors interfere with the complete appearance of genotypic potential, it is called stress. There are two types of stress, the first one is biotic (pathogens, pests, weeds, etc.) and second is abiotic depending on their biological/abiological nature. "Abiotic stresses outcome due to moisture, temperature (high/low), minerals (deficiency/toxicity), salinity, soil pH, air pollution, etc. Abiotic stress situations cause wide losses to agricultural production worldwide" [1]. "Exclusively, stress conditions such as drought, salinity or heat have been the subject of intense research" [2]. However, in field condition, crops and other plants are usually subjected to a combination of different type abiotic stresses. In drought-prone areas, for example, many crops encounter a mixture of drought and other stresses, such as soil salinity or heat.

2. CHARACTERISTICS OF ABIOTIC STRESSES

1. The physiognomies of an abiotic stress may vary significantly depending on the location.
2. The relative importance of different abiotic stresses is mainly based on location.
3. During the crop season variation may occur in the degree of some stresses.
4. A given abiotic factor may affect the level of another abiotic stress, e.g., in a saline soil, moisture stress would enhance salinity stress.
5. Different plant/crop species having different abilities to withstand in a given stress.
6. Different varieties of a crop also can have difference in their capabilities to tolerate abiotic stresses.
7. One abiotic stress can overlap the effect of another stress. For example, salinity stress generated some features produce by drought stress, so that strains develop for salt resistance also show enhanced drought tolerance.

Drought: Drought is a condition of unusually dry weather within a geographic area resulted from the lack of rains. It is directed by various factors like high temperature, photon irradiance and

scarcity of water. Drought is a low water potential condition resulted from high solute concentration.

3. MECHANISM OF DROUGHT TOLERANCE

Drought escape: Drought escape can define as the aptitude of a plant or crop species to complete its life cycle before supply of water in soil is washed-out and became dormant seeds before the arrival of dry season. That type of plants is known as drought escapers since they skip the drought condition by rapid development.

Drought avoidance: These type of plants can store high amount of water to maintain relatively high tissue- water potential despite a shortage of soil-moisture. Drought avoidance is governed by maintenance of turgor through roots grow in deep soil, transpiration controlled by stomata to decrease water and by reduction of water loss through reduced epidermal i.e. reduced surface by smaller and thick leaves.

Drought tolerance: Those plant having ability to withstand in drought condition with low tissue water potential. Drought tolerance is the manner of turgor preservation via osmotic adjustment (a manner which induces solute accumulation in cell), increase in elasticity and reduce in cell size.

4. EFFECT OF DROUGHT STRESS

Effect on Growth: Fall in turgor Pressure, due to cell sizes will be smaller.

Effect on Photosynthesis: Photosynthesis decreases due to disruption of PS II (Photo System II), stomatal closure, decrease in electron transport.

Decrease in nuclear acids and proteins: Protease activity increase, free amino acid increase, RNAase activity increase, RNA hydrolysis, DNA content falls down.

Effect on Nitrogen Metabolism: Nitrate reductase activity decrease, nitrite reductase activity insensitive.

Effect on Carbohydrate metabolism: Loss of starch and increase in simple sugars, carbohydrate translocation decreases.

Salt stress: “Salt stress occurs when there is an excessive amount of salt in the soil, which inhibits crop growth and finally leads to crop death. Reduction of water use efficacy, increase ions, induction of heat stress and reduces stem extension in plants caused by salt stress. Biological macromolecules are broken down due to generation of free ions. It was noticed that salinity stress resulted in decrease of high water content, production of hydrogen peroxide and electrolyte generate in plants” [3]. “Salt accumulation is one of the hazardous factor in soil decreasing the production and quality of plants, causes adverse effects on sprouting, plant growth and development. Throughout the world, nearby 45 million hectares of irrigated land is affected by salt stress” [4]. Salt stress affects plants in various means such as; ion toxicity, physiological drought, nutritional disorders, modification of metabolic processes, oxidative stress, membrane incompetence and reduction of cell division.

The response of ornamental plants to salt stress: Salt effects on plants are the combined outcome of the composite interaction among various morphological, physiological and biochemical processes. Among all outcomes of salt stress reduction in rate of leaf growth is one of the first responses of plants to salinity [5] “primarily due to the osmotic effect of salt around the roots, which leads to a reduction in water supply to leaf cells. Availability of higher salt concentrations in soil can also inhibit root growth [6], with a reduction in length and total mass of plant roots” [7] and of function. “Reduction in rate of cell elongation and division in leaves reduces their final size, resulting in a decrease in leaf area” [8,9]. Leaf area reduction could be produced by a decrease in turgor in the leaves, as a significance of changes in cell wall properties or a reduction in photosynthetic rate. Such concerns are seen in ornamental plants. As per suggested by Cassaniti et al., [10] “Loss in dry weight of shoot and leaf area were noticed as the first visible effects of salinity both in sensitive and tolerant species such as *Cotoneaster lacteus* and *Eugenia myrtifolia*, respectively”. “Leaf thickening is another common effect of heavy salt level, which appeared in ornamental plants such as *Coleus blumei* and *Salvia splendens*” [11].

Heat stress: Heat stress is commonly characterised as a temperature rise that exceeds a threshold level for a long enough period of time to cause irreparable damage to plant growth and

development. The negative effect of temperature higher than the optimal is called as heat stress. Plant survival, growth and development and the physiological processes would be affected by heat, the nature and extent of the effects depending mainly on the temperature, the plant species and the process in question. Heat stress can also define as the induction in temperature beyond a threshold level for a period of time sufficient to cause unalterable destruction to plant growth and development [12]. Heat stress affects factory growth throughout its ontogeny, though heat- threshold position varies vastly at different experimental stages. Heat stress due to high ambient temperature is a serious trouble to crop product worldwide. Different worldwide circulation models forecast that greenhouse gases will gradually increase world’s average ambient temperature. Because of very high temperatures, severe cellular damage and even cell death may occur within few minutes. At relatively high temperatures, direct injuries include protein declination, aggregation and increased fluidity of membrane lipids. Circular injuries include inactivation of enzymes, inhibition of protein conflation, protein declination and loss of membrane integrity.

Cold stress: Cold stress, which includes chilling (0–15°C) and freezing (<0°C), is an abiotic stress that has a negative impact on plant growth and agricultural productivity. Plant growth and development are frequently restricted by chilling stress, which has numerous important effects on plant cells. Cold stress is a thoughtful threat to the sustainability of crop yields. It can lead to major crop losses. Poor germination, stunted seedlings, yellowing of leaves, reduced leaf expansion and wilting, and may lead to death of tissue (necrosis) are different phenotypic symptoms of plants in response to cold stress. Cold stress also harshly affects the development of plants reproductive systems. “Severe membrane damage is one of the major negative effect of cold stress. Plants display cold or chilling stress at temperatures from 0-15 °C. Under such situations, plants try to maintain homeostasis to acquire freezing tolerance and this involves extensive reprogramming of gene expression and metabolism” [13,14].

5. BREEDING METHODS FOR ABIOTIC STRESS TOLERANCE

Wide distant hybridization: “Wide distant (interspecific and intergeneric) hybridization has become a promising approach to increase

tolerance power against abiotic stresses of some crops through introducing stress-tolerant trait from relative wild species to cultivated, for the wild species usually have high abiotic stress-tolerant trait that cultivars do not possess" [15,16]. "For example, an improved drought-tolerant perennial ryegrass (*Lolium perenne*) was developed through intergeneric hybridization by inheriting drought tolerance of Atlas fescue (*Festuca mairei*)" [17]. "In addition, many cultivars of chrysanthemum are developed with upgraded drought tolerant have been bred through interspecific hybridization" [18]. Develop interspecific hybrids between *Dendrenthema morifolium* and *Dendrenthema nankingense* by using ovary rescue, to improve cold tolerance in cultivated species [19].

In vitro mutagenesis: To improve stress tolerance as well as yield and quality of crop plants, *In vitro* mutagenesis is an important technique. *In vitro* mutagenesis technique is a combination of both tissue culture technique and induce mutation approach. One NaCl tolerant chrysanthemum species (*Chrysanthemum morifolium* Ramat.) variant (E2) has been converted in a stable form by using *in vitro* mutagenesis using ethylmethanesulfonate (EMS) as the chemical mutagen. Salt stress tolerance was displayed by the ability of the plant to maintain both flower quality and yield under stress conditions. Enhanced tolerance of the E2 variant has been attributed to the increased activity of superoxide dismutase (SOD), ascorbate peroxidase (APX), and dehydroascorbate reductase (DHAR), and, to a lesser extent of membrane damage than NaCl treated control plants. Isoform analysis shown that a rise in total SOD exertion in the E2 variant was solely due to significant activation of the Cu/Zn isoform. Elevated situations of carotenoids and ascorbate in E2 leaves have been reflected in their advanced free radical scavenging capacity (RSC) expressed in terms of DPPH (-diphenyl-1 picrylhydrazyl) scavenging capability. Data reflect that a proper balance between enzymatic and non-enzymatic defence systems is needed for combating salt stress in chrysanthemum. "More performance of the E2 get under same salt stress condition, indeed in the alternate time, confirms the inheritable stability of the swab- forbearance character. On the whole, the E2 variant, developed through 0.025 EMS treatment, might be considered as a NaCl tolerant strain showing positive characters towards NaCl stress" [20].

Genetic engineering for abiotic stress tolerance: Development of plant varieties with a high level of tolerance against abiotic stresses is essential for getting full yield potential and to stabilize production. Due to the crowd of abiotic stresses and their complex genetic control the development of breeding for tolerance to abiotic stresses using conventional methodologies has been very disappointing. "Stress tolerant crop production by genetic engineering need to identify the key of genetic factors underlying stress tolerance in plants and introducing these genes into crops. Introduction of molecular change in crop plants by genetic engineering takes less time as compared to plant breeding methods, only wanted gene can be transferred, whereas, in traditional breeding methodology is connected with simultaneous transfer of undesired gene" [21]. Recent advances in cellular and molecular biology have made it possible to clone important genes and mobilize them in any organism across barriers of sexual hybridization for stable expression and transmission. All living organisms have evolved mechanisms for avoidance or forbearance to one or further of the abiotic stresses. Plants producing pivotal enzymes or proteins from different organisms involved in abiotic stress forbearance mechanisms have shown significant advantage over their wild type controls under stressed-out terrain. The enhanced position of compatible osmolytes, radical scavengers and other transgene products identified with the degree of forbearance. Farther understanding of the molecular mechanisms of stress perception, signal transduction and response by shops and other organisms may help to mastermind plants with high situations of forbearance to multiple stresses. Perspectives and fresh approaches for further perfecting the forbearance to abiotic stresses through inheritable engineering are banded [22].

6. CONCLUSION

Abiotic stress caused by temperature, drought or salinity represents a major constraint over crop growth and productivity. Breeding methods like interspecific, intergeneric hybridization, *in vitro* mutagenesis and genetic engineering for improving abiotic stress tolerance in ornamental crops have proven to be the potential approaches. Genetic engineering improved abiotic stress tolerance through adding one or more new traits that are not already found in that organism.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Boyer JS. Plant productivity and environment. *Science*. 1982;218:443-448.
2. Cushman JC, Bohnert HJ. Genomic approaches to plant stress tolerance. *Current Opinion in Plant Biology*. 2000;3:117-124.
3. Mandhania S, Madan S, Sawhney V, Haryana CCS. Antioxidant defense mechanism under salt stress in wheat seedlings. *Biologia Plantarum*. 2006;50(2):227-231.
4. Munns R, Tester M. Mechanisms of salinity tolerance. *Annual Review of Plant Biology*. 2008;59:651-681.
5. Blum A. Salinity resistance, In: *Plant Breeding for Stress Environments*, 1163-1169, CRC Press, Boca Raton; 1986.
6. Wild A. Russell's soil conditions and plant growth. 11th edn. Harlow, Longman; 1988.
7. Shannon MC, Grieve CM. Tolerance of vegetable crops to salinity. *Scientia Horticulturae*. 1999;78:5-38.
8. Alarcon J., Sanchez-Blanco MJ, Bolarin MC, Torrecillas A. Water relations and osmotic adjustment in *Lycopersicon esculentum* and *L. pennellii* during short term salt exposure and recovery, *Physiologia Plantarum*. 1993;89:441-447.
9. Matsuda, K, Riazi A. Stress-induced osmotic adjustment in growing regions of barley leaves. *Plant Physiology*. 1981;68:571-576.
10. Cassaniti C, Li Rosi A, Romano D. Salt tolerance of ornamental shrubs mainly used in the Mediterranean landscape. *Acta Horticulturae*. 2009;807:675-680.
11. Ibrahim KM, Collins JC, Collin HA. Effects of salinity on growth and ionic composition of *Coleus blumei* and *Salvia splendens*. *Journal of Horticulture Science*. 1991;66:215-222.
12. Hall AE. Breeding for heat tolerance. *Plant Breeding Reviews*. 1992;10:129-168.
13. Cook D, Fowler S, Fiehn O, Thomashow MF. A prominent role for the CBF cold response pathway in configuring the low temperature metabolome of *Arabidopsis*. *Proceedings of the National Academy of Sciences (USA)*. 2004;101:15243-15248.
14. Thomashow MF. Plant cold acclimation: freezing tolerance genes and regulatory mechanisms. *Annual Review of Plant Biology*. 1999;50:571-599.
15. Abraham EM, Huang B, Bonos SA, Meyer WA. Evaluation of drought resistance for Texas bluegrass, Kentucky bluegrass, and their hybrids. *Crop Science*. 2004;44:1746-1753.
16. Cattivelli L, Rizza F, Badeck FW, Mazzucotelli E, Mastrangelo AM, Francia E, Mare C, Tondelli A, Stanca AM. Drought tolerance improvement in crop plants: an integrated view from breeding to genomics. *Field Crops Research*. 2008;105:1-14.
17. Wang JP, Bughrara SS. Evaluation of drought tolerance for Atlas fescue, perennial ryegrass, and their progeny. *Euphytica*. 2008;164:113-122.
18. Cheng X, Chen S, Chen F, Deng Y, Fang W, Tang F, Liu Z, Shao W. Creating novel chrysanthemum germplasm via interspecific hybridization and backcrossing, *Euphytica*. 2010;177:45-53.
19. Cheng X, Chen SM, Chen FD, Fang WM, Deng YM, She LF. Interspecific hybrids between *Dendranthema morifolium* (Ramat.) Kitamura and *D. nankingense* (Nakai) Tzvel achieved using ovary rescue and their cold tolerance characteristics. *Euphytica*. 2009;172:101-108.
20. Hossain Z, Mandal AKA, Datta SK, Biswas AK. Development of NaCl-tolerant strain in *chrysanthemum morifolium* through *in vitro* mutagenesis. *Plant Biology*. 2006;8(4):450-461.
21. Roy B, Noren SK, Mandal AB, Basu AK. Genetic engineering for abiotic stress tolerance in Agricultural crops. *Biotechnology*. 2011;10:1-22.
22. Dunemann F, Illgner R, Stange I. Transformation of *rhododendron* with genes for abiotic stress tolerance. *Acta horticulturae*. 2002;113-120.

© 2022 Atal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/87700>