



Effect of Nitrogen and Spacing on Growth and Yield of Sunflower (*Helianthus annuus* L.)

Eslavath Sandhya^{a++*}, Rajesh Singh^{a#},
Akankhya Pradhan^{b†} and Thakur Indu^{b†}

^a Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Naini, Prayagraj-211007, Uttar Pradesh, India.

^b Sam Higgin Bottom University of Agriculture, Technology and Naini, Prayagraj-211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i62855

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97642>

Original Research Article

Received: 12/01/2023

Accepted: 17/03/2023

Published: 22/03/2023

ABSTRACT

A field experiment was carried out during *kharif* season of 2022, at crop research farm of Department of Agronomy at Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj in North Eastern plains of Eastern Uttar Pradesh, India with the objective to study the effect of nitrogen and spacing on growth, yield and economics of Sunflower (*Helianthus annuus* L.) with 10 treatments of which treatments (T₁-T₉) with different combination of nitrogen along with spacing and T₁₀. The soil experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1). The result revealed that application of 70 kg N/ha + 55 X 20 cm recorded Maximum plant height (148.48 cm), plant dry weight (60.53 g), test weight (41.40 g), capitulum diameter (15.13 cm).

⁺⁺ P.G. Scholar;

[#] Associate Professor;

[†] Ph.D. Scholar;

*Corresponding author: E-mail: sandhyaeslavathi09250@gmail.com;

Keywords: Economics; growth; sunflower; nitrogen (N); spacing; yield.

1. INTRODUCTION

“Sunflower is one of the most important oil seed crop grown in temperate countries. It is a major source of vegetable oil in the world. In India it has gained popularity due to the national priority of vegetable oil production. India is one of the largest producers of oil seed crop in the world. Oil seeds occupy an important position in the Indian agricultural economy. It is an important oil seed crop contributes 14% of the total oilseed production from other major oil seed crops” [1-6].

“Among nutrients, nitrogen plays an important role in growth and yield of sunflower. Nitrogen is crucial for growth and development while, sulphur fertilization is most critical for oil and protein synthesis besides seed yield enhancement. The nitrogen requirement for sunflower is high. Oil content in sunflower seed is also affected by changes in the proportion of nitrogen [7-18]. Nitrogen deficiency reduces vegetative and generative growth, thereby potentially reducing yields” [19]. “Increasing Nitrogen increase number of seeds per head seed mass and overall oil yield [20] is a major essential element and is responsible for increasing the photosynthetic surface area and in turn increases the translocation of photosynthates to sink and results in increase in productivity”. “low application of nitrogen enhanced the duration but decreased leaf expansion and distribution of dry matter” [21].

The plant spacing is an important factor, the optimum plant population should be maintained in the field for getting higher yield per unit area. Diepenbrock et al. [22] reported that achene yield increased with increasing row spacing. Overcrowding effects sunflower yield and achenes oil percentage [23]. The number of seeds per head and the mean seed weight decrease significantly with an increasing plant density (Barros et al. 2004) Sunflower crop express its full potential when it is also backed up by plant density [24-31]. Optimum plant density provides better conditions for proper light interception throughout the crop growth period [32-41]. Further, it is important to realize that plant density should be defined not only in terms of number of plants per unit area but also in terms of arrangement of these plants on the

ground (planting geometry/spatial arrangement) as it helps in efficient harvesting of solar energy with least competition [42-49].

“Plant spacing effects are highly pronounced in sunflower because there is no possibility of covering gaps between plants by branching or tillering. Thus, an optimum plant stand helps in harnessing the natural resourced in efficient manner towards achieving high crop yields” [1]. Beside this, spacing influences crop yield through its influence on light interception, rooting pattern and moisture extraction pattern [50-59]. The high-density planting system (HDPS) is now being conceived as an alternate production system having a potential for improving productivity, profitability, efficiency, reducing input costs and minimizing the risks associated with crop production system.

2. MATERIALS AND METHODS

A field experiment was conducted during kharif season of 2022, at Crop research farm of Department of Agronomy at Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj which is located at 25°24' 42" N latitude, 81°50' 56" E longitude and 98 m altitude above the mean sea level (MSL). To assess the effect of nitrogen and spacing on growth and yield of Sunflower (*Helianthus annuus* L.). The experiment was laid out in Randomized Block Design comprising of 10 treatments which are replicated thrice. Each treatment net plot size is 3m × 3m. The treatment are categorized as with recommended dose of Nitrogen through Urea, Phosphorus through DAP and Potash through Muriate of Potash, in addition with sulphur when applied in combinations as follows, (T₁) 50 kg N/ha + 45 X 20 cm, (T₂) 50 kg N/ha + 55 X 20 cm, (T₃) 50 kg N/ha + 65 X 20 cm, (T₄) 60 kg N/ha + 45 X 20 cm, (T₅) 60 kg N/ha + 55 X 20 cm, (T₆) 60 kg N/ha + 65 X 20 cm, (T₇) 70 kg N/ha + 45 X 20 cm, (T₈) 70 kg N/ha + 55 X 20 cm, (T₉) 70 kg N/ha + 65 X 20 cm, (T₁₀) Control. “The sunflower crop was harvested treatment wise at harvesting maturity stage. Growth parameters viz. plant height (cm), dry matter accumulation g plant/ha were recorded manually on five randomly selected representative plants from each plot of each replication separately and after harvesting, seeds were separated from each net plot and were dried under sun for three days. Later winnowed, cleaned and grain yield per ha was

computed and expressed in kgs per hectare. After complete drying under sun for 10 days stover yield from each net plot was recorded and expressed in kgs per hectare. The data was computed and analysed by following statistical method" of Gomez and Gomez (1984). The benefit: cost ratio was worked out after price value of seed with stover and total cost included in crop cultivation.

3. RESULTS AND DISCUSSION

3.1 Effect on Yield of Sunflower

The observation related to yield attributing parameters were shown in Table 1. Treatment of 70 kg N/ha + 55 X 20 cm resulted in significantly highest capitulum diameter (15.13 cm). However, 60 kg N/ha + 55 X 20 cm was found to be statistically at par with 70 kg N/ha + 55 X 20 cm. The statistical analysis on test weight was found to be significant. However, highest test weight (41.40 g) was recorded with treatment of 70 kg N/ha + 55 X 20 cm and whereas treatment of 60 kg N/ha + 55 X 20 cm, control (41.40 and 40.33 g respectively) was found to be statistically at par with treatment of 70 kg N/ha + 45 X 20 cm.

Significant increase in Capitulum diameter and test weight was favoured by higher levels of nitrogen. "Improved stem elongation and accumulated photosynthates as manifested by higher LAI and dry matter might have been responsible for larger head (Capitulum) diameter. Thus, higher Photosynthetic activity with adequate nitrogen fertilization enabled the plant to accumulate more dry matter and greater translocation of photosynthates to the developing head resulting in larger flower heads" [1]. Similar results were also obtained by Reddy et al. [60], Reddy and Reddy [61] and Sarkar and Mallick [62]. Maximum head diameter and test weight was observed in spacing (55 X 20 cm), it might be due to less competition exerted for light, moisture and nutrients, Sufficient interception of sunlight promotes efficient photosynthesis activities and ultimately greater accumulation of photosynthates under wider spacing.

Narrow spacing with dense plant population resulted in the lower value of yield attributes. The

reduction in yield at with increase in plant density could be attributed to keen competition for moisture, photosynthesis and solar radiation. In wider spacing might be attributed to relatively less inter plant competition because of more space availability to individual plant. Similar results were also obtained by Sen et al. [63], Kumar et al. [64], Ali et al. (2011). The seed and stover yield showed increasing trend with the application of nitrogen in sunflower. The highest seed yield was obtained with the treatment 70 kg N/ha + 45 X 20 cm (2102 Kg), however no other treatment was found to be statistically on par with 70 kg N/ha + 45 X 20 cm. Highest stover yield (4287.9 kg/ha) was recorded 70 kg N/ha + 45 X 20 cm, however, no other treatment was found to be statistically on par with 70 kg N/ha + 45 X 20 cm.

The data showed significant difference in harvest index, however, 70 kg N/ha + 45 X 20 cm recorded highest value of (33.45%) and lowest value (32.81%) was recorded 50 kg N/ha + 65 X 20 cm. "Higher seed and stover yield under higher nitrogen application was due to good growth and availability of adequate nitrogen might lead to increased accumulation of amino acid and amide substance and their translocation to the reproductive organs has improved the seed yield through increased seed setting and filling" [1]. It was evident that plant spacing increased up to 45 X 20 cm showed highest seed and stover yield but increased spacing beyond this decreased the seed and stover yield per plant to noticeable extent. The results are similar with results obtained by Seshadri Reddy et al. [65], Reddi Ramu and Maheswara Reddy [61] and Sarkar and Mallick [62], Kumar et al. [66], Ali et al. (2011).

3.2 Effect on Economics of Sunflower

The observation related to economics were shown in Table 2. Treatment 70 kg N/ha + 45 x 20 cm recorded highest net return (88.737 x 10³ INR / ha), gross return (126.120 x 10³ INR/ha) and benefit: cost ratio (2.37). While lowest gross return (INR 83.644), Net return (INR 47.197) and benefit cost ratio 1.29 was recorded in T₃ with application of 50 kg N/ha + 65 x 20 cm spacing [67-71].

Table 1. Effect of Nitrogen levels and Spacing on yield and yield attributes of Sunflower

S. No	Treatments	Capitulum diameter (cm)	Test weight (g)	Seed yield (Kg/ha)	Stover yield (Kg/ha)	Harvest index (%)
1	50 kg N/ha + 45 X 20 cm	12.70	36.27	1883.35	3830.76	32.96
2	50 kg N/ha + 55 X 20 cm	13.30	38.60	1692.81	3448.24	32.93
3	50 kg N/ha + 65 X 20 cm	12.03	35.22	1394.07	2855.08	32.81
4	60 kg N/ha + 45 X 20 cm	13.67	39.12	2003.72	4067.13	33.00
5	60 kg N/ha + 55 X 20 cm	14.67	40.33	1812.11	3642.39	33.22
6	60 kg N/ha + 65 X 20 cm	12.37	35.52	1514.87	3085.84	32.93
7	70 kg N/ha + 45 X 20 cm	14.03	39.25	2102.00	4287.92	32.89
8	70 kg N/ha + 55 X 20 cm	15.13	41.40	1918.65	3818.31	33.45
9	70 kg N/ha + 65 X 20 cm	13.00	38.36	1638.20	3310.80	33.10
10	Control	14.47	39.81	1646.63	3293.34	33.33
	SEm(±)	0.18	0.64	29.60	59.60	0.04
	CD (5%)	0.53	1.90	87.95	177.07	0.12

Table 2. Effect of Nitrogen and Spacing on economics of Sunflower

S. No.	Treatments	Gross return (x 10 ³ INR/ha)	Net return (x 10 ³ INR/ha)	Benefit : Cost ratio
1	50 kg N/ha + 45 X 20 cm	113.001	75.879	2.04
2	50 kg N/ha + 55 X 20 cm	101.569	64.847	1.77
3	50 kg N/ha + 65 X 20 cm	83.644	47.197	1.29
4	60 kg N/ha + 45 X 20 cm	120.223	82.970	2.23
5	60 kg N/ha + 55 X 20 cm	108.726	71.873	1.95
6	60 kg N/ha + 65 X 20 cm	90.892	54.314	1.48
7	70 kg N/ha + 45 X 20 cm	126.120	88.737	2.37
8	70 kg N/ha + 55 X 20 cm	115.119	78.136	2.11
9	70 kg N/ha + 65 X 20 cm	98.292	61.584	1.68
10	Control	98.798	63.414	1.79

**Economics not subjected to data analysis*

4. CONCLUSION

It was concluded that Treatment of 70 kg N/ha + 55 X 20 cm recorded Maximum plant height (148.48 cm), plant dry weight (60.53 g), test weight (41.40 g), capitulum diameter (15.13 cm) and Number of seeds/capitulum which may be more preferable for farmers .since it is economically more profitable.

ACKNOWLEDGEMENT

I express my gratitude to my advisor Dr. Rajesh Singh for constant support, guidance and for his valuable suggestions for improving the quality of this research work and to all the faculty members of Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh (U.P). For providing all necessary facilities for their cooperation, encouragement and support.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lakshmi Deepika Chandu, Singh R, Singh E. Effect of nitrogen and sulphur levels on growth and yield of sunflower (*Helianthus annuus* L.). The Pharma. Innov J. 2022;11(3):2049-52.
2. Thanki JD, Patel SN, Raman S. Effect of moisture regime in conjunction with nitrogen levels on consumptive use in summer sunflower (*Helianthus annuus* L.). J Oilseeds Resreach. 2002;19(1):44-5.
3. Vavaliya DA, Patel VJ, Patel BD, Panchal PS. Growth and yield of rabi maize (*Zea mays* L.) as influenced by different varieties and crop geometries. Int J Chem Stud. 2018;6(5):2890-3.
4. Vega AJ, Hall AJ. Effects of planting date, genotype and their interactions on sunflower yield determinants of oil-corrected achene yield. Crop Sci. 2002;42(4):1191-201.
5. Vyas AK, Billore SD, Joshi OP, Pachnlania N. Productivity of soybean (*Glycine max.*) genotypes as influenced by nitrogen and sulphur nutrition. Indian J Agric Sceinces. 2006;76(4):272-5.
6. Yadav RP, Tripathi MP, Trivedi SK. Effect of irrigation and nutrient levels on productivity and profitability of sunflower (*Helianthus annuus* L.). Indian J Agron. 2009;54(3):332-225.
7. Özer H, Polat T, Öztürk E. Response of irrigated sunflower (*Helianthus annuus* L.) hybrids to nitrogen fertilization: growth, yield and yield components. Plant Soil Environ. 2004;50(5):205-11. doi: 10.17221/4023-PSE.
8. Pal MS. Effect of plant density on seed yield and harvest index of sunflower in tarai of Uttarachal. Annals Agricultral Res. 2002;23(1):155-8.
9. Palanjiya RR, Solanki RM. Effect of spacing and nitrogen levels on growth, yield and quality of rabi fodder sorghum (*Sorghum bicolor* L.). M.Sc.(Agri.) [thesis] (Published). Junagarh: JAU; 2019.
10. Patel RM, Patel MM, Patel GN. Effect of spacing and nitrogen levels on rabi castor (*Ricinus communis* L.) grown under different cropping sequences in North Gujarat agroclimatic conditions. J Oilseeds Res. 2009;26(2):123-5.
11. Patel SR, Thakur DS. Effect of crop geometry, phosphorus levels and PSB on growth, yield and oil content of Sunflower (*Helianthus annuus* L.). J Oilseeds Res. 2003;20(1):133-56.
12. Patra AK, Mishra A. Effect of variety, nitrogen ad spacing on yield attributes and yield of sesame (*Sesamum indicum* L.) during post-rainy season. J Oilseeds Res. 2000;17(1):113-6.
13. Pavani S. Bhanu rekha K, Sudhakara babu S and Madhu M. Crop Res 45. 2013, Effect of nitrogen and sulphur on growth, yield and quality of sunflower (*Helianthus annuus* L.);152-154:(1, 2 & 3).
14. Poonia KL. Effect of planting geometry, nitrogen and sulfur on growth and yield of sunflower (*Helianthus annuus* L.). J Ecol Physiol. 2000;3:59-71.
15. Prasad PV, Satyanarayama V, Murthy VRK, Boote KJ. Maximizing yields in rice ground cropping sequence through integrated nutrient management. Field Crop Resreach. 2002;75:9-21.
16. Rathod M, Bavalgave VG, Tandel B, Gudadhe NN. Effect of spacing and INM practices on growth, yield and economics of rabi sweet corn (*Zea mays* L.) under south Gujarat condition. Int J Chem Stud. 2018;6(5):247-50.
17. Ravichandran S, Srinivasan N. Effect of sowing density on the growth, yield and nutrient uptake of hybrid sunflower (*Helianthus annuus* L.) in rabi season. Int J

- Curr Microbiol Appl Sci. 2017;6(7):4193-200. doi: 10.20546/ijcmas.2017.607.434.
18. Rasool FU, Hassan B, Jahangir A. Growth and yield of sunflower (*Helianthus annuus* L.) as influenced by nitrogen, sulphur and farmyard manure under temperate conditions. SAARC J Agric. 2013;11(1):81-9. doi: 10.3329/sja.v11i1.18386.
 19. Abbadi C, Sadras A. Response of Sunflower to Nitrogen and Phosphorus in North. 1992;110:1-11.
 20. Steer RA, Beck AT, Garrison B. Applications of the beck depression inventory. Assess Depress. 1986:123-42.
 21. Basha, Allam, Barros. Planting density and nitrogen rates optimization for growth and yield of sunflower (*Helianthus annuus* L.). Hybrids. J Anim Plant Sci. 2012;22(4).
 22. Diepenbrock W, Lang M, Feil B. Yield and quality of sunflower as affected by row orientation, row spacing and plant density. Die Bodenkultur. J Land Manag Food Environ. 2001;52(1).
 23. Kumar VM, Subban P, Singh SSD, Prabhakaran NK. Influence of plant population and N splits on seed, oil yield and economic of rainfed sunflower (*Helianthus annuus* L.). Madras Agriculture [journal]. 2004;91(7-12):365-9.
 24. Ishfaq M, Ali A, Khaliq A, Yaseen M. Allometry, agronomic traits and yield of autumn planted sunflower hybrids under varying row spacing. Pak J Agric Sci. 2009;46(4):0552-9034.
 25. Jahangir AA, Mondal RK, Nada K, Afoze S, Hakim MA. Response of nitrogen and phosphorus fertilizer and plant spacing on growth and yield contributing character of sunflower. Bangladesh J Sci Ind Res. 2006;41(1):33-40. DOI: 10.3329/bjsir.v41i1.258.
 26. Jat RL, Dahama AK, Meena RP. Productivity and economics of mustard (*Brassica juncea* L.) as influenced by phosphorus and sulphur nutrition in association with farmyard manure. Annals Agric Res. 2005;26(4):481-4.
 27. Kar PP, Barik KC, Mahapatra PK, Garnayak LM. Effect of planting geometry and nitrogen on yield, economics and nitrogen uptake of sweet corn (*Zea mays* L.). Indian J Agron. 2006;51(1):43-5.
 28. Kazemeini SA, Edalat M, Shekoofa A. Interaction effects of deficit irrigation and row spacing on sunflower (*Helianthus annuus* L.) growth, seed yield and oil yield. Afr J Agric Res. 2009;4(11):1165-70.
 29. Faith K. Influence of different nitrogen levels on productivity of oil seed and confection sunflower under varying plan populations. Int J Agric Biol. 2004:1560-8530/2004/06-4-594-598.
 30. Kumar SG, Tewari SK, Singh SS. Effect of spacing sources and levels of sulphur on the growth, yield, oil content and economics of sunflower (*Helianthus annuus*) var. "Morden" in sandy loam soil of Allahabad (Uttar Pradesh). Plant Arch. 2012;12(1):249-52.
 31. Kumar AK, Reddy MD. Effect of time of fertilizer application on performance of winter sunflower (*Helianthus annuus* L.). Indian J Agron. 1997;42:512-54.
 32. Riahinia S, Dehdashti SM. Row spacing effects on light extinction coefficients, leaf area index, leaf area index affecting in photosynthesis and grain yield of corn (*Zea mays* L.) and sunflower (*Helianthus annuus* L.). J Biol Sci. 2008;8(5):954-7. doi: 10.3923/jbs.2008.954.957.
 33. Sabale RN. Response of sunflower cultivars to fertilization in rabi season. J Maharashtra Agriculture Univ. 2003;28(3):309-10.
 34. Sawant VB, Sonani VV, Raskar SS, Patil PA. Effect of different levels of nitrogen, phosphorus and sulphur on yield and yield attributes of summer sesame (*Sesamum indicum* L.). Int J Curr Res. 2013;5(10):2744-5.
 35. Sumanthi V, Koteswara Rao DS. Effect of organic and inorganic sources of nitrogen with different irrigation schedules on growth and yield of sunflower (*Helianthus annuus* L.). Indian J Agron. 2007;52(1):77-9.
 36. Syed TH, Gani MR, Ali T, Mi AH. Effect of nitrogen and sulphur fertilization on yield and nutrient uptake by sunflower. J Indian Soc Soil Sci. 2006;54(3):375-6.
 37. Raghavendra S, Sen A, Sharma SN, Singh RK. Effect of plant spacing on Growth, Yield and Seed Quality in restorer Lines of Sunflower Hybrids. Indian J Plant Physiol. 2003;8(4).
 38. Singh B, Mundra SL, Singh D, Kaushik MK, Jat SL. Plant nutrient content and uptake as influenced by plant densities and nutrient management approaches of QPM hybrids. Int J FO Chem Stud. 2018;6(5):2020-2.
 39. Srikanth M, Amanullah MM, Muthukrishnan P, Subramanianik S. Nutrient uptake and yield of hybrid maize (*Zea mays* L.) and soil nutrient status as influenced by plant

- density and fertilizer level. Int J Agric Sci. 2009;5(1):193-6.
40. Taha M, Mishra BK, Acharya N. Effect of irrigation on yield and yield attributing characters of sunflower. Annals Agric Res. 2001;22(2):182-6.
 41. Thavaprakash N, Siva Kumar SD, Raja K, Sentil Kumar G. Effect of nitrogen and phosphorus levels and ratios on seed yield and nutrient uptake of sunflower hybrid DHSI. Pak J Bot. 2004;37:59-68.
 42. Chaniyara NJ, Solanki RM, Bhalu VB. Effect of inter and intra row spacings on yield of mustard. Agric Sci Dig. 2002;22(1):48-50.
 43. Dar EA, Rather SA, Harika AS. Growth and yield of babycorn (*Zea mays* L.) as affected by different crop geometry and level of nitrogen application. Int J Sci Res. 2014;3:7-9.
 44. De Giorgio D, Montemurro V, Fornaro F. Four- year field experiment on nitrogen application to sunflower genotypes grown in semiarid conditions. Eur J Agron. 2007;30:15-26.
 45. Farid SS, Hussain Z, Farida S, Khan S. Performance of sunflower intercropped in ratoon sugarcane under NP fertilizers. Sarhad J Agric. 2000;16(4):431-4.
 46. Gokhale DN, Wadhvane SV, Kalegore NK, Khalgeand S, FG. Response of linseed, (*Linum usitatissimum* L.) varieties to row spacing and phosphorus level under irrigated condition. J Oilseeds Res. 2008;25(1):94-5.
 47. Gunri SK, Biswas T, Mandal GS, Nath R, Kundu CK. Effect of spacing on improved cultivars of summer growing groundnut (*Arachis hypogaea* L.) in red and laterite zone of West Bengal. Karnataka J Agric Sci. 2010;23(5):687-8.
 48. Hada N, Wansik VK, Bhadauria SS, Singh KV. Influence of balanced nutrition, seed rate and plant geometry on fodder maize in south-eastern Rajasthan. Range Manag Agroforestry. 2016;37(2):243-7.
 49. Howlader SH, Bashir HMK, Islam MS, mamun MH, Jahan SMH. Effect of plant spacings on the yield and yield attributes of groundnut. International Journal of sustainable. Crop Poduction. 2009;4(1):41-4.
 50. Kumar S, Tewari SK, Singh SS. Effect of sources and levels of sulphur and spacing on the growth, yield and quality of spring sunflower (*Helianthus annus* L.). 56. 2011;3:242-6.
 51. Mahavishanan K, Prasad M, Bhanu Rekha K. Yield and nutrient uptake of cotton and sunflower as influenced by integrated nutrient management in cotton sunflower cropping system. J Oilseeds Res. 2006;23(1):63-8.
 52. Malik MA, Saleem MF, Sana M, Rehman A. Suitable level of N, P and K for harvesting the maximum economic returns of sunflower (*Helianthus annuus* L.). Int J Agric Biol. 2004;6(2):240-2.
 53. Manjunatha SB, Angadi VV, Palled YB, Hosamani SV. Nutritional quality of multicut fodder sorghum (CoFS-29) as influenced by different row spacing's and nitrogen levels under irrigated condition. Res Environ Life Sci. 2014;7(3):179-82.
 54. Mojiri A, Arzani A. Effect of nitrogen rate and plant density on yield and yield components of sunflower. J Crop Prod Process. 2003;7(2):115-25.
 55. Nagavani AV. Rama Chandra Reddy P. Soundra Rajan, M.S. and Anjaneyulu, A. Growth and yield of sunflower as influenced by irrigation and nitrogen management. Journal of Oilseed Research. 1997;14:315-7.
 56. Nannajkar RR, Sinare BT, Ughade SR. Effect of different varieties and spacing on growth, yield and quality of sunhemp (*Crotalaria juncea* L.). Trends Biosci.9(4). 2016;8431:0974.
 57. Nawaz N, Sarwara G, Yousaf M, Naseeb T, Ahmed A, Shah J. Yield and yield components of sunflower as affected by various NPK levels. Asian J Plant Sci. 2003;2:123-76.
 58. Nawaz R, Ahemad R, Cheema ZA, Mehmood T. Effect of row spacing on sunflower and its weeds. Int J Agric Biol. 2001;3(4):360-2.
 59. Nehra PL, Yadav PS. Integrated weed control in cotton. National conference of Paradigm Shift in cotton Research and Cultivation, October 19-21; 2010.
 60. Reddy MD, Kumar KA, Krishna A. Response of sunflower (*Helianthus annus* L.) to different levels of nitrogen during rabi in northern Telanagana. J Oilseeds Res. 2002;19(1):117-8.
 61. Reddi Ramu Y, Kumar MR P et al. 2004. Ali et al. Growth and yield of sunflower as influenced by nitrogen and sulphur nutrition. Indian J Dry Land Agric Res Dev. 2003. 2011;18(2):192-5.
 62. Sarkar RK, Mallick RB. Effect of nitrogen, sulphur and foliar spray of nitrate salts on

- performance of spring sunflower (*Helianthus annuus*). Indian J Agric Sci. 2009;79(12):986-90.
63. Sen A, Ali SR, Sharma SN, Mishra PJ, Singh AK. Effect of planting density on yield attributes, yield and nutrient uptake of restorer lines of sunflower (*Helianthus annuus* L.). J Oilseeds Res. 2002;19(1):46-9.
64. Kumar J, Badiyala V. Studies on the effect of seed rate, row spacing and sowing time on dry matter accumulation and nutrient uptake in soybean (*Glycine max* L.). J Oilseeds Res. 2004;21(2):290-2.
65. Seshadri Reddy S, Yadahalli YH, Kiran Kumar VK, Kumara O, Boraiah B. Effect of fertilizer, gypsum and boron on dry matter accumulation, yield and nutrient content in sunflower hybrids. Karnataka J Agric Sci. 2002;15(3):569-72.
66. Bastia DK. Response of cotton hybrid Savitha to spacing and NPK treatment under rainfed condition of Orissa. Indian J Agricultural Sci. 2000;70(8):541-2.
67. Abbadi J, Gerendás J. Nitrogen use efficiency of safflower as compared to sunflower. J Plant Nutr. 2009;32(6): 929-45.
68. Ali CK, Nadeem MA, Javid MA. Effect of nitrogen levels and P application strategy on sunflower Department of Fertilizer. 1997;42(3):43-7.
69. Ali H, Randhawa HA, Yousaf M. Quantitative and qualitative traits of sunflower (*Helianthus annuus* L.) as influenced by planting dates and nitrogen application. Int J Agric Biol. 2004;6(2):410-2.
70. Ali M, Khali SK, Nawab K. Response of sunflower hybrids to various levels of nitrogen and phosphorus. Sarhad J Agric. 2000;16(5):477-83.
71. Narwal, Malik. Response of irrigated sunflower (*helianthus annuus*) hybrids to nitrogen fertilization: growth, yield and yield components. Plant Soil Environ. 1985; 50,2004 (5):205-11.

© 2023 Sandhya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97642>