



## **Effect of Probiotics (*Lactobacillus acidophilus* and *Bifidobacterium bifidum*) as Immune Stimulant on Hybrid Catfish *Heteroclarias***

T. A. Yisa<sup>1\*</sup>, O. A. Ibrahim<sup>1</sup>, S. M. Tsadu<sup>1</sup> and U. P. Yakubu<sup>1</sup>

<sup>1</sup>Department of Water Resources, Aquaculture and Fisheries Technology, School of Agriculture and Agricultural Technology, Federal University of Technology, Minna, Niger State, Nigeria.

### **Authors' contributions**

This work was carried out in collaboration between all authors. Author TAY designed the study and wrote the first draft of the manuscript. Author OAI performed the statistical analysis, author SMT wrote the protocol and author UPY managed literature searches. Authors TAY, OAI, SMT and UPY managed the analyses of the study and literature searches. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/BMRJ/2015/17703

#### Editor(s):

(1) Marcin Lukaszewicz, Department of Biotransformation, Faculty of Biotechnology, University of Wroclaw, Wroclaw, Poland and Division of Chemistry and Technology Fuels, Wroclaw University of Technology, Wroclaw, Poland.

#### Reviewers:

- (1) Anonymous, Ocean University of China, China.  
(2) M. Angels Calvo Torras, Univeristat Autònoma de Barcelona, Spain.  
(3) Anonymous, Malaysia.

Complete Peer review History: <http://www.sciencedomain.org/review-history.php?iid=1216&id=8&aid=9674>

**Original Research Article**

**Received 24<sup>th</sup> March 2015**  
**Accepted 25<sup>th</sup> May 2015**  
**Published 9<sup>th</sup> June 2015**

### **ABSTRACT**

Study on the effect of probiotics (*Lactobacillus acidophilus* and *Bifidobacterium bifidum*) as feed supplement and immune stimulant on hybrid *Heteroclarias* was carried out with 240 hybrid *Heteroclarias* fingerlings with mean weight  $1.90 \pm 0.40$  g fed probiotic supplement (GNC Ultra 25) for fifty six (56) days in 12 number 30 x 60 x 30 cm rectangular aquarium tanks. There were four treatments and each replicated three times. The inclusion level of probiotic fed supplement was at 0 g/kg, 1 g/kg, 2 g/kg and 3 g/kg. The experimental fish were fed at 3%, 5%, 7%, and 9% body weight at two weeks interval, The result showed that there was significant difference  $p(<0.05)$  between control and other treatments in terms of specific growth rate and percentage survival. The microbial analysis showed that there was presence of Gram positive bacterial in the gut of the hybrid *Heteroclarias*. The treatments fed with probiotics have the higher plate count as compared to the control, the haematological analysis showed that blood parameters of control differed

\*Corresponding author: E-mail: [atyisa@yahoo.com](mailto:atyisa@yahoo.com);

significantly  $p(<0.05)$  from other treatments. The inclusion levels of the probiotic supplement use in this study enhanced the growth and immune response of hybrid *Heteroclarias*, however inclusion level at 1 g/kg is sufficient to enhance the proliferation of the required bacteria.

**Keywords:** *Heteroclarias*; immune booster; fish feed (coppens); culture media and *Lactobacillus acidophilus* and *Bifidobacterium bifidum*.

## 1. INTRODUCTION

Probiotics are microbes which have a beneficial effect on their host by modifying the host ambient microbial community, by ensuring improved use of the feed or enhancing its nutritional value, *visa-viz* the host response towards disease, and improving the quality of its ambient environment [1]. This implies that probiotics may include microbial additives that prevent pathogenic organism from proliferating or establishing in the gut, on the superficial structure and in culture environment of the cultured species that ensure optimal use of feed by aiding digestion, improve water quality and stimulate the immune system of the host [2]. They also stated that there is suggestive evidence that several probiotic strains such as *Lactobacillus acidophilus* are useful in boosting the immune response.

They are also selected pure culture isolates of one or more living microorganisms added to feed to proliferates in the host gastrointestinal (GI) tract [3]. They confer a healthy benefit on the host as significant microbial feed supplement in the field of prophylaxis [4]. The research on probiotics for animal health is increasing with the demand for environment-friendly microorganism. Some probiotics were designed to treat the rearing medium, like bio control when the treatment is antagonistic to pathogenic organism or bioremediation when water quality is improved to support the aquatic organism. Most probiotics have been studied by isolating and selecting strains from aquatic environment [5]. For example *Bifidobacterium* spp Gram positive, anaerobic, branched or pleomorphic rods can be isolated from a variety of materials such as human and animal faces, sewage and from the oral cavity [5]. There are wide range of micro algae, yeast (*Debaryomyces*, *Phaffia* and *Saccharomyces*) and Gram positive (*Bacillus*, *Lactococcus*, *Micrococcus*, *Carnobacterium*, *Enterococcus*, *lacobacillus*, *Streptococcus*, *Weisslla*) and gram negative bacteria (*Aeromonas*, *Alteromonas*, *Photorhodobacterium*, *Pseudomonas* and *Vibrio*) that have been evaluated and known to be probiotics [5].

The use of antibiotic to treat or control disease in aquaculture have been reported to affect the fish intestinal microbiota [6,7] and cause prevalence of fish resistance to antibiotic [8,9]. This notorious adverse effect associated with the use of antibiotic in aquaculture has made European Union to ratify a ban for the use of all sub-therapeutic antibiotics as growth promoting agents or prophylaxes in animal production. There is an urgent need in aquaculture to look for alternative to antibiotic in disease control; therefore, probiotics seems to prefer a solution and is increasingly viewed as an alternative to antibiotics in animal production and particularly aquaculture [10]. One of the common probiotics is *Lactobacillus and bifidobacterium sp* which are a Lactic Acid Bacteria (LAB) [11].

Although the idea of prebiotics and probiotics were used in animals for the purpose of enhancing growth and reducing early mortality [12]; in humans their potential nutritional advantages consist of preventive and curative effects against diseases, including intestinal dysfunctions, inflammatory bowel diseases and colon cancer amongst others [12,13]. In Nigeria apart from the problems of high cost of fish feeds and quality seed, disease outbreak is a major challenge in fish farming particularly in early life stages [14]. This study was carried out to evaluate the effect of *Lactobacillus and bifidobacterium sp* probiotics drug *GNC ultra 25* as feed supplement when incorporated into fish feed and immune stimulant on hybrid *Heteroclarias*. The choice of *Heteroclarias* in this study was due to its high market value, fast growth rate, improved hybrid vigour and resistant to disease amongst others.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Site

The experiment was carried out at Toxicology unit, Fish Farm, Department of Water Resources, Aquaculture and Fisheries Technology, Bosso Campus, Federal University of Technology, Minna, for 56 days.

## 2.2 Experimental Fish and Feed

Two hundred and forty (240) *Heteroclarias* fingerlings of average (Mean initial weight  $1.9 \pm 0.4$  g) were purchased from private commercial fish farm in Ilorin, Kwara State. The fish were acclimatized for 5 days and fed with commercial diet (0.6-0.8 mm size coppers). Twenty (20) fish samples were randomly selected and distributed in 12 aquarium tanks of 30 liters water holding capacity, and each of the tanks was filled with approximately 15 L of clean water. Four treatments were maintained and each treatment was replicated three times. The tested *ultra GNC 25* probiotic drug (*Lactobacillus acidophilus* and *Bifidobacterium bifidum*) was added at inclusion level of 0 g, 1 g, 2 g and 3 g/kg diet. This was then mixed thoroughly and stored in a cool dry place. The experimental fish were fed at 3%, 5%, 7%, and 9% body weight at two weeks interval. Specific Growth Rate (SGR), Feed Conversion Ratio (FCR) and cumulative percentage survival and mortality rates was determined using the formulae below:

$$\text{SGR} = \frac{\text{Log final weight} - \text{Log initial weight}}{T_2 - T_1} \times 100$$

Log = Natural logarithm,  $T_2$  = Time two and  $T_1$  = Time one.

$$\text{FCR} = \frac{\text{Total feed consumed by fish (g)}}{\text{Total weight gain by fish (g)}}$$

Percentage Mortality=

$$\frac{\text{Cumulative Mortality} \times 100}{\text{Total number stocked}}$$

$$\text{Percentage Survival} = \frac{\text{Cumulative Survival} \times 100}{\text{Total number stocked}}$$

## 2.3 Standard Plate Count (SPC) and Gut Sampling

Intestinal content of *Heteroclarias* was taken from 40 fish sampling and total plate count were enumerated using method described by Onwuka, [15]. Isolation of bacteria from *Heteroclarias* intestinal content after oral administration of probiotics for microbial count was carried out by taking one fish per replicate from each treatment, killed and gutted. Selective media (solid and broth media) was adopted according to method of Nebra and Blanch, [16]. Their gut content was obtained using a swab stick which was immediately transferred to 1 ml of 0.09% saline

solution, thereafter homogenized in sterile distilled water. The homogenized sample was diluted serially up to ( $5 \times 10^5$ ) fold in sterile water and then inoculated on nutrient agar plates by pour plate method as described by Awan and Rahman, [17]. The inoculated plate was incubated at 37°C for 24 hours; the number of visible colonies was counted. The results were typically expressed as colony forming units (C.F.U.)/g. or /ML.

## 2.4 Haematological Examination

### 2.4.1 Blood sampling

5 ml of blood sample were collected from the experimental fish by severing them at the tail region and dripped blood samples into clean EDTA bottle. The blood samples were analyzed for Red Blood Cell (RBC), White Blood Cell (WBC) and Pack Cell Volume (PCV) according to the methods of (Kaplow, 1955) as described by Onwuka, [15]. Haematocrit value was determined by the standard micro haematocrit method, and expressed in percentage. Duplicate blood samples were loaded into standard heparinized capillary tubes, spun in a micro haematocrit centrifuge at 12,000 rpm for 5 minutes and measured on a haematocrit reader.

## 2.5 Experimental Design and Statistical Analysis

Completely Randomized Design (CRD) was used for the experiment. The data obtained were subjected to one way analysis of variance (ANOVA) and all differences in mean values of parameters were determined at  $P = 0.05$  level of significance. The coefficient regression equation was used to determine the length/weight relationships. Also Duncan Multiple range Test was used for mean separation.

## 3. RESULTS

The result in Table 1 shows that there was no significant difference ( $p > 0.05$ ) of mean weight gain and feed conversion ratio among all treatments in this study. There was a significant difference ( $p < 0.05$ ) in treatment I (0 g inclusion level probiotic drug) compared to others in terms of specific growth rate and percentage of survival. The result in Table 2 shows isolation and identification of bacterial organisms from *Heteroclarias* gut (stomach) fed probiotics inclusion level at 0, 1, 2 and 3 g/kg respectively.

It shows that standard plate count (SPC) differed significantly ( $p < 0.05$ ) among the treatments. The result of haematological parameters tested in the blood sample of experimental fish is indicated in Table 3. Result indicated shows that the red blood cell (RBC), white blood cell (WBC), pack cell volume (PCV) and haemoglobin (Hb) differed significantly ( $p < 0.05$ ) in between treatment I compared to other treatments. The result in Table 4 above shows that gram positive cocci was present in all the treatments as gram negative cocci were absent in all. However, gram positive rod was absent in treatment I as gram negative rod was present.

#### 4. DISCUSSION

The least survival rate in the control could be attributed to the absence of beneficial microorganisms in the intestinal tract because the ability to control the micro flora in the intestinal tract with beneficial microorganisms was triggered by probiotics which increased the survival rate of *Heteroclaris* (Table 1). The combined effect of *Lactobacillus acidophilus* and *Bifidobacteria bifidum* used in this study as immune stimulant revealed to have enhanced the growth and resistance to disease of *Heteroclaris*, hence the higher percentage

**Table 1. Growth response of *Heteroclaris* fingerlings Fed with probiotics at various inclusion levels for 56 days**

Growth parameters	Treatment I	Treatment II	Treatment III	Treatment IV
Mean initial weight (MIW(g))	1.88 <sup>a</sup> ±0.75	1.86 <sup>a</sup> ±0.05	1.79 <sup>a</sup> ±0.38	1.87 <sup>a</sup> ±0.04
Mean final weight (MFW(g))	15.84 <sup>b</sup> ±0.27	20.67 <sup>a</sup> ±0.82	19.92 <sup>a</sup> ±0.24	20.10 <sup>a</sup> ±0.96
Mean weight gain (MWG(g))	12.87 <sup>b</sup> ±0.27	18.48 <sup>b</sup> ±0.86	17.85 <sup>b</sup> ±0.32	17.89 <sup>b</sup> ±1.10
Feed conversion ratio (FCR)	0.65 <sup>a</sup> ±0.02	0.64 <sup>a</sup> ±0.06	0.57 <sup>a</sup> ±0.04	0.62 <sup>a</sup> ±0.41
Specific growth rate (SGR(%/day))	3.80 <sup>b</sup> ±0.12	4.29 <sup>a</sup> ±0.14	4.29 <sup>a</sup> ±0.08	4.29 <sup>a</sup> ±0.16
% Survival rate	63.33 <sup>b</sup> ±2.8	85 <sup>a</sup> ±5.00	86.66 <sup>a</sup> ±2.88	85 <sup>a</sup> ±5.00

Legend: Treatment I=0 (g) probiotic per 1 kg diet; Treatment II= 1 (g) probiotic per 1 kg diet; Treatment III= 2 (g) probiotic per 1 kg diet and Treatment IV =3 (g) probiotic per 1 kg diet.

Mean data on the same row carrying different superscript differs significantly ( $p < 0.05$ ) from each other.

**Table 2. Microbial count of gut sample of *Heteroclaris* fingerlings fed probiotics at various inclusion levels for 56 days**

Microbial count	Treatment 1	Treatment II	Treatment III	Treatment IV
Colony forming unit (CFU)	0.8 <sup>a</sup> ×10 <sup>5</sup> ±0.70	1.0 <sup>b</sup> ×10 <sup>5</sup> ±0.36	1.2 <sup>c</sup> ×10 <sup>5</sup> ±0.16	1.1 <sup>c</sup> ×10 <sup>5</sup> ±0.63

Mean data on the same row carrying different superscripts differs significantly ( $p < 0.05$ ) from each other

**Table 3. Haematological parameter of *Heteroclaris* fingerlings fed probiotic at various inclusion levels for 56 days**

Blood Parameters	Treatment I	Treatment II	Treatment III	Treatment IV
Red blood cell (RBC(10 <sup>6</sup> mm <sup>-3</sup> ))	1.01 <sup>a</sup> ±0.03	1.08 <sup>b</sup> ±0.08	1.08 <sup>b</sup> ±0.03	1.07 <sup>b</sup> ±0.08
White blood cell count (WBC( 10 <sup>3</sup> mm <sup>-3</sup> ))	14.98 <sup>a</sup> ±0.24	16.67 <sup>b</sup> ±0.44	19.14 <sup>c</sup> ±0.21	20.10 <sup>c</sup> ±0.30
Pack cell volume (% PCV)	24.24 <sup>a</sup> ±0.61	26.77 <sup>b</sup> ±0.41	26.91 <sup>b</sup> ±0.20	26.95 <sup>b</sup> ±0.24
Haemoglobin (Hb( g/100ml))	8.96 <sup>a</sup> ±0.39	10.11 <sup>b</sup> ±0.20	11.80 <sup>b</sup> ±0.15	11.66 <sup>b</sup> ±0.33

Mean data on the same row carrying different superscript differed significantly ( $p < 0.05$ ) from each other

**Table 4. Gram staining for the culture of the gut content of *Heteroclaris* fed probiotic at various inclusion levels for 56 days**

Bacteria Species	Treatment I	Treatment II	Treatment III	Treatment IV
Gram positive cocci	present	present	present	present
Gram positive rod	absent	present	present	present
Gram negative rod	present	absent	absent	absent
Gram negative cocci	absent	absent	absent	absent

survival in treatments II, III and IV. Similarly, [18] observed that a strain of *Lactococcus rhamnosus* administered to rainbow trout enhanced survival rate and stimulates the respiratory boost activity after 2 weeks of feeding. The presence of gram positive rod shaped bacteria in *Heteroclaris* fed probiotics led to the establishment of a normal gut micro biota which add to the effective function of digestive system and serve as a barrier against invading pathogens as stated by Farzanfar, 2006 [19]. The high value of haematological parameters observed in the treatments fed probiotics on *Heteroclaris* showed that probiotics actively stimulate the proliferation of lymphocytes and immunoglobulin production in fish as similar observation was made by [20].

## 5. CONCLUSION

From the research, it could be concluded that probiotics (*Lactobacillus acidophilus* and *Bifidobacterium bifidum*) fed *Heteroclaris* at inclusion level of 1 g/kg diet was sufficient to stimulate immune and proliferate beneficial microorganisms in the gut.

## 6. RECOMMENDATION

Probiotics should be incorporated in commercial feeds at least 1 g/kg diet by fish farmers because of its ability to improve gut health and stimulating immune function.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Fuller R. Probiotics in man and animals. Journal of Applied Bacteriology. 1989;66: 365-378.
- Lee MC, Lin LH, Hung KL, Wu HY. Oral bacteria therapy promotes recovery from acute diarrhea in children. Acta Paediatr. 2001;42(30):1-5.
- Lin C. Progression of intensive marine shrimps culture in Thailand in swimming through troubled water. Proceeding of the special program on shrimp farming. Aquaculture Research. 2005;13-23.
- Balcazar JL, De Blas I, Ruiz-Zarzuola I, Vendrell D, Muzquiz JL. Probiotics a tool for the future of fish and shellfish health management. Journal of Aquaculture in the Tropics. 2004;19:239-242.
- Gatesoupe FJ. The use of probiotics in aquaculture. Aquaculture. 1999;180:147-165.
- Strom E, Ringo E. Changes in the bacteria composition of early developing cod, *gadus morhua* larva following inoculation of *Lactobacillus plantarum* into the water in physiological and biochemical aspect of fish development. Eds walter BT, fyhn HJ. (University of Bergen, bergen, Norway). 1993;226-228.
- Direkbusarakom S, Yoshimizu M, Ezura Y, Ruangpan L, Danayadol Y. The dominant flora in shrimp hatchery against some fish pathogenic viruses. Journal Marine Biotechnology. 1998;6:266.
- Cabello FC. Heavy use of prophylactic antibiotics in aquaculture, a growing problem for human and animal health and for the environment. Journal of Environmental Microbiology. 2006;8:1137-1144.
- Serum H. Antimicrobial drug resistance in fish pathogen In: Aarestrup FM. (ed). Antimicrobial resistance in bacteria of animal origin. ASM Press Science. 2006; 6:75-82.
- Verschuere L, Rombaut G, Sorgeloos P, Verstraete W. Probiotic bacteria as biological control agents in aquaculture. Journal of Microbiology and Molecular Biology. 2000;64:655.
- Aly SM, Abd-El-Rahman AM, John G, Mohammed MF. Characterization of some bacteria isolated from *Oreochromis niloticus* and their potential use as probiotics in aquaculture. Aquaculture Research. 2008a;277:1-6.
- Chabrillo'n M, Rico RM, Balebona MC, Morinigo MA. Adhesion to sole, *Solea senegalensis*, mucus of micro-organisms isolated from farmed fish, and their interaction with photo bacterium. Journal of Fish Diseases. 2005;28:229.
- Lara-Flores M, Aguirre-Guzman G. The use of probiotic in fish and shrimp aquaculture. Aquaculture Research. 2009; 44:456-564.
- Servin AL, Coconnier MH. Adhesion of probiotic strains to the intestinal mucosa and interaction with pathogens. Best Pract Res Clin Gastroenterol. 2003;17:741\_54.
- Onwuka GI. Food analysis and instrumentation. Adeniyi Jones Close

- Surulere, Lagos, Naphthali Publications. 2005;171-174.
16. Nebra Y, Blanch AR. A new selective medium for *Bifidobacterium* spp. Journal of Applied and Environmental Microbiology. 1999;65(11):5173-5176.
  17. Awan JA, Rahman SU. Microbiology manual Uni. tech communication, Faisalabad Pakistan. 2005;10:49-51.
  18. Nikoskelainen S, Ouwehand AC, Bylund G, Salminen S, Lilius EM. Immune enhancement in rainbow trout (*Oncorhynchus mykiss*) by potential probiotic bacteria (*Lactobacillus rhamnosus*). Shellfish Immunology. 2003; 15:443-452.
  19. Farzanfar A. The use of probiotics in shrimp aquaculture. Journal of Immunology and Medical Microbiology. 2006;48:149-158.
  20. Al-Dohail MA, Hashim R, Aliyu-Paiko M. Effects of the probiotic, *Lactobacillus acidophilus*, on the growth performance, haematology parameters and immunoglobulin concentration in African Catfish (*Clarias gariepinus*, Burchell, 1822) fingerling. Aquaculture Research. 2009;40: 1642-1652.

© 2015 Yisa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<http://www.sciencedomain.org/review-history.php?iid=1216&id=8&aid=9674>