



# **Surgical Repositioning of Developing Canine and Circummandibular Wire Fixation of Parasymphysis Fracture in a Four Year Old Child: A 1 - Year Follow up Case Report**

**K. Maheshkumar<sup>a++\*</sup>, Nagalakshmi Chowdhary<sup>a#</sup>,  
Maheshkumar<sup>b#</sup>, Rajashekar Reddy<sup>a†</sup>, Supriyo Pal<sup>c‡</sup>  
and S. Varsha<sup>c++</sup>**

<sup>a</sup> Department of Pedodontics and Preventive Dentistry, Sri Siddhartha Dental College and Hospital, Sri Siddhartha Academy of Higher Education, Tumakuru, 572107, Karnataka, India.

<sup>b</sup> Department of Oral Maxillofacial Surgery, Sri Siddhartha Dental College and Hospital, Sri Siddhartha Academy of Higher Education, Tumakuru, 572107, Karnataka, India.

<sup>c</sup> Department of Oral and Maxillofacial Surgery, Sri Siddhartha Dental College, Tumkur, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102303>

**Case Report**

**Received: 20/05/2023**

**Accepted: 23/07/2023**

**Published: 31/07/2023**

## **ABSTRACT**

Mandibular fractures are the most common (56%) facial skeletal injury with a prevalence of 0.6-1.4% below the age of five. This is attributed to the constant supervision of parents and is seen to increase as child begin school peaking during puberty and adolescence due to increased sport and unsupervised physical activity.

<sup>++</sup> Post Graduate Student;

<sup>#</sup> Professor and H.O.D;

<sup>†</sup> Reader;

<sup>‡</sup> Senior Lecturer;

\*Corresponding author: E-mail: kmaheshkumarbds@gmail.com;

The goal while treating Paediatric fractures is to ensure bony union, normal occlusion, restore normal form and function, and avoid impediments to normal growth. Management of mandibular fractures in paediatric patients depends upon the fracture type, site of fracture & the phase of dental and skeletal development.

The Authors managed to treat a 4-year old Parasymphyseal fracture associated with inversion of permanent canine tooth bud using surgical repositioning and Circummandibular wire fixation.

**Keywords:** Facial trauma; mandibular fracture; circummandibular-wire fixation; displaced tooth bud.

## 1. INTRODUCTION

In hospitalised paediatric trauma patients, mandibular fractures are the most common (56%) facial skeletal injury [1]. Their prevalence is low below the age of five (0.6-1.4%) and increases as children begin schooling, with peaks during puberty and adolescence, with a predilection for boys, owing to increased sport and unsupervised physical activity [2-3]. The etiological factors reported are motor accidents, sports injuries, falls and victims of child abuse [1].

The most concerning aspect for while treating paediatric patients is the effect of trauma/treatment on growth and development, which differs from adults [4]. Other factors affecting growth and development are anatomic and physical factors which are equally significant, and can have diverse effects on management [5]. Frontal prominence reduces in size making the facial bones to emerge from the shelter of the cranial base [3,6]. Pediatric maxillofacial complex is influenced greatly by cancellous-to-cortical ratio and has higher osteogenic and bone remodeling potential [7].

This article, presents effective management of Paediatric parasymphysis fractures associated with traumatic inversion of permanent canine tooth bud by circummandibular wire fixation.

## 2. CASE REPORT

A 4-year-old female child reported to the Outpatient Department of Pedodontics and Preventive Dentistry, with a chief complaint of pain in her lower left face. On eliciting history, a fall from travelling autorickshaw was reported following which she sustained injuries to her face.

On extra oral examination, multiple wounds over the left face was noted with a solitary ovoid swelling associated with step deformity at the symphyseal region measuring approximately 2cm in diameter which was tender on palpation.

Intraoral examination revealed deranged occlusion and segmental mobility noted between 71 -72 region and associated vestibular tenderness (Fig. 1). On radiographic examination, inversion of 33 tooth bud along the fracture line was noted (Fig. 2). Based on the above finding, a diagnosis of left parasymphyseal fracture with traumatic inversion of permanent canine tooth bud was made.

Informed consent was taken from the parent before starting the treatment. Impressions of maxilla and mandible were taken for the fabrication of closed acrylic cap splint. Under general anaesthesia, the fracture segments were carefully exposed and displaced to facilitate for instrumentation of surgical repositioning of the tooth bud in 33. (Figs. 3,4) Closure of the wound was done using 3-0 Vicryl suture and fracture segments were later reduced and held in place by means of circummandibular wire fixation using the cap splints (Fig. 5).

Postoperatively, patient was continued on analgesic and antibiotic regimen. Soft diet, avoidance of physical activities, and antibacterial mouth rinse were prescribed. No signs of complications were observed during the healing period. Post operative radiograph shows the successful reposition of the 33 tooth bud and reduction of fracture segments. (Fig. 6) 1-year follow-up shows the unhindered eruption pattern of the 33tooth bud (Fig. 7).

## 3. DISCUSSION

Maxillofacial fractures in the pediatric population comprise of less than 15% of all facial fractures. Numerous studies have reported almost similar data of a decreased occurrence of mandibular fractures in children when compared to adults, ranging from 1% to 15%[2,3,8].

Management of mandibular fractures in pediatric patients depends upon the type of fracture and the phase of dental as well as skeletal development. Main concerns while treating the

pediatric mandible fracture are mandibular growth and development of dentition. Factors that can increase the risk of managing mandibular fractures are small jaw size, the presence of developing permanent tooth buds, and existing active growth centres [6].

To be noted is that children have a higher osteogenic potential and a rapid healing rate when compared to adults. Thus, anatomic reduction must be achieved earlier and immobilization periods must have a lesser time period (two weeks instead of four–six weeks for adults)[2,3,9]. Based on the type of fracture and the patient's stage of development, immobilization and fixation of the fracture segments can be achieved by means of maxillomandibular fixation (MMF) or internal skeletal fixation or a combination of both these methods [2,3,5].

Intermaxillary fixation (IMF) using the teeth in pediatric facial fracture patient may prove to be more difficult than in adults. This may be due to decreased availability of teeth, resorption of roots of deciduous teeth, surfaces of the teeth not being retentive for etching procedure, and unfavourable form of the crowns of deciduous teeth for the fixation of interdental wires and arch bars [3,10,11]. Posnick stated that approximately 42% of mandibular fractures in his series were managed by closed reduction, specially with the help of maxillomandibular fixation (MMF) [12].

In present times, open reduction and internal fixation (ORIF) have become the gold standard for the treatment of displaced pediatric mandible fractures [3,4,10,11]. This procedure uses fixation with miniplates, microplates, or bio-degradable plates. Although ORIF provides three-dimensional stability, promotes primary healing, and shortens the treatment time, several risks are associated with the ORIF for the management of the pediatric mandibular fracture such as damage to the developing tooth buds; however, in a specific age group, plate fixation is possible at the inferior border of the mandible away from the developing tooth germs [10,13]. In addition this also carries the risk of interference with growth, plate migration and stress shielding owing to the placement of the plates. Allergic reaction to the metal leads to inflammatory sequelae, which necessitates further elimination of the plating hardware. Corrosion and freeing of metal ions can also be a cause to avoid the internal fixation devices [10].


At the same time this case shows, surgical repositioning is a treatment option for displaced developing teeth germ to avoid dilacerations with subsequent impaction. Other methods for the treatment of an impacted permanent tooth are surgical exposure followed by orthodontic extrusion or the extraction of the impacted tooth [14,15]. Kuroe et al. reported that an early time of operation favors a better ability to mobilize the tooth, and the risk of damaging the developing root is smaller[16]. Any damage to the periodontal ligament during the operation can result in root resorption and/or ankylosis or in arresting of further root development. As this case shows, a repositioning is also possible even if root development has not started. Therefore the risk of damaging the root is even more reduced [16,17].

Surgical repositioning is a fast and cost effective treatment alternative for displaced permanent tooth germs and can be successful in avoiding later malformation of the permanent teeth [18].


Complications such as postoperative infection, non-union, and malunion are less common when closed reduction is done properly, due to the higher osteogenic potential, rapid healing rate, and less common necessity for open reduction and rigid internal fixation. In addition, a large number of fractures are minimally displaced to undisplaced. However, TMJ dysfunction, restricted condylar translation, deviation upon mouth opening and growth disturbances like hypoplasia of mandible or asymmetry, and secondary midface deformity usually occur with severe comminuted fractures in pediatric patients [3,10].




**Fig. 1. Fracture seen in between in 71-72 region**



**Fig. 2. Preoperative panoramic radiograph showing fracture line and inversion of permanent tooth germ**




**Fig. 3. The site of fracture was expanded to gain access to 33 tooth bud**



**Fig. 4. The displaced tooth bud (33) was repositioned**






**Fig. 5.** Repositioned and the site of fracture was reduced to normal anatomical site and tied ss wire



**Fig. 6.** One year follow up



**Fig. 7.** One year follow up OPG shows reduction of fracture segments and unhindered eruption pattern of the 33tooth bud

#### **4. CONCLUSION**

In conclusion close reduction is preferred in mandibular fractures in a young child, whenever

intervention is needed. In a younger pediatric patient, an acrylic splint fixed to the mandible with the help of circummandibular wire fixation can successfully eliminate the need of IMF.

## CONSENT

As per international standard or university standard, Parental written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Priya Vellore K, Gadipelly S, Dutta B, Reddy VB, Ram S, Parsa A. Circummandibular wire fixation of symphysis fracture in a five-year-old child. *Case Rep Dent*. 2013;2013:930789.
2. Glazer M, Joshua BZ, et al. Mandibular fracture in children: Analysis of 61 cases and review of the literature. *Int J Pediatr Otorhinolaryngol* 2011;75(1):62–64. DOI: 10.1016/j.ijporl.2010.10.008
3. Zimmermann CE, Troulis MJ, et al. Pediatric facial fractures: Recent advances in prevention, diagnosis and management. *Int J Oral Maxillofac Surg*. 2006;35:2–13.
4. Myall RW. Management of Mandibular fractures in children. *Oral Maxillofacial Surg Clin North Am*. 2009;21(2):197–201.
5. Haug HR, Foss J. Maxillofacial injuries in the pediatric patient. *Oral Surg Oral Med Oral Pathol*. 2000;90(2):126–134.
6. Sharma A, Patidar DC, Gandhi G, Soodan KS, Patidar D. Mandibular Fracture in Children: A New Approach for Management and Review of Literature. *Int J Clin Pediatr Dent*. 2019;12(4):356-359.
7. Aziz SR, Ziccardi VB. Management of Pediatric Facial Fractures. Fonseca RJ. *Oral and Maxillofacial Trauma*, 4<sup>th</sup> ed., Missouri: Elsevier St. Louis; 2013;735–748.
8. Singhal R, Singh V, et al. Pediatric maxillofacial injuries- If a new look is required? *Int J Pediatr Otorhinolaryngol*. 2013;77:1333–1336.
9. Cole P, Kaufman Y, et al. Principles of pediatric mandibular fracture management. *Plast Reconstr Surg*. 2009;123:1022–1024.
10. John B, John RR, et al. Management of mandibular body fractures in pediatric patients: A case report with review of literature. *Contemp Clin Dent*. 2010;1(4): 291–296. DOI: 10.4103/0976-237X.76406
11. Joshi S, Kshirsagar R, et al. Clinical efficacy of open reduction and semirigid internal fixation in management of displaced pediatric mandibular fractures: A series of 10 cases and surgical guidelines. *J Indian Soc Pedod Prev Dent*. 2015;33(2): 161–165.
12. Posnick JC. Management of facial fractures in children and adolescents. *Ann Plast Surg*. 1994;33(4):442-457.
13. Infante Cossio P, Espin Galvez F, et al. Mandibular fractures in children. A retrospective study of 99 fractures in 59 patients. *Int J Oral Maxillofac Surg*. 1994; 23:329–331. DOI: 10.1016/s0901-5027(05) 80047-5
14. Andreasen JO, Petersen JK, Laskin DM. *Textbook and Color Atlas of Tooth Impactions: Diagnosis, Treatment, Prevention*, 1<sup>st</sup> edn. Copenhagen: Munksgaard; 1997.
15. Kim S, Kim J, Song JS, Choi H, Choi B, Kim S. Continued root development of a surgically repositioned human incisor tooth germ. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2013;115:e11–e15.
16. Kuroe K, Tomonari H, Soejima K, Maeda A. Surgical repositioning of a developing maxillary permanent central incisor in a horizontal position: Spontaneous eruption and root formation. *Eur J Orthod*. 2006;28: 206–209.
17. Tsai T. Surgical repositioning of an impacted dilacerated incisor in mixed dentition. *J Am Dent Assoc*. 2002;133: 61–66.
18. Spierri A, Signorelli C, van Waes H. Surgical repositioning of a developing maxillary central incisor. A case report. *Int J Paediatr Dent*. 2015;25(4):305-308.

© 2023 Maheshkumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/102303>