



A Case Report on Virilizing Adrenal Tumor

Shikhil Puzhakkal ^a^o, Niya Narayanan ^b[#], Ani Praveen ^c[#]
and Pradeep Puthen Veetil ^a^{¥*}

^a Endocrine Surgery Department, Baby Memorial Hospital, Kozhikode, Kerala, India.
^b Medical Endocrinology Department, Baby Memorial Hospital, Kozhikode, Kerala, India.
^c Pathology Department, Baby Memorial Hospital, Kozhikode, Kerala, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author SP designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors NN and AP managed the analyses of the study. Author PPV managed the literature searches. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93339>

Case Study

Received: 12/09/2022

Accepted: 03/11/2022

Published: 07/11/2022

ABSTRACT

Objective: To discuss a rare case of virilizing Oncocytoma of Adrenal Gland.

Background: Hirsutism is a common problem affecting 5-10% of the general population. It causes immense psychological burden. The commonest cause of hirsutism is poly cystic ovary syndrome (PCOS). It is seen in 3 out of 4 hirsute women. Sometimes it will be due to rare endocrine diseases like Cushing's syndrome or non- classical congenital adrenal hyperplasia or androgen secreting ovarian tumor and adrenal tumors.

Presentation: we report a 49 Year old lady who presented with hirsutism and virilization features to the Dermatology department. On further hormonal evaluation, she was found to have an elevated Di hydro epiandrosterone - sulfate (DHEA-S). Computerized Tomography (CT) of abdomen was done which showed 45x35x38 mm right adrenal mass. She underwent open right adrenalectomy

^o Associate Consultant;

[#] Consultant;

[¥] Senior Consultant;

*Corresponding author: E-mail: pradeepputhenveetil@yahoo.co.in, pradeepputhenveetil@yahoo.co.in;

and histopathology was suggestive of adrenal oncocytoma, a rare adrenal tumor. Functional adrenal oncocytoma is very rare in literature and it is valuable to report.

Conclusion: Adrenal oncocytomas are rare tumors which are commonly present as adrenal incidentalomas. Only histopathology can diagnose oncocytomas and its clinical behaviour. Treatment of choice is excision as open or laparoscopic adrenalectomy. As there are no clear cut guidelines for follow up, it needs further studies and high levels of evidence.

Keywords: Hirsutism; virilization; DHEA-S; adrenal oncocytoma.

1. INTRODUCTION

Oncocytic adrenocortical tumors (OATs) or neoplasms are extremely rare. Most of the cases are nonfunctional and benign, predominantly affecting middle-aged women. About 147 cases have been reported globally in the literature according to a systematic review by Mearini et al. and it is reported that most of the affected patients have been in the age group of 40 to 60 years. Most of these tumors were detected incidentally with 20% of them demonstrating evidence of malignancy. Approximately 10-20% of the OATs were functional in nature. Here we present a 49 year old lady with a functional OAT.

2. CASE PRESENTATION

A 49 year old lady with no co-morbidities came with history of excessive hair growth over chin and upper lip area of 8 months duration which was progressive in nature. She had history of recent change in voice for 6 months. She attained menopause 1 year back till then she had regular cycles. She had no symptoms of easy bruising, striae or proximal myopathy. Her medical and family history were unremarkable. She had history of thyroidectomy 25 years back, but was not on any thyroxine supplementation. On examination she was moderately built and nourished with Body Mass Index (BMI) of 28.2kg/m². She had hirsutism with features of virilization like male pattern hair loss, acne, atrophied breast and hypertrophied clitoris (Modified Ferriman Gallwey score was 12) (Fig 1A). She had a large Café-au-lait spot in the upper abdomen and an umbilical hernia which was reducible. There was no mass palpable in the abdomen. There were no features of hypercortisolism like moon facies, facial plethora, ecchymosis, purple striae or acanthosis nigricans. She had a neck scar which was healed with primary intention and had a grade 2 thyroid swelling, which was moved with deglutition and lower border was palpable. Hormonal evaluation was done (Table 1) which showed elevated testosterone and Dihydroepiandrosterone – sulphate (DHEA-S). Contrast Enhanced

Computerized Tomography of abdomen was done which showed heterogeneous enhancing mass right adrenal mass of size 45x35x32 mm with absolute washout of 58% and relative washout of 32% (Fig. 1B and 1C). Left adrenal gland was seen separate and normal. Lesion was in close contact with inferior vena cava (IVC) and has indiscernible fat planes with segment VI of liver. No obvious infiltration to liver or adjacent structures was visualized. Minimal ascites was also seen. Features were suggestive of adrenocortical carcinoma or lipid poor adenoma. As ascites and indiscernible fat planes with liver segment was demonstrated in CT abdomen, Positron Emission Tomography (PET) was done. It showed Fluoro Deoxy glucose (FDG) avid, heterogeneously enhancing well defined soft tissue density lesion in right adrenal of size 43x34x31 mm with SUV max 5.1. Hormonal evaluation was done for assessing functioning status of adrenal mass which was showed raised Free testosterone: 33.12pg/mL (Normal range :0.49-2.87 pg/mL) (Table 1). She underwent open right adrenalectomy. Intra operatively minimal free fluid was noticed in the abdominal cavity. Liver and other viscera were normal. Approximately 58x40x40 mm mass in the Right adrenal gland was seen and it was partly retro-caval with single right adrenal vein (Fig. 1D). She had smooth post operative recovery and her Serum cortisol was low (0.52 mcg/dl) on post operative day 1, for which hydrocortisone replacement was started. She was discharged on post operative day 5 and she was on follow up. Histopathology was consistent with adrenal adenoma- oncocytic variant (Fig. 2A,B,C and D) with absence of major and minor criteria. Her Lin- Weiss-Bisceglia score (Modified Weiss score) of 0. On follow up her hydrocortisone was tapered and at present she is off hydrocortisone and symptomatically better. Her post operative free testosterone came to normal.

3. DISCUSSION

Hirsutism is defined as an unwanted androgen-dependent hair growth in females [1]. Most of the causes of hirsutism are benign like

polycystic ovary syndrome, however it can rarely be life threatening [2]. Causes of hirsutism is summarized in Table 2. Excess androgen secretion in a pre- menopausal lady is mainly from ovary or adrenal gland. Testosterone is produced from ovary while DHEA-S is exclusively from adrenals. In post menopausal women adrenal gland becomes major source for androgens [3]. Oncocytomas are tumors composed of oncocytic cells; these cells are first

described by Hamprel in 1931 in thyroid gland as large eosinophilic, granular cells [4]. These granular cytoplasm is due to large number of mitochondria. The oncocytomas are commonly seen in kidney, salivary glands, thyroid and parathyroid glands. Adrenal oncocytomas are an extremely rare cause of adrenal tumors and the functional ones are only 30% as per the literature [5]. They usually present as benign, non functional adrenal incidentalomas.

Table 1. Lab reports

Hormone	Plasma Levels	Reference
FSH	5.01 mIU/ml	19.3-100.6mIU/mL
LH	5.01 mIU/ml	14.2-52.3mIU/L
S cortisol Random :	16.6mcg/dl	5-23mcg/dl
Cortisol (Over night Dexamethasone Suppression Test)	0.66 mcg/dl	Suppressed - Normal
Free testosterone	33.12pg/ml	0.49-2.87 pg/ml
Prolactin	4.7ng/ml	Up to 27 ng/ml
Plasma free metanephrine	8.41ng/L	7.9-88.7 ng/dl
Plasma free Nor-metanephrine	33.80ng/L	20.1-136.4 ng/dl
Thyroid stimulating Hormone (TSH)	0.316 mlu/L	0.5-4.5 mlu/L
T4	8.88mcg/dl	7.3-11.6 mcg/dl
17 OH Progesterone	2.11 ng/dl	<200 ng/dl

Table 2. Causes of hirsutism

Androgenic hirsutism	Non-androgenic hirsutism
Polycystic ovary syndrome (PCOS)	Idiopathic
Non-classical congenital adrenal hyperplasia	Drug-induced
Cushing syndrome	Testosterone, danazol, progestins, anabolic steroids, valproic acid, methyl dopa
Androgen-secreting ovarian tumor (eg, Sertoli-Leydig cell tumor)	
Androgen-secreting adrenal tumor (eg, adrenocortical carcinoma)	




Fig. 1. Clinical pictures of the patient (1 A) : Patient is having alopecia and virilization (1 B and C) : CT Abdomen showing right adrenal tumor (1 D):Specimen photograph of right adrenalectomy




Fig. 2. Microscopic features (2A): well circumscribed lesion showing capsule (2B): Nests of cells separated by thin vascular fibrous septae (2C and 2 D) : (High power)Cells have abundant granular eosinophilic cytoplasm and irregular vesicular nuclei with prominent nucleoli

Recent studies have shown that 65% of them were either malignant or had malignant potential [6]. These tumors are commonly seen in females and present with virilizing features and or hirsutism. In our case DHEA-S was elevated as in all other cases of adrenal oncocytoma reported in the literature. There are no specific signs in CT/MRI to reach the diagnosis. Mostly these may confuse with adreno cortical carcinoma (ACC) in non invasive investigations like CT/MRI. As we have a differential diagnosis of adreno cortical carcinoma, guided biopsy is contraindicated. FDG PET scan is used to differentiate benign and malignant lesions in adrenals, but only few reports are there in literature for adrenal oncocytomas and the findings are paradoxical [7]. The treatment for adrenal oncocytomas is surgical excision. As it has a malignant potential and a differential diagnosis of adreno cortical carcinoma, a complete resection without disrupting the capsule is mandatory. Recent studies have shown that laparoscopic surgeries are safe in large adrenal masses of more than 6 cm [8]. Even though the experience in laparoscopic approach could be taken for these type tumors, we opted for an open approach in view of ACC possibility and thereby getting a complete resection.

The final diagnosis of oncocytoma depended on the histopathological examination.

Microscopically, they are composed of large cell with eosinophilic and granular cytoplasm and central pyknotic nuclei. There are specific criteria for benign, borderline or malignant oncocytomas as described by Bisceglia et al in 2004 [9]. Prognosis of these tumors will depend up on the cited criteria for adreno cortical oncocytic neoplasm known as Lin-weiss-Bisceglia system. These system includes 3 major criteria, 4 minor criteria and definitional criteria. If any one major criteria is present, it is carcinoma. Presence of any four minor criteria indicative of uncertain malignant potential and if there is no major or minor criteria, it is indicative of benign tumor. Because of the rarity of these tumors, clear cut guidelines are not there for post operative treatment and follow up. As far as benign oncocytomas concerned prognosis is excellent. As for borderline oncocytomas only one publication showed a recurrence. For malignant oncocytomas survival is up to 58 months which is more than the survival of adreno cortical carcinoma [10].

4. CONCLUSION

Adrenal oncocytomas are rare tumors which are commonly present as adrenal incidentalomas. Only histopathology can diagnose oncocytomas and its clinical behaviour. Treatment of choice is excision as open or laparoscopic adrenalectomy.

As there are no clear cut guidelines for follow up, it needs further studies and high levels of evidence.

CONSENT

All authors declare that 'written informed consent was obtained from the patient for publication of this case report and accompanying images.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hunter MH, Carek PJ. Evaluation and treatment of women with hirsutism. American Family Physician. 2003 Jun 15;67(12):2565-72.
2. Martin KA, Anderson RR, Chang RJ, Ehrmann DA, Lobo RA, Murad MH, Pugeat MM, Rosenfield RL. Evaluation and treatment of hirsutism in premenopausal women: An Endocrine Society clinical practice guideline. The Journal of Clinical Endocrinology & Metabolism. 2018 Apr;103(4):1233-57.
3. Shifren JL, Davis SR. Androgens in postmenopausal women: A review. Menopause. 2017 Aug 1;24(8):970-9.
4. Chang A, HARAWAI S. Oncotype, oncocytosis, and oncocytic tumors. Pathology Annual. 1992; 27:263-304.
5. Geramizadeh B, Norouzzadeh B, Bolandparvaz S, Sefidbakht S. Functioning adrenocortical oncocytoma: A case report and review of literature. Indian Journal of Pathology & Microbiology. 2008 Apr 1;51(2):237-9.
6. Kanitra JJ, Hardaway JC, Soleimani T, Koehler TJ, McLeod MK, Kavuturu S. Adrenocortical oncocytic neoplasm: A systematic review. Surgery. 2018 Dec 1;164(6):1351-9.
7. Kim DJ, Chung JJ, Ryu YH, Hong SW, Yu JS, Kim JH. Adrenocortical oncocytoma displaying intense activity on 18F-FDG-PET: A case report and a literature review. Annals of Nuclear Medicine. 2008 Nov;22(9):821-4.
8. Machado NO, Al Qadhi H, Al Wahaibi K, Rizvi SG. Laparoscopic adrenalectomy for large adrenocortical carcinoma. JSLS: Journal of the Society of Laparoendoscopic Surgeons. 2015 Jul;19(3).
9. Bisceglia M, Ludovico O, Di Mattia A, Bendor D, Sandbank J, Pasquinelli G, Lau SK, Weiss LM. Adrenocortical oncocytic tumors: Report of 10 cases and review of the literature. International Journal of Surgical Pathology. 2004 Jul;12(3):231-43.
10. Wong DD, Spagnolo DV, Bisceglia M, Havlat M, McCallum D, Platten MA. Oncocytic adrenocortical neoplasms—A clinicopathologic study of 13 new cases emphasizing the importance of their recognition. Human Pathology. 2011 Apr 1;42(4):489-99.

© 2022 Puzhakkal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93339>