



Response of Cucumber (*Cucumis sativus* L.) Plants to Vermi Tea as Organic Fertilizer Supplementation

Gleza Mae Z. Dela Cerna^a and Leo C. Radores^{b*}

^a BS in Agriculture Malita, Davao Occidental, Philippines.

^b Faculty SPAMAST Malita, Davao Occidental, Philippines.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAAR/2022/v18i130209

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84241>

Original Research Article

Received 20 January 2022

Accepted 25 March 2022

Published 09 April 2022

ABSTRACT

Aims: To determine the performance of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer

Study design: The experimental design was a completely randomized block design.

Place and Duration of Study: Southern Philippines Agribusiness and Marine and Aquatic School of Technology (SPAMAST) Institute of Agricultural Technology and Entrepreneurial Studies (IATES) Buhangin, Malita, Davao Occidental from October 2019 to January 2020.

Methodology: Randomized Complete Block Design with five treatments and replicated three times as follows: Treatment 1 (Control), Treatment 2 (150 ml of vermin-tea/liter of water), Treatment 3 (200 ml of vermin-tea/liter of water), Treatment 4 (250 ml of vermin-tea/liter of water) and Treatment 5 (300 ml of vermin-tea/liter of water).

Results: Results showed that there was a significant effect of vermi tea as organic fertilizer application of cucumber (*Cucumis sativus* L.) in terms of weight of fruits harvested while there was no significant effect on the number of days to emergence, number of flowers per plant, length of fruits, the diameter of fruits, and number of fruits harvested was no significant effect.

Conclusion: Based on the results, the application of vermi tea as organic fertilizer gives the best result only on the weight of fruit harvested where vermi tea as a source of Nitrogen, Phosphorous, and potassium has helped to develop cucumber (*Cucumis sativus* L.) fruits.

Keywords: Performance; cucumber; vermin- tea; organic fertilizer.

1. INTRODUCTION

Cucumber (*Cucumis sativus* L.) was an economically important family of botanical Cucurbitaceae, commonly called cucurbits and gourds. Cucumber is a hot plant growing wine and annual crops in the season. The cucurbit order belongs botanically to the magnoliophyte group, the class Magnoliopsida. The edible and economically important part were immature fruit and therefore its potential yield would be affected by its flowering characteristics, particularly its femininity [1].

Nowadays, cultivators have developed vermicompost (VC) aqueous extracts which are commonly called 'vermicompost' there are which can improve crop tolerance to disease, and plant growth and are much more easily applied to soils and plants [2]. Vermicompost is much better than traditional thermophilic compost than microbial productivity and development [3]. while microbial organic matter can biochemically deteriorate, earthworms are significant derivatives to the process, reinforce the substratum and switch biological functions [4]. Throughout vermicomposting, nutrients are generated and converted into soluble and available crops [5].

This study was conducted to evaluate the impact of vermin-tea as organic fertilizer on the performance of cucumber (*Cucumis sativus* L.).

1.1 Statement of the Problem

This study aimed to determine the performance of cucumber (*Cucumis sativus* L.) as applied with vermin-tea as organic fertilizer. Specifically, it sought to answer the following questions.

1. Is there a significant difference in the number of days to germinate cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer?
2. Is there a significant difference in the number of flowers per plant of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer?
3. Is there a significant difference in the length of fruits of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer?
4. Is there a significant difference in the diameter of fruits of cucumber (*Cucumis*

sativus L.) as applied with vermi tea as organic fertilizer?

5. Is there a significant difference in the number of fruits harvested of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer?
6. Is there a significant difference in the weight of the fruit of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer?

1.2 Significance of the Study

This study imparts and as a supplement some information gathered by other researchers and references to those people who are interested in adopting new technology. It shared knowledge with the small- and large-scale farmers on the production management of cucumber (*Cucumis sativus* L.) production through low-cost fertilizer and so it produced an organic product that is healthier and has no harmful effect on the human body.

1.3 Scope and Limitation of the study

The study was conducted in the experimental area of SPAMAST, Buhangin, Malita, and Davao Occidental in October 2019.

This was limited only to the effect of different levels of vermin-tea as applied organic fertilizer on the yield performance of cucumber (*Cucumis sativus* L.) with the following parameters: number of days to germinate, number of flowers per plant, length of fruits, the diameter of fruits, number of fruits harvested, and weight of fruits harvested.

2. METHODOLOGY

The study was conducted at SPAMAST Buhangin, Malita, Davao Occidental from October 2019 to January 2020. Buhangin, Malita, Davao Occidental is bounded by the Municipality of Sta. Maria on the North and Barangay Tubalan on the South while on the East by Davao Gulf. It occupies an area of 8.0 hectares and comprises about 3.58% of the total land area of the municipality of Malita.

The soil type of barangay Buhangin is characterized by an alluvial sandy loam soil. However, a portion of this area is silt, clay loam that extends to the shoreline of Davao gulf. The

experimental area has an elevation of more or less 53 meters above sea level.

Based on the Agro-climatic Map developed by the Philippine Atmospheric Geophysical and Astronomical Service Administration (PAGASA) Davao Occidental has a mean average rainfall of 87.85 mm per month with a mean temperature of 34.3^oC. It is blessed with a favorable climate and its rainfall pattern generally conforms to Types IV which is characterized by more or less evenly distributed rainfall.

2.1 Experimental Design and Treatments

This study was laid out in Randomized Complete Block Design (RCBD) with five (5) treatments replicated three times. Each treatment was randomly selected using the draw-lot technique.

The treatments were as follows:

- T₁ – Control
- T₂ – 150 ml of vermi tea/liter of water
- T₃ – 200 ml of vermi tea/liter of water
- T₄ – 250 ml of vermi tea/liter of water
- T₅ – 300 ml of vermi tea/liter of water

2.1.1 Experimental field lay-out

This study has a total area of 120 square meters measuring 10 meters wide and 12 meters in length. It consisted of 15 plots measuring 2 meters x 3 meters with a corresponding distance of 0.5-meter alley.

2.1.2 Randomization

Draw-lot technique was employed in the assignment of treatments to their respected plots and blocks.

2.1.3 Procurement of materials

Cucumber seeds and other materials such as a seed tray were purchased in Digos City, particularly in agricultural supplies. Vermicast was purchased in Basiawan, Sta. Maria, Davao Occidental.

2.1.4 Preparation of vermi tea

Vermi- tea was prepared by placing two (2) kilograms of vermicast, two (2) kilograms of molasses in a bucket of five (5) gallons, dissolving two (2) liters of water, and positioning the aerator. Covered the bucket to keep it free of

insects and animals. Placed the brewing in a shaded region out of the immediate sunlight. Let it aerate, stirring occasionally, for 18 to 24 hours of fermentation.

2.2 Cultural Management and Practices

2.2.1 Land preparation

The area was slashed to remove unwanted plants and it was followed by plowing and harrowing twice to reduce the soil into a fine tilt. Thereafter, it was leveled and shallow furrows were made. Sticking was done after measuring the size of the plot and the planting distance. Digging was followed by 0.50 meters between furrows and 0.50 between hills.

2.2.2 Planting

The direct seeding method of planting was adapted with two (2) seeds per hill. The planting distance was 0.50 meters between furrows. Thinning out was done a week after the plants have germinated leaving only one plant per hill. Replanting was done after one week for missing hills.

2.2.3 Weeding and cultivation

Manual weeding was employed periodically to eliminate weeds that cause competition for nutrients. Weeding was employed late in the afternoon to prevent the plant stress. Shallow cultivation was done once a month to loosen up the soil and control the growth of weeds. This was done regularly to avoid competition and ensure the penetration and absorption of nutrients.

2.2.4 Fertilization

Vermi tea was sprayed three (3) days after planting and at seven (7) days intervals until the study was terminated. This was done early in the morning or before sunrise using a hand sprayer to ensure sufficient results.

2.2.5 Pest and diseases control

Cucumber plants are prone to the beetle, leaf folder, aphids, and powdery mildew that was attacked anytime the vegetable and reproductive stages of plants. To prevent the occurrence of these pests and diseases, insecticides and fungicides were sprayed regularly following the recommended dosage stipulated by the manufacturers. This was done using hand spray.

2.2.6 Trellis establishment

Cucumber was a vine crop belonging to the *Cucurbitaceae* family which needs a trellis or any materials that the plants could climb for their survival. A fence-type trellis was conducted made up of round numbers with a GI which was connected along the poles of every plant. This was done right after the seeds were germinated.

2.2.7 Watering

Cucumber was a characterized as high-water requirement crop. Heavy watering is desirable to ensure proper penetration to the root zones. Periodic watering was done especially during the drought period and early stages of plants. This was done using a water sprinkler with fine holes.

2.2.8 Harvesting

Harvesting was done when cucumber fruits reached their marketing or vegetable stage or when the pale green color changed to dark green. This was done by cutting the fruits one inch away from the stem.

2.3 Data Gathered

2.3.1 Number of days to emerge

This was determined by counting the average number of days of emergences per hill in every treatment per replication.

2.3.2 Number of flowers per plant

This was determined by counting the average number of flowers per plant per hill in every treatment per replication.

2.3.2 Length of fruits

This was determined by measuring the fruits per plant in every treatment per replication using a ruler.

2.3.3 Diameter of fruits

This was done by measuring the fruit diameter of cucumber from the base, mid, and tip of the ten sample plants using a tape measure and caliper.

2.3.4 Number of fruits harvested

This was determined by counting the fruits harvested.

3. RESULTS AND DISCUSSION

3.1 Number of Days to Emergence

Table 1 shows the yield performance of cucumber (*Cucumis sativus* L.) in terms of the number of days to emergence as affected by the application of vermi tea as organic fertilizer.

Treatment 3 got the lowest number of days to germinate cucumber (*Cucumis sativus* L.) with a mean value of 3.27 days followed by treatment 1 with a mean of 3.43 days, treatment 5 with a mean of 3.40 days, treatment 2 with a mean of 3.37 days while treatment 4 got the highest with a mean value of 3.60 days.

Analysis of Variance (ANOVA) exhibited that there was no significant difference among treatments on the growth and yield performance of cucumber (*Cucumis sativus* L.) in terms of the number of days to germinate as affected by the application of vermi -tea as organic fertilizer.

3.2 Number of Flowers per Plant

Table 2 shows the yield performance of cucumber (*Cucumis sativus* L.) in terms of the number of flowers per plant as affected by the application of vermi tea as organic fertilizer.

Treatment 3 got the highest number of flowers per plant of cucumber (*Cucumis sativus* L.) with a mean value of 38.40 flowers followed by treatment 4 with a mean of 38.07 flowers, treatment 1 with a mean of 37.87 flowers, treatment 2 with a mean of 37.73 flowers while treatment 5 got the lowest with a mean value of 34.20 flowers.

Analysis of Variance (ANOVA) exhibited that there was no significant difference among treatments on the growth and yield performance of cucumber (*Cucumis sativus* L.) in terms of the number of flowers per plant as affected by the application of vermi tea as organic fertilizer.

3.3 Length of Fruits

Table 3 shows the yield performance of cucumber (*Cucumis sativus* L.) in terms of the length of fruits as affected by the application of vermi tea as organic fertilizer.

Treatment 4 got the highest length of fruits of cucumber (*Cucumis sativus* L.) with a mean

value of 91.88 mm followed by treatment 3 with a mean of 81.45 mm, treatment 5 with a mean of 73.57 mm, treatment 2 with a mean of 62.60 mm while treatment 1 got the lowest with a mean value of 59.56 mm.

Analysis of Variance (ANOVA) exhibited that there was no significant difference among treatments on the growth and yield performance of cucumber (*Cucumis sativus* L.) in terms of length of fruits as affected by the application of vermi tea as organic fertilizer.

3.4 Diameter of Fruits

Table 4 shows the yield performance of cucumber (*Cucumis sativus* L.) in terms of the

diameter of fruits as affected by the application of vermi tea as organic fertilizer.

Treatment 3 got the highest diameter of fruits of cucumber (*Cucumis sativus* L.) with a mean value of 43.30 mm followed by Treatment 4 with a mean of 42.01 mm, treatment 1 with a mean of 39.53 mm, treatment 5 with a mean of 38.90 mm while treatment 2 got the lowest with a mean value of 38.23 mm.

Analysis of Variance (ANOVA) exhibited that there was no significant difference among treatments on the growth and yield performance of cucumber (*Cucumis sativus* L.) in terms of diameter of fruits as affected by the application of vermin- tea as organic fertilizer.

Table 1. Number of days to emergence of cucumber (*Cucumis sativus* L.) as applied with vermi -tea as organic fertilizer

Treatment	Mean (day)	CF	Tabular F	
			5%	1%
T ₁ – Control	3.43			
T ₂ – 150 ml Vermi Tea/liter of water	3.37	1.350 ^{ns}	3.84	7.01
T ₃ – 200 ml Vermi Tea/liter of water	3.27			
T ₄ – 250 ml Vermi Tea/liter of water	3.60			
T ₅ – 300 ml Vermi Tea/liter of water	3.40			

ns = not significant
CV = 5.31%

Table 2. Number of flowers per plant of cucumber (*Cucumis sativus* L.) as applied with vermi -tea as organic fertilizer

Treatment	Mean (flower)	CF	Tabular F	
			5%	1%
T ₁ – Control	37.87			
T ₂ – 150 ml Vermi Tea/liter of water	37.73	0.068 ^{ns}	.84	7.01
T ₃ – 200 ml Vermi Tea/liter of water	38.40			
T ₄ – 250 ml Vermi Tea/liter of water	38.07			
T ₅ – 300 ml Vermi Tea/liter of water	34.20			

ns = not significant
CV = 30.84%

Table 3. Length of fruits of cucumber (*Cucumis sativus* L.) as applied with vermi -tea as organic fertilizer

Treatment	Mean (mm)	CF	Tabular F	
			5%	1%
T ₁ – Control	59.56			
T ₂ – 150 ml Vermi Tea/liter of water	62.60	2.328 ^{ns}	.84	7.01
T ₃ – 200 ml Vermi Tea/liter of water	81.45			
T ₄ – 250 ml Vermi Tea/liter of water	91.88			
T ₅ – 300 ml Vermi Tea/liter of water	73.57			

ns = not significant
CV = 20.55%

3.5 Number of Fruits Harvested

Table 5 shows the yield performance of cucumber (*Cucumis sativus* L.) in terms of the number of fruits harvested as affected by the application of vermi tea as organic fertilizer.

Treatment 3 got the highest number of fruits harvested of cucumber (*Cucumis sativus* L.) with a mean value of 5.33 fruits followed by treatment 2 with a mean of 5.17 fruits, treatment 4 with a mean of 5.13 fruits, treatment 5 with a mean of 4.20 fruits while treatment 1 got the lowest with a mean value of 3.03 fruits.

Analysis of Variance (ANOVA) exhibited that there was no significant difference among treatments on the growth and yield performance of cucumber (*Cucumis sativus* L.) in terms of the number of fruits harvested as affected by the application of vermi tea as organic fertilizer.

3.6 Weight of Fruits Harvested

Table 6 shows the yield performance of cucumber (*Cucumis sativus* L.) in terms of the

weight of fruits harvested as affected by the application of vermi tea as organic fertilizer.

Treatment 4 got the highest weight of fruits harvested of cucumber (*Cucumis sativus* L.) with a mean value of 100.73 grams followed by treatment 3 with a mean of 100.45 grams, treatment 1 with a mean of 90.69 grams, treatment 2 with a mean of 80.33 grams while treatment 5 got the lowest with a mean value of 70.74 grams.

Analysis of Variance (ANOVA) exhibited that there was a significant difference among treatments on the growth and yield performance of cucumber (*Cucumis sativus* L.) in terms of the weight of fruits harvested as affected by the application of vermi tea as organic fertilizer.

The Least Significant Difference (LSD) showed that treatment 3 and treatment 4, were significantly different from treatment 1 (control). Whereas treatment 2 and treatment 5 were not significantly different from treatment 1 (control). It implies that there was a significant effect on the weight of fruits harvested of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer.

Table 4. Diameter of fruits of cucumber (*Cucumis sativus* L.) as applied with vermi -tea as organic fertilizer

Treatment	Mean (mm)	CF	Tabular F	
			5%	1%
T ₁ – Control	39.53			
T ₂ – 150 ml Vermi Tea/liter of water	38.23	0.779 ^{ns}	.84	7.01
T ₃ – 200 ml Vermi Tea/liter of water	43.30			
T ₄ – 250 ml Vermi Tea/liter of water	42.01			
T ₅ – 300 ml Vermi Tea/liter of water	38.90			

ns = not significant

CV = 10.51%

Table 5. Number of fruits harvested of cucumber (*Cucumis sativus* L.) as applied with vermi -tea as organic fertilizer

Treatment	Mean (fruit)	CF	Tabular F	
			5%	1%
T ₁ – Control	3.03			
T ₂ – 150 ml Vermi Tea/liter of water	5.17	1.753 ^{ns}	.84	7.01
T ₃ – 200 ml Vermi Tea/liter of water	5.33			
T ₄ – 250 ml Vermi Tea/liter of water	5.13			
T ₅ – 300 ml Vermi Tea/liter of water	4.20			

ns = not significant

CV = 27.76%

Table 6. Weight of fruits harvested of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer

Treatment	Mean (gram)	CF	Tabular F	
			5%	1%
T ₁ – Control	90.69			
T ₂ – 150 ml Vermi Tea/liter of water	80.33 ^{ns}	4.321*	.84	7.01
T ₃ – 200 ml Vermi Tea/liter of water	100.45*			
T ₄ – 250 ml Vermi Tea/liter of water	100.73*			
T ₅ – 300 ml Vermi Tea/liter of water	77.74 ^{ns}			

* = significant
CV = 11.62%

According to Sinha *et al.* [6], vermicast has millions of valuable soil microbes including nitrogen, potassium, and phosphorus (NPK) which significantly the growth of plants. It implies that potassium enhances dry biomasses of the plant through enhancing photosynthetic pigments content and cell metabolic processes including; cell growth and elongation [7]. Numerous studies indicate application of potassium, increases the plant dry weight of plants [8]. As regards the physical fruit traits, many efforts illustrated that potassium improved fruit weight and dry matter contents [9] as well as increased cucumber fruit biomasses [10].

4. SUMMARY, CONCLUSION, AND RECOMMENDATION

This study was conducted to evaluate the performance of cucumber (*Cucumis sativus* L.) as applied with vermi tea as organic fertilizer at the Southern Philippines Agribusiness and Marine and Aquatic School of Technology (SPAMAST) Institute of Agricultural Technology and Entrepreneurial Studies (IATES) Buhangin Malita Davao Occidental from September 2019 to January 2020.

Complete Randomized Design was laid out with five treatments and replicated three times as follows: Treatment 1 (Control), Treatment 2 (150 ml of vermi tea/liter of water), Treatment 3 (200 ml of vermi tea/liter of water), Treatment 4 (250 ml of vermi tea/liter of water) and Treatment 5 (300 ml of vermi tea/liter of water).

The following parameters were taken: number of days to germinate, number of flowers per plant, length of fruits, the diameter of fruits, number of fruits harvested, and weight of fruits harvested. The result showed that there was a significant effect of vermi tea as an organic fertilizer application on the yield of cucumber (*Cucumis sativus* L.) in terms of weight of fruits harvested

while the number of days to germinate, the number of flowers per plant, length of fruits, the diameter of fruits, and several fruits harvested was insignificant,

4.1 Conclusion

Based on the study results, the application of vermi tea as organic fertilizer gives the best result only on the weight of fruit harvested cucumber (*Cucumis sativus* L.) where vermi tea is a source of Nitrogen, Phosphorous and potassium has helped to develop cucumber (*Cucumis sativus* L.) fruits.

4.2 Recommendation

Based on the result the researcher hereby recommends using Treatment 4 (250 ml of vermi tea/liter of water) as it gives a significant result on the weight of fruits harvested. It also recommends conducting the further study with additional organic fertilizer to supplement the yield of cucumber (*Cucumis sativus* L.).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tanurdzic M, Banks JA. Sex-determining mechanisms in land plants. *The Plant Cell*. 2004;16: S61–S71.
2. Ingham ER. *The Compost Tea Brewing Manual* 4th ed., Corvallis: Soil Foo web Inc; 2003.
3. Edward T. *Suppressing Plant Parasitic Nematodes and Arthropod Pest with Vermicompost Teas*; 2004.
4. Aira M, Monroy F, Dominguez J. Eisenia fetida (*Oligochaeta: Lumbricidae*) Modifies the Structure and Physiological

- Capabilities of Microbial Communities Improving Carbon Mineralization during Vermicomposting of Pig Manure. *Microbe Ecol.* 2002;662-671.
5. Ndegwa PM, Thompson SA. Effects of C-to-N ratio on vermicomposting of biosolids. *Bioresource technology.* 2001; 75:7–12.
 6. Sinha RK, Bharambe G, Chowdhary U. Sewage Treatment by Vermi-filtration With Synchronous Treatment of Sludge by Earthworms: A Low-Cost Sustainable Technology Over Conventional Systems with Potential for Decentralization; Springer, USA. *The Environmentalist*; UK. 2008; 28:409– 420.
 7. Farago ME. Plants and the chemical elements: biochemistry, uptake, tolerance, and toxicity. Wiley-Blackwell, New York, USA; 2008.
 8. Botella MA, Arévalo L, Mestre TC, Rubio F, García-Sánchez F, Rivero RM, Martínez V. Potassium fertilization enhances pepper fruit quality. *Journal of Plant Nutrition.* 2017; 40:145-155.
 9. Gupta CR, Sengar SS. Response of tomato (*Lycopersicon esculentum mill.*) to nitrogen and potassium fertilization in acidic soil of Bastar. *Journal of Vegetation Science.* 2000; 27:94-95.
 10. Akram MS, Ashraf M. Exogenous application of potassium dihydrogen phosphate can alleviate the adverse effects of salt stress on sunflower (*Helianthus annuus L.*). *Journal of Plant Nutrition.* 2011; 34:1041-1057.

© 2022 Cerna and Radores; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84241>