



## Outcome of Unilateral Sinonasal Masses in University of Port Harcourt Teaching Hospital

A. C. Nwogbo <sup>a\*</sup> and J. Kpobene <sup>a</sup>

<sup>a</sup> Department of Otolaryngology, University of Port Harcourt Teaching Hospital, Nigeria.

### Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

### Article Information

#### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94571>

Original Research Article

Received 19 October 2022  
Accepted 17 December 2022  
Published 24 December 2022

### ABSTRACT

**Background:** Unrelenting nasal obstruction sometimes associated with rhinorrhoea may suggest the presence of sinonasal growth. Two major groups commonly exist, neoplastic or inflammatory.

**Aim:** The aim of this study is to present the clinical pattern of unilateral nasal masses in University of Port Harcourt Teaching Hospital.

**Materials and Methods:** Patients managed for unilateral nasal masses were considered, between January 2017 – January 2021. The following demographic data were recorded; age, gender, occupation, presenting symptoms, duration of symptoms. Histological type and treatment given were also analysed.

**Results:** Forty three (43) cases were reviewed in this study. There were (28) males and females (15) with gender ratio of 2:1. Most common presentation were rhinorrhoea, nasal obstruction and presence of nasal growth. Common histological type was inflammatory polyp.

**Conclusion:** Most common unilateral sinonasal mass was inflammatory polyps. This was followed by inverted papilloma.

*Keywords:* Inflammatory; polyp; nasal obstruction; sinonasal mass.

### 1. INTRODUCTION

Sinonasal mass may present with nasal obstruction and rhinorrhoea. Persistent unilateral

obstruction may suggest the presence of sinonasal lesion, which may be inflammatory or neoplastic [1].

\*Corresponding author: E-mail: [nwogbo\\_augustine@yahoo.com](mailto:nwogbo_augustine@yahoo.com);

Most times inflammatory polyps present commonly than neoplastic lesions in unilateral nasal lesions. It is usually common to conclude that unilateral sinonasal lesion in adult is either neoplastic or inverted papilloma [2].

However, some workers have reported nasal polyp and squamous cell carcinoma as common sinonasal lesion.

Common clinical conditions associated with inflammatory polyp include asthma, allergy, infection, and cystic fibroma.

Patients who have unilateral nasal masses may present with nasal obstruction, rhinorrhoea, hyposmia, facial pain. Sometimes it may be difficult to differentiate with common cold, especially at early stage. Hence full evaluation of patient is mandatory [3,4].

The aim of this study is to present the clinical pattern and outcome of unilateral nasal masses in University of Port Harcourt Teaching Hospital.

## 2. MATERIALS AND METHODS

This was a retrospective study of patients who were managed for unilateral nasal masses at the Department of Otolaryngology of the University of Port Harcourt Teaching Hospital, which is a tertiary health institution in the South-South Nigeria.

The records of patients managed for unilateral nasal masses were retrieved and the following information were recorded; age, gender, occupation, presenting symptoms and duration of symptoms. Other recorded data were plain radiological findings and CT Scan findings. Histological results, type of treatment given and clinical status of the patient were also recorded.

Each patients had full history, clinical examination, appropriate imaging and examination under anaesthesia with biopsy. All information were recorded on tables.

## 3. RESULTS

This is a five (5) year retrospective study of forty three (43) cases that qualified as unilateral sinonasal masses. Twenty eight (28) male and fifteen (15) female with m:f ratio of 2:1. Most common age group affected was 56-65 years whereas the least affected was 16-25.

Most common symptoms were rhinorrhoea and nasal obstruction (Table 2), seen in almost all benign sinonasal masses.

Table 3 showed the common histological type of sinonasal masses (inflammatory polyp). This is followed by inverted papilloma. Squamous cell carcinoma presented as most common malignant unilateral sinonasal masses.

**Table 1. Age/gender distribution**

Age Group (years)	Male	Female	Total (%)
16-25	1	1	4.5
26-35	4	2	13.9
36-45	5	4	20.9
46-55	7	2	20.9
56-65	10	5	34.8
66-75	1	1	4.5
<b>Total</b>	<b>28</b>	<b>15</b>	<b>43</b>

**Table 2. Symptoms at presentation**

Symptoms	Non-neoplastic	Benign	Malignant	Total (%)
Rhinorrhoea	19	17	7	43
Nasal obstruction	19	16	8	43
Mass	19	14	8	41
Epistaxis	0	9	10	19
Diplopia	0	1	3	4
Proptosis	0	0	1	1
Cheek swelling	0	1	1	2

**Table 3. Histological types of unilateral nasal masses**

<b>Tumour Type</b>	<b>Number of Patients</b>
Non neoplastic (inflammatory polyp)	18 (42%)
<b>Benign</b>	
Inverted Papilloma	11 (26%)
Pleomorphic Adenoma	3 (7%)
Angiofibroma	1 (2.3%)
Cavernous Haemangioma	1 (2.3%)
<b>Malignant</b>	
Squamous cell carcinoma	6 (14%)
Rhabdomyosarcoma	1 (2.3%)
Lymphoma	1 (2.3%)
Adenocarcinoma	1 (2.3%)

**Table 4. Duration of Nasal mass**

<b>Duration(yrs)</b>	<b>Non-neoplastic</b>	<b>Benign</b>	<b>Malignant</b>	<b>Total</b>
1-3	15	8	6	29
4-6	3	2	2	7
7-9	0	2	0	2
10-12	0	5	0	5
<b>Total (100%)</b>	<b>18(41.8%)</b>	<b>17 (39.5%)</b>	<b>8(18.6%)</b>	<b>43</b>

**4. DISCUSSION**

Neoplastic and non-neoplastic lesions are common causes of unilateral sinonasal masses. Majority of patient in this study presented in the fifth decade (about 34.8%). This has been identified in various other studies [4,5].

The male/female ratio in our study was 2:1. In a study of Shuaibu et al, they had male/female ratio of 1:9:1 [6]. Despite numerous presentation like nasal obstruction, epistaxis, hyposmia, rhinorrhoea, facial pain, anosmia, however nasal blockage and rhinorrhoea and nasal mass remain the most common presentations observed in our study [6,7]. These were observed in both neoplastic and non-neoplastic cases. Epistaxis presentations dominated in our study for neoplastic unilateral sinonasal masses [8]. Nair et al reported nasal obstruction as most common presentation in non-neoplastic cases [8,9].

In our study, epistaxis was noted to be common presentation in neoplastic unilateral masses. Pazsilva et al in their study, demonstrated that epistaxis was commonly associated with neoplastic unilateral sinonasal masses [9,10].

In our study, non-neoplastic cases constituted about 42%. Similar findings were also demonstrated by Kucur et al in their study [11]. While our study showed inflammatory polyp as most common non-neoplastic lesion, Kucur et al reported inflammatory polyp, chronic sinusitis and antrochoanal polyp as most common [11,12].

Inverted papilloma was the most common benign tumour in our study, though benign, but usually aggressive in its presentation. This tumour is notorious for recurrence after excision [13]. This was followed by Pleomorphic Adenoma. Others were Angiofibroma and Cavernous Haemangioma (2.3%).

In another study by Humayun et al. [14], they reported inverted papilloma, meningioma followed by haemangioma as most common benign unilateral nasal masses.

In this study nine (9) patients had sinonasal malignant condition. Squamous Carcinoma (14%), Rhabdomyosarcorma (2.3%) lymphoma (2.3%), followed by Adenocarcinoma (2.3%).

However, in another study Belli et al, squamous carcinoma was reported as commonest, followed

by adenocystic carcinoma was reported and B-cell non-Hodgkin lymphoma [15].

We observed that majority of our patients presented within 1-3 years of the onset of symptoms. Those with malignancy presented earlier [16]. It could be as a result of symptoms like epistaxis, facial pain which are more alarming to the patients [17]. The benign unilateral lesion tend to present later, probably because grave symptoms are not experienced by the patient, like epistaxis [18].

## 5. CONCLUSION

Most common amongst benign unilateral sinonasal mass is inflammatory polyp, followed by inverted papilloma. Rhinorrhoea and nasal obstruction presented commonly in all patients. Patient with neo-plastic masses commonly presented with epistaxis. The outcome of benign lesion was better than in malignant cases. Any sinonasal mass associated with epistaxis should be suspicious of malignant lesion.

## CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Bachert C, Hormann ., Mosges R. An update on the diagnosis and treatment of sinusitis and nasal polyposis. *Allergy* 2003;58;176-91.
2. Badia L, Lund V. Topical corticosteroids in nasal polyposis. *Drugs*. 2001;61:573-8.
3. Karthikeya P, Mahima VG, Bhavna G. Sinonasal verrucous carcinoma with oral invasion. *Indian J Dent Res*. 2006;1:82-6.
4. Kristensen S, Vorre P, Elbrond O, Sogaard H. Nasal schneiderian papillomas. *Clin Otolaryngol*. 1985;10: 12534.
5. Uysal İÖ, Misir M, Polar K, Altuntas EE, Atalar MH, Tuncer E, et al. Primary malignant melanoma of the nasal cavity. *J Craniofac Surg*. 2012;23:e2-5.
6. Shuaibu I Y, Usman M A, Ayiye A. Unilateral sinonasal masses: Review of clinical presentation and outcome Niger Med J. 2020;61:16-21.
7. Hulse KE, Stevens WW, Tan BK, Schleimer RP. Pathogenesis of nasal polyposis. *Clin Exp Allergy*. 2015;45:328-46.
8. Kim WA, Tumours of the nose and sinuses. In: Hussain SM, editor. *Logan turner's Diseases of the Nose, Throat and Ear Head and Neck Surgery*. 11<sup>th</sup> ed. New York: CRC Press. 2016;119-29.
9. Nair S, James E, Awasthi S, Nambiar S, Goyal S. A review of the clinicopathological and radiological features of unilateral nasal mass. *Indian J Otolaryngol Head Neck Surg*. 2013;65: 199-204.
10. Paz Silva M, Pinto JM, Covey JP, Mhoon EE, Barody FM, Naclerio RM. Diagnostic algorithm for unilateral sinus disease: A 15-year retrospective review. *Int Forum Allergy Rhinol*. 2015;5:590-6.
11. Kucur C, Oghan F, Özbay I, Erdogan O, Tok S, Sanal B, et al. Unilateral nasal pathologies: Clinical presentation and management. *ENT Updates*. 2015;5:23-9.
12. Kahveci OK, Duran A, Miman MC. Our histopathological result for intranasal masses: Retrospective study of 6 years. *J Clin Anal Med*. 2012;3:289-92.
13. Gomes P, Gomes A, Salvador P, Lombo C, Caselhos S, Fonseca R. Clinical assessment, diagnosis and management of patients with unilateral sinonasal disease. *Acta Otolaryngol Esp*. 2020;71:16-25.
14. Humayun AH, Huq AH, Ahmed SM, Kamal MD, Kyaw KU, Nilakanta B. Clinicopathological study of sinonasal masses. *Banladesh J Otorhinolaryngol*. 2010;16:15-22.
15. Belli S, Yildirim M, Eroglu S, Emre FK. Single-sided sinonasal mass: A retrospective study. *North Clin Istanbul*. 2018;5:139-43.
16. Bohman A, Oscarsson M, Holmberg K, Johansson L, Millqvist E, Nasic S, et al. Heredity of nasal polyps. *Rhinology*. 2015;53:25-8.
17. Kerewala C, Clarke P, Newbold K. Nasal cavity and paranasal sinus malignancy. In: Watkinson JC, Clarke RW, editors. *Scott-*

- Brown's Otorhinolaryngology Head and Neck Surgery. 8<sup>th</sup> ed. New York: CRC Press. 2009;73-91.
18. Leoncini G, Zanetti L. The papillomas of the sinonasal tract. A comprehensive review. *Pathologica*. 2017;109:31-4.

---

© 2022 Nwogbo and Kpobene; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/94571>*