



Fixed Point Theorems Using Soft Multiplicative Generalized Weak Contractive Mappings

Shalini Nagpal ^{a,b}, Sushma Devi ^c and Manoj Kumar ^{a*}

^a Department of Mathematics, Baba Mastnath University, Asthal Bohar, Rohtak, India.

^b Govt. P.G. College for Women, Rohtak, India.

^c Department of Mathematics, Kanya Mahavidyalya, Kharkhoda, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMCS/2023/v38i11739

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95326>

Original Research Article

Received: 20/10/2022

Accepted: 28/12/2022

Published: 19/01/2023

Abstract

In the present paper, we establish fixed point theorems for mappings satisfying generalized weak contractive conditions in soft multiplicative metric space.

Keywords: Soft metric space; soft multiplicative metric space; soft multiplicative weak contractive mapping; fixed point.

MSC: 47H10, 54H25

*Corresponding author: Email: manojantil18@gmail.com;

1 Introduction

In 1874, Cantor defined set theory as a branch of mathematics. This theory deals with the problems that contain certain results. But in real life situations, there are various uncertain problems which have imprecise results. To deal with these problems, Molodtsov [1], in 1999, introduced soft set theory and applied this theory in various fields like game theory, operations research etc. In 2002, Maji et al. [2,3] “worked on soft set theory and its applications in decision making problems”. In 2011, Ali et al. [4] “defined various operations in this theory”. In 2013, Wardowski [5] worked on soft mappings with the fixed point theorems.

As metric space is one of the prominent branches of mathematics, thus to explore this idea using soft sets, Das and Samanta [6,7] investigated the properties of soft real numbers in 2012 and in 2013, they introduced the concept of soft metric space. After this, several researchers worked on soft metric spaces and their properties. In 2008, Bashirov et al. [8] defined multiplicative metric space. Then various authors worked on this space [9-14]. Rathee et al. [15] combined “soft metric space and multiplicative metric space and generated a new space called soft multiplicative metric space”.

Fixed point theory plays a vital role in various fields of mathematics. In 2016, Wadkar et al. [16] proved fixed point results related to soft sets and in the same year, Yazar et al. [17] proved some fixed point theorems of soft contractive mappings. In 2017, Hosseinzadeh [18] proved fixed point theorems in soft metric space. Then, Abbas et al. [19] introduced various results on fixed point theorems in soft metric spaces. After this, in 2021, Bhardwaj et al. [20] investigated some new fixed point results in soft metric space. Rathee et al. [15] derived some fixed point theorems in soft multiplicative metric space. In 2017, Solanki et al. [21] “generalized the concept of soft weak contractive mapping and proved various fixed point theorems in soft metric space”. Extending the work of Solanki et al. [21] and Rathee et al. [15], we generate some new fixed point theorems using generalized multiplicative weak contraction mapping in soft multiplicative metric space.

2 Preliminaries

This section contains some basic definitions and results which are useful for our research work.

Definition 2.1[6]. “Let I be an initial universal set and Ω be the non-empty parameter set. Then, a pair (T, Ω) is called a soft set over I if T is a set valued mapping on Ω taking values in 2^I i.e., $T: \Omega \rightarrow 2^I$.”

Definition 2.2[6]. “A soft set (T, Ω) over I is said to be an absolute soft set if $T(\alpha) = I \quad \forall \alpha \in \Omega$. It is denoted by \tilde{I} .”

Definition 2.3[6]. “A soft set (T, Ω) over I is said to be a soft point if there is exactly one $\alpha \in \Omega$ such that $T(\alpha) = \{i\}$ for some $i \in I$ and $T(\beta) = \emptyset \quad \forall \beta \in \Omega \setminus \{\alpha\}$. Such a soft point is denoted by \tilde{T}_α^i .”

NOTE. The collection of all soft points of a soft set (T, Ω) is denoted by $SP(T, \Omega)$.

Definition 2.4[6]. “Let \mathbb{R} be the set of real numbers and $B(\mathbb{R})$ be the collection of all non-empty bounded subsets of \mathbb{R} . Then, the function by $T: \Omega \rightarrow B(\mathbb{R})$ is called a soft real set and is denoted by (T, Ω) . If T is a single valued function on Ω taking values in \mathbb{R} , then the pair (T, Ω) or simply T is called a soft real number. We denote soft real number and soft constant real number by $\tilde{r}, \tilde{s}, \tilde{t}$ and $\bar{r}, \bar{s}, \bar{t}$ respectively where \bar{r} will denote a particular type of soft real number such that $\bar{r}(\alpha) = r$ for all $\alpha \in \Omega$.”

Definition 2.5[6]. “For two soft real numbers \tilde{p} and \tilde{q} , the following conditions hold for all $\alpha \in \Omega$:

- (a) $\tilde{p} \lesssim \tilde{q}$ if $\tilde{p}(\alpha) \lesssim \tilde{q}(\alpha)$;

- (b) $\tilde{p} \succeq \tilde{q}$ if $\tilde{p}(\alpha) \succeq \tilde{q}(\alpha)$;
- (c) $\tilde{p} < \tilde{q}$ if $\tilde{p}(\alpha) \prec \tilde{q}(\alpha)$;
- (d) $\tilde{p} > \tilde{q}$ if $\tilde{p}(\alpha) \succ \tilde{q}(\alpha)$.”

Definition 2.6[6]. “A mapping $\tilde{\rho}: SP(\tilde{I}) \times SP(\tilde{I}) \rightarrow R(\Omega)^*$ is a soft metric on the absolute soft set \tilde{I} if $\tilde{\rho}$ satisfies the following conditions:

1. $\tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) \succeq \bar{0}$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$;
2. $\tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) = \bar{0}$ if and only if $\alpha = \beta$ and $i = j$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$;
3. $\tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) = \tilde{\rho}(\tilde{T}_\beta^j, \tilde{T}_\alpha^i)$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$;
4. $\tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\gamma^k) \preceq \tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) + \tilde{\rho}(\tilde{T}_\beta^j, \tilde{T}_\gamma^k)$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k \in SP(\tilde{I})$.

The soft set \tilde{I} together with soft metric $\tilde{\rho}$ is called a soft metric space and is denoted by $(\tilde{I}, \tilde{\rho}, \Omega)$ or simply by $(\tilde{I}, \tilde{\rho})$.”

Definition 2.7[8]. “A mapping $\rho^*: I \times I \rightarrow \mathbb{R}^*$ is multiplicative metric if ρ^* satisfies the following conditions:

1. $\rho^*(u, v) \geq 1$ for all $u, v \in I$;
2. $\rho^*(u, v) = 1$ if and only if $u = v$ for all $u, v \in I$;
3. $\rho^*(u, v) = \rho^*(v, u)$ for all $u, v \in I$;
4. $\rho^*(u, w) \leq \rho^*(u, v) \cdot \rho^*(v, w)$ for all $u, v, w \in I$.

The pair (I, ρ^*) is called a multiplicative metric space.”

Definition 2.8[15]. “A function $\tilde{\rho}^*: SP(\tilde{I}) \times SP(\tilde{I}) \rightarrow R(\Omega)^*$ is soft multiplicative metric on the absolute soft set \tilde{I} if $\tilde{\rho}^*$ meets the following properties:

1. $\tilde{\rho}^*(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) \succeq \bar{1}$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$;
2. $\tilde{\rho}^*(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) = \bar{1}$ if and only if $\alpha = \beta$ and $i = j$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$;
3. $\tilde{\rho}^*(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) = \tilde{\rho}^*(\tilde{T}_\beta^j, \tilde{T}_\alpha^i)$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$;
4. $\tilde{\rho}^*(\tilde{T}_\alpha^i, \tilde{T}_\gamma^k) \preceq \tilde{\rho}^*(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) \tilde{\rho}^*(\tilde{T}_\beta^j, \tilde{T}_\gamma^k)$ for all $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k \in SP(\tilde{I})$.

The soft set \tilde{I} together with soft multiplicative metric $\tilde{\rho}^*$ is called a soft multiplicative metric space and is denoted by $(\tilde{I}, \tilde{\rho}^*, \Omega)$.”

Definition 2.9[15]. “Suppose $(\tilde{I}, \tilde{\rho}^*)$ is a soft multiplicative metric space. Then, a sequence $\{\tilde{T}_{\alpha_n}^{i_n}\}$ in $(\tilde{I}, \tilde{\rho}^*)$ is soft multiplicative convergent to a soft point $\tilde{T}_\beta^j \in \tilde{I}$ if for given $\tilde{\varepsilon} \succeq \bar{1}$, we have a unique positive integer n_0 such that $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_\beta^j) \prec \tilde{\varepsilon}$ for all $n \geq n_0$ i.e., $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_\beta^j) \rightarrow \bar{1}$ as $n \rightarrow \infty$.”

Definition 2.10[15]. “Suppose $(\tilde{I}, \tilde{\rho}^*)$ is a soft multiplicative metric space. Then, a sequence $\{\tilde{T}_{\alpha_n}^{i_n}\}$ in $(\tilde{I}, \tilde{\rho}^*)$ is soft multiplicative Cauchy sequence if for given $\tilde{\varepsilon} \gtrsim \bar{1}$, we have a unique positive integer n_0 such that $\tilde{\rho}^*(\tilde{T}_{\alpha_m}^{i_m}, \tilde{T}_{\alpha_n}^{i_n}) \lesssim \tilde{\varepsilon}$ for all $m, n \geq n_0$ i.e., $\tilde{\rho}^*(\tilde{T}_{\alpha_m}^{i_m}, \tilde{T}_{\alpha_n}^{i_n}) \rightarrow \bar{1}$ as $m, n \rightarrow \infty$.”

Definition 2.11[15]. “A soft multiplicative metric space $(\tilde{I}, \tilde{\rho}^*)$ is complete, if every soft multiplicative Cauchy sequence in \tilde{I} converges to some soft point in \tilde{I} .”

Definition 2.12[17]. “Let $(\tilde{I}, \tilde{\rho}, \Omega)$ and $(\tilde{I}', \tilde{\rho}', \Omega')$ be two soft metric spaces. Then, $(h, \psi): (\tilde{I}, \tilde{\rho}, \Omega) \rightarrow (\tilde{I}', \tilde{\rho}', \Omega')$ is a soft mapping where $h: I \rightarrow I'$ and $\psi: \Omega \rightarrow \Omega'$ are two mappings.”

Definition 2.13[15]. “Consider a soft multiplicative metric space $(\tilde{I}, \tilde{\rho}^*, \Omega)$. A function $(h, \psi): (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$ is said to be soft multiplicative contraction mapping if for every soft point $\tilde{T}_\alpha^i, \tilde{T}_\beta^j \in \tilde{I}$, there exists a soft real number $\bar{\lambda}$, $\bar{0} \lesssim \bar{\lambda} \lesssim \bar{1}$ such that $\tilde{\rho}^*\{(h, \psi)(\tilde{T}_\alpha^i), (h, \psi)(\tilde{T}_\beta^j)\} \lesssim \{\tilde{\rho}^*(\tilde{T}_\alpha^i, \tilde{T}_\beta^j)\}^{\bar{\lambda}}$.”

Definition 2.14[21]. “A mapping $(h, \psi): (\tilde{I}, \tilde{\rho}, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}, \Omega)$, where $(\tilde{I}, \tilde{\rho}, \Omega)$ is a soft metric space, is said to be soft weakly C-contractive or a soft weak contraction if $\forall \tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$,

$$\tilde{\rho}\{(h, \psi)(\tilde{T}_\alpha^i), (h, \psi)(\tilde{T}_\beta^j)\} \lesssim \frac{\bar{1}}{2} \left[\tilde{\rho}\{\tilde{T}_\alpha^i, (h, \psi)(\tilde{T}_\beta^j)\} + \tilde{\rho}\{\tilde{T}_\beta^j, (h, \psi)(\tilde{T}_\alpha^i)\} \right] - \xi \left[\tilde{\rho}\{\tilde{T}_\alpha^i, (h, \psi)(\tilde{T}_\beta^j)\}, \tilde{\rho}\{\tilde{T}_\beta^j, (h, \psi)(\tilde{T}_\alpha^i)\} \right],$$

where $\xi: [\bar{0}, \infty)^2 \rightarrow [\bar{0}, \infty)$ is a continuous mapping such that $\xi(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) = \bar{0}$ if and only if one of $\tilde{T}_\alpha^i, \tilde{T}_\beta^j = \bar{0}$.”

Definition 2.15[21]. “A mapping $(h, \psi): (\tilde{I}, \tilde{\rho}, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}, \Omega)$, where $(\tilde{I}, \tilde{\rho}, \Omega)$ is a soft metric space, is said to be soft generalized weakly contractive or a soft generalized weak contraction if $\forall \tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$,

$$\tilde{\rho}\{(h, \psi)(\tilde{T}_\alpha^i), (h, \psi)(\tilde{T}_\beta^j)\} \lesssim \bar{\eta} \left[\max \left\{ \tilde{\rho}\{\tilde{T}_\alpha^i, (h, \psi)(\tilde{T}_\alpha^i)\}, \tilde{\rho}\{\tilde{T}_\beta^j, (h, \psi)(\tilde{T}_\beta^j)\}, \right. \right. \\ \left. \left. \tilde{\rho}\{\tilde{T}_\alpha^i, (h, \psi)(\tilde{T}_\beta^j)\}, \tilde{\rho}\{\tilde{T}_\beta^j, (h, \psi)(\tilde{T}_\alpha^i)\}, \tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) \right\} \right] - \xi \left[\left\{ \tilde{\rho}\{\tilde{T}_\alpha^i, (h, \psi)(\tilde{T}_\alpha^i)\}, \tilde{\rho}\{\tilde{T}_\beta^j, (h, \psi)(\tilde{T}_\beta^j)\}, \right. \right. \\ \left. \left. \tilde{\rho}\{\tilde{T}_\alpha^i, (h, \psi)(\tilde{T}_\beta^j)\}, \tilde{\rho}\{\tilde{T}_\beta^j, (h, \psi)(\tilde{T}_\alpha^i)\}, \tilde{\rho}(\tilde{T}_\alpha^i, \tilde{T}_\beta^j) \right\} \right],$$

where $\bar{\eta} \in \left[\bar{0}, \frac{\bar{1}}{2} \right)$, $\xi: [\bar{0}, \infty)^5 \rightarrow [\bar{0}, \infty)$ is a continuous mapping such that $\xi(\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\varepsilon^m) = \bar{0}$ if and only if one of $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\varepsilon^m = \bar{0}$.”

3 Main Results

In this section, we define soft multiplicative generalized weakly contractive mappings and prove fixed point results using these mappings.

Theorem 3.1. Let $(\tilde{I}, \tilde{\rho}^*, \Omega)$ be a complete soft multiplicative metric space and $(h, \psi): (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$ be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping:

$$\tilde{\rho}^*\{(h, \psi)(\tilde{T}_\alpha^i), (h, \psi)(\tilde{T}_\beta^j)\} \lesssim \frac{\left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j\} \right]^{\bar{p}} \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\beta^j\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\alpha^i\} \right]^{\bar{q}}}{\xi \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\beta^j\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\alpha^i\} \right]} \forall \tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I}),$$

where \bar{p}, \bar{q} are non-negative soft real numbers such that $\bar{p} + \bar{q} < \frac{1}{2}$ and $\xi: [\bar{1}, \infty)^2 \rightarrow [\bar{1}, \infty)$ is a continuous function such that $\xi(\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l) = \bar{1}$ iff one of $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l = \bar{1}$. Then, there exists a unique fixed point of (h, ψ) .

Proof. Let $\tilde{T}_{\alpha_0}^{i_0}$ be any soft point in $SP(\tilde{I})$. Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi)\tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi)\tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi)\tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) &= \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}})\} \\ &\quad \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n}\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\} \right]^{\bar{p}} \\ &\quad \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_n}^{i_n}\} \right]^{\bar{q}} \\ &\stackrel{\lesssim}{\leq} \xi \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n}\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\}, \right. \\ &\quad \left. \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_n}^{i_n}\} \right] \\ &\quad \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{p}} \\ &\quad \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{q}} \\ &\stackrel{\lesssim}{\leq} \xi \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \right. \\ &\quad \left. \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}) \right\} \\ &\quad \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{p}} \\ &= \xi \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{q}} \\ &\quad \left. \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \bar{1} \right\} \end{aligned}$$

Since ξ satisfies the given condition, thus

$$\xi \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \bar{1} \right\} = \bar{1}$$

and thus

$$\begin{aligned}
 \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{p}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{q}} \\
 &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{p}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{q}} \\
 \Rightarrow \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{1}-\bar{p}-\bar{q}} &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{p}+\bar{q}} \\
 \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\frac{\bar{p}+\bar{q}}{\bar{1}-\bar{p}-\bar{q}}} \\
 \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{p}+\bar{q}}{\bar{1}-\bar{p}-\bar{q}} \\
 \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n-2}}^{i_{n-2}}) \right\}^{\bar{\eta}^2} \\
 &\vdots \\
 \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_1}^{i_1}, \tilde{T}_{\alpha_0}^{i_0}) \right\}^{\bar{\eta}^n}.
 \end{aligned}$$

For any $m > n$, where $m, n \in \mathbb{N}$

$$\begin{aligned}
 \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\
 &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\
 &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_{n+3}}^{i_{n+3}}) \right\} \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\
 &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+1}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+2}} \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{m-1}} \\
 &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n(1+\bar{\eta}+\bar{\eta}^2+\dots+\bar{\eta}^{m-n-1})}} \\
 &\leq \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\frac{\bar{\eta}^n}{1-\bar{\eta}}}.
 \end{aligned}$$

Since $\bar{p} + \bar{q} < \frac{1}{2}$, thus $\bar{\eta} < \bar{1}$ and hence $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) \rightarrow \bar{1}$ as $m, n \rightarrow \infty$. So, the soft sequence $\{\tilde{T}_{\alpha_n}^{i_n}\}$ is soft multiplicative Cauchy sequence in \tilde{I} . Being the completeness of $(\tilde{I}, \tilde{\rho}^*, \Omega)$, there exists a soft point $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$ such that $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$ as $n \rightarrow \infty$.

Also,

$$\begin{aligned}
 \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} &\geq \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), (h, \psi)(\tilde{T}_{\alpha_n}^{i_n})\} \left\{ \tilde{\rho}^*(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}) \right\} \right] \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} \\
 &\quad \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n})\} \right]^{\bar{p}} \\
 &\leq \frac{\left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*})\} \right]^{\bar{q}} \left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*})\} \right]}{\sqrt[n]{\left[\tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n})\}, \right. \\
 &\quad \left. \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*})\} \right]}
 \end{aligned}$$

$$\begin{aligned} & \left\{ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{p}} \\ & \lesssim \frac{\left\{ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}} \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right)}{\left[\begin{array}{l} \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \\ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \end{array} \right]} \\ & \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{p}} \\ & \left[\tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}-\bar{q}} \right] \lesssim \frac{\left\{ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}} \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}-\bar{q}}}{\left[\begin{array}{l} \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \\ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \end{array} \right]} \\ & \left[\tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}-\bar{q}} \right] \lesssim \bar{1} \text{ as } n \rightarrow \infty. \end{aligned}$$

Since $\bar{p} + \bar{q} < \frac{1}{2}$, thus $\bar{1} - \bar{p} - \bar{q} > \bar{0}$ and hence $\tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$. This signifies that $\tilde{T}_{\alpha^*}^{i^*}$ is a “soft fixed point” of (h, ψ) .

Now, if $\tilde{T}_{\alpha'}^{i'}$ be another “soft fixed point” of (h, ψ) . Then,

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right) \right\} \\ & \left[\tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\} \right]^{\bar{p}} \\ & \lesssim \frac{\left[\left\{ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\} \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{q}}}{\left[\begin{array}{l} \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\}, \\ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \end{array} \right]} \\ & = \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{2\bar{q}}}{\left[\begin{array}{l} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right\} \end{array} \right]} \\ & \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}-2\bar{q}} \lesssim \bar{1} \end{aligned}$$

Since $\bar{p} + \bar{q} < \frac{1}{2}$ and $\bar{p} > \bar{0}$, thus $\bar{1} - 2\bar{q} > \bar{0}$ and hence $\tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}$.

Hence, there is one and only one soft fixed point of (h, ψ) .

Theorem 3.2. Let $(\tilde{I}, \tilde{\rho}^*, \Omega)$ be a complete soft multiplicative metric space and $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$ be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping $\forall \tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j \in SP(\tilde{I})$,

$$\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), (h, \psi)(\tilde{T}_\beta^j) \right\} \lesssim \frac{\left[\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j \right\} \right]^{\bar{q}} \left[\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j \right\} \right]^{\bar{r}}}{\xi \left[\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j \right\} \right]}$$

where \bar{p} , \bar{q} and \bar{r} are non-negative soft real number such that $\bar{p} + 2\bar{q} + 2\bar{r} < \bar{1}$ and $\xi: [\bar{1}, \infty) \rightarrow [\bar{1}, \infty)$ is a continuous function such that $\xi(\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\kappa^m) = \bar{1}$ iff one of $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\kappa^m = \bar{1}$. Then, there exists a unique fixed point of (h, ψ) .

Proof. Let $\tilde{T}_{\alpha_0}^{i_0}$ be any soft point in $SP(\tilde{I})$. Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi)\tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi)\tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi)\tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) &= \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\} \\ &\quad \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left[\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n} \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\} \right]^{\bar{q}} \\ &\lesssim \frac{\left[\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_n}^{i_n} \right\} \right]^{\bar{r}}}{\xi \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n} \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\}, \right.} \\ &\quad \left. \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_n}^{i_n} \right\} \right]} \\ &\quad \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{q}} \\ &\lesssim \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{r}}}{\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \right.} \\ &\quad \left. \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}} \\ &\quad \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{q}} \\ &\lesssim \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{r}}}{\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \right.} \\ &\quad \left. \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \bar{1} \right\}} \end{aligned}$$

Since ξ satisfies the given condition, thus

$$\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \bar{1} \right\} = \bar{1}$$

and thus

$$\begin{aligned} & \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{1}-\bar{q}-\bar{r}} \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}+\bar{q}+\bar{r}} \\ \Rightarrow & \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\frac{\bar{p}+\bar{q}+\bar{r}}{\bar{1}-\bar{q}-\bar{r}}} \\ \Rightarrow & \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{p}+\bar{q}+\bar{r}}{\bar{1}-\bar{q}-\bar{r}} \\ \Rightarrow & \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n-2}}^{i_{n-2}} \right) \right\}^{\bar{\eta}^2} \\ & \quad \vdots \\ \Rightarrow & \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_1}^{i_1}, \tilde{T}_{\alpha_0}^{i_0} \right) \right\}^{\bar{\eta}^n}. \end{aligned}$$

For any $m > n$, where $m, n \in \mathbb{N}$

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) & \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_{n+3}}^{i_{n+3}} \right) \right\} \dots \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n+1}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n+2}} \dots \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{m-1}} \\ & \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n \left(\bar{1} + \bar{\eta} + \bar{\eta}^2 + \dots + \bar{\eta}^{m-n-1} \right)}} \\ & \lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\frac{\bar{\eta}^n}{\bar{1}-\bar{\eta}}}. \end{aligned}$$

Since $\bar{p} + 2\bar{q} + 2\bar{r} < \bar{1}$, thus $\bar{\eta} < \bar{1}$. Therefore, $\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) \rightarrow \bar{1}$ as $m, n \rightarrow \infty$. So, the soft sequence $\{\tilde{T}_{\alpha_n}^{i_n}\}$ is soft multiplicative Cauchy sequence in \tilde{I} . Being the completeness of $(\tilde{I}, \tilde{\rho}^*, \Omega)$, there exists a soft point $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$ such that $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$ as $n \rightarrow \infty$.

Also,

$$\left[\tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}-\bar{r}} \lesssim \bar{1} \text{ as } n \rightarrow \infty.$$

Since $\bar{p} + 2\bar{q} + 2\bar{r} < \bar{1}$ and $\bar{p} > \bar{0}$, therefore $\bar{1} - \bar{q} - \bar{r} > \bar{0}$ and hence $\tilde{T}_{\alpha^*}^{i^*}$ is a “soft fixed point” of (h, ψ) .

Now, if $\tilde{T}_{\alpha'}^{i'}$ be another “soft fixed point” of (h, ψ) . Then,

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) & = \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right\} \\ & \quad \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left[\tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right\} \right]^{\bar{q}} \\ & \lesssim \frac{\left[\left\{ \tilde{\rho}^* \left((h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\} \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right\} \right]^{\bar{r}}}{\xi \left[\begin{array}{l} \left\{ \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right\} \right\}, \\ \left\{ \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right\} \right\} \end{array} \right]} \end{aligned}$$

$$\begin{aligned} & \left\{ \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'}) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}) \tilde{\rho}^* (\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'}) \right\}^{\bar{q}} \\ &= \frac{\left\{ \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'}) \right\}^{\bar{p} + \bar{q}}}{\xi \left[\begin{array}{l} \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'}), \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}), \tilde{\rho}^* (\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'}) \\ \left\{ \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'}) \right\}, \tilde{\rho}^* (\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*}) \end{array} \right]} \\ & \left\{ \tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'}) \right\}^{\bar{1} - \bar{p} - \bar{2}\bar{r}} \lesssim \bar{1} \end{aligned}$$

Since $\bar{p} + 2\bar{q} + 2\bar{r} < \bar{1}$ and $\bar{q} > \bar{0}$, therefore $\bar{1} - \bar{p} - \bar{r} > \bar{0}$. Thus, $\tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'}) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}$.

Hence, there is one and only one soft fixed point of (h, ψ) .

Theorem 3.3. Let $(\tilde{I}, \tilde{\rho}^*, \Omega)$ be a complete soft multiplicative metric space and $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$ be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping:

$$\begin{aligned} & \left[\max \left\{ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\alpha}^i) \right\}, \tilde{\rho} \left\{ \tilde{T}_{\beta}^j, (h, \psi) (\tilde{T}_{\beta}^j) \right\}, \tilde{\rho}^* (\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j) \right\} \right]^{\bar{p}} \\ \tilde{\rho}^* \left\{ (h, \psi) (\tilde{T}_{\alpha}^i), (h, \psi) (\tilde{T}_{\beta}^j) \right\} & \lesssim \frac{\left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\beta}^j) \right\} \tilde{\rho}^* \left\{ (\tilde{T}_{\beta}^j), (h, \psi) (\tilde{T}_{\alpha}^i) \right\} \right]^{\bar{q}}}{\xi \left[\begin{array}{l} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\alpha}^i) \right\}, \tilde{\rho} \left\{ \tilde{T}_{\beta}^j, (h, \psi) (\tilde{T}_{\beta}^j) \right\}, \tilde{\rho}^* (\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j) \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\beta}^j) \right\}, \tilde{\rho}^* \left\{ (\tilde{T}_{\beta}^j), (h, \psi) (\tilde{T}_{\alpha}^i) \right\} \end{array} \right]} \forall \tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j \in SP(\tilde{I}), \end{aligned}$$

where \bar{p}, \bar{q} are non-negative soft real numbers such that $\bar{p} + 2\bar{q} < \bar{1}$ and $\xi : [\bar{1}, \infty)^2 \rightarrow [\bar{1}, \infty)$ is a continuous function such that $\xi(\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j, \tilde{T}_{\gamma}^k, \tilde{T}_{\delta}^l, \tilde{T}_{\kappa}^m) = \bar{1}$ iff one of $\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j, \tilde{T}_{\gamma}^k, \tilde{T}_{\delta}^l, \tilde{T}_{\kappa}^m = \bar{1}$. Then, there exists a unique fixed point of (h, ψ) .

Proof. Let $\tilde{T}_{\alpha_0}^{i_0}$ be any soft point in $SP(\tilde{I})$. Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi) \tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi) \tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi) \tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\rho}^* (\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &= \tilde{\rho}^* \left\{ (h, \psi) (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), (h, \psi) (\tilde{T}_{\alpha_n}^{i_n}) \right\} \\ & \left[\max \left\{ \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), (h, \psi) (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}, \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_n}^{i_n}), (h, \psi) (\tilde{T}_{\alpha_n}^{i_n}) \right\}, \tilde{\rho}^* (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\} \right]^{\bar{p}} \\ & \lesssim \frac{\left[\tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), (h, \psi) (\tilde{T}_{\alpha_n}^{i_n}) \right\} \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_n}^{i_n}), (h, \psi) (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\} \right]^{\bar{q}}}{\xi \left[\begin{array}{l} \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), (h, \psi) (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}, \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_n}^{i_n}), (h, \psi) (\tilde{T}_{\alpha_n}^{i_n}) \right\}, \tilde{\rho}^* (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \\ \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), (h, \psi) (\tilde{T}_{\alpha_n}^{i_n}) \right\}, \tilde{\rho}^* \left\{ (\tilde{T}_{\alpha_n}^{i_n}), (h, \psi) (\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\} \end{array} \right]} \end{aligned}$$

$$\begin{aligned} & \left[\max \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\} \right]^{\bar{p}} \\ & \leq \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{q}}}{\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \right.} \\ & \quad \left. \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}} \\ & \quad \left[\max \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \right]^{\bar{p}} \\ & \leq \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}}}{\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \bar{1} \right\}} \end{aligned}$$

Since ξ satisfies the given condition, thus

$$\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \bar{1} \right\} = \bar{1}$$

and thus

$$\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq M^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}}$$

where $M = \max \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}$

Now, two cases will be arised:

CASE 1. If $M = \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right)$, then

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}} \\ \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}-\bar{p}-\bar{q}}} & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{q}} \\ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\bar{q}}{\bar{1}-\bar{p}-\bar{q}}} \\ \Rightarrow \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{q}}{\bar{1}-\bar{p}-\bar{q}} \\ \Rightarrow \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-2}}^{i_{n-2}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{\eta}^2} \\ & \vdots \\ \Rightarrow \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n}. \end{aligned}$$

CASE 2. If $M = \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right)$, then

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}} \\ \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{1-\bar{q}} & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{p}+\bar{q}} \\ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) & \leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\bar{p}+\bar{q}}{1-\bar{q}}} \end{aligned}$$

$$\begin{aligned} \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{p} + \bar{q}}{1 - \bar{q}} \\ \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-2}}^{i_{n-2}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{\eta}^2} \\ &\vdots \\ \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^n}. \end{aligned}$$

Using both the cases, for any $m > n$, where $m, n \in \mathbb{N}$, we have

$$\begin{aligned} \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_{n+3}}^{i_{n+3}}) \right\} \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+1}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+2}} \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{m-1}} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n(\bar{1} + \bar{\eta} + \bar{\eta}^2 + \dots + \bar{\eta}^{m-n-1})}} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\frac{\bar{\eta}^n}{1 - \bar{\eta}}}. \end{aligned}$$

Since $\bar{p} + 2\bar{q} < \bar{1}$, therefore $\bar{\eta} < \bar{1}$ and hence $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) \rightarrow \bar{1}$ as $m, n \rightarrow \infty$. So, the soft sequence $\{\tilde{T}_{\alpha_n}^{i_n}\}$ is soft multiplicative Cauchy sequence in \tilde{I} . Being the completeness of $(\tilde{I}, \tilde{\rho}^*, \Omega)$, there exists a soft point $\tilde{T}_{\alpha^*}^{i^*} \tilde{\in} \tilde{I}$ such that $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$ as $n \rightarrow \infty$.

Also,

$$\begin{aligned} \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})) &\lesssim \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})) \\ &\lesssim \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \tilde{\rho}^*(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), (h, \psi)(\tilde{T}_{\alpha^*}^{i^*}) \\ &\tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \max \left[\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi)(\tilde{T}_{\alpha_n}^{i_n})), \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}) \right]^{\bar{p}} \\ &\lesssim \frac{\left[\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})) \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha_n}^{i_n})) \right]^{\bar{q}}}{\xi \left[\begin{array}{l} \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi)(\tilde{T}_{\alpha_n}^{i_n})), \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}) \\ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})), \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha_n}^{i_n})) \end{array} \right]} \\ &\tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \max \left[\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}), \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}) \right]^{\bar{p}} \\ &\lesssim \frac{\left[\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})) \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right]^{\bar{q}}}{\xi \left[\begin{array}{l} \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}), \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})), \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}) \\ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi)(\tilde{T}_{\alpha^*}^{i^*})), \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \end{array} \right]} \end{aligned}$$

$$\Rightarrow \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}} \rightarrow \bar{1} \text{ as } n \rightarrow \infty.$$

Since $\bar{p} + 2\bar{q} < \bar{1}$ and $\bar{p} > \bar{0}$, therefore $\bar{1} - \bar{q} > \bar{0}$ which indicates that $\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$ and hence $\tilde{T}_{\alpha^*}^{i^*}$ is a “soft fixed point” of (h, ψ) .

Now, if $M_2 = \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}$. Then, from equation (3.1), we have

$$\begin{aligned} \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}} &\leq \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{q}} \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ &\leq \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}}{\xi \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ &\leq \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{q}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ &\leq \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}}{\xi \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ &\rightarrow \bar{1} \text{ as } n \rightarrow \infty. \end{aligned}$$

Since $\bar{p} + 2\bar{q} < \bar{1}$, therefore $\bar{1} - \bar{q} - \bar{p} > \bar{0}$ which indicates that $\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$ and hence $\tilde{T}_{\alpha^*}^{i^*}$ is a “soft fixed point” of (h, ψ) .

Thus, in all the cases, we get $\tilde{T}_{\alpha^*}^{i^*}$ as a “soft fixed point” of (h, ψ) . Now, if $\tilde{T}_{\alpha'}^{i'}$ be another “soft fixed point” of (h, ψ) . Then,

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right\} \\ &= \left[\max \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho} \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right] \right]^{\bar{p}} \\ &= \frac{\left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right\} \right]^{\bar{q}}}{\xi \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho} \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right]} \\ &= \frac{\left[\max \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho} \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\} \right]^{\bar{p}}}{\xi \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]^{\bar{q}}} \\ &= \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho} \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}}{\xi \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}} \\ &= \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left\{ \bar{1}, \bar{1} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}} \\ &\leq \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}+2\bar{q}} \end{aligned}$$

$$\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{1} - \bar{p} - 2\bar{q}} \leq \bar{1}$$

Since $\bar{p} + 2\bar{q} < \bar{1}$, therefore $\bar{1} - \bar{p} - 2\bar{q} > \bar{0}$. Thus, $\tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}$.

Hence, there is one and only one soft fixed point of (h, ψ) .

Theorem 3.4. Let $(\tilde{I}, \tilde{\rho}^*, \Omega)$ be a complete soft multiplicative metric space and $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$ be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping

$$\tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha}^i \right), (h, \psi) \left(\tilde{T}_{\beta}^j \right) \right\} \leq \frac{\left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) \left(\tilde{T}_{\beta}^j \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) \left(\tilde{T}_{\alpha}^i \right) \right\} \right]^{\frac{\bar{p}}{2}} \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) \left(\tilde{T}_{\beta}^j \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) \left(\tilde{T}_{\alpha}^i \right) \right\} \right]^{\frac{\bar{q}}{2}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j \right) \right\}^{\bar{r}}}{\xi \left[\begin{array}{l} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) \left(\tilde{T}_{\alpha}^i \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) \left(\tilde{T}_{\beta}^j \right) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) \left(\tilde{T}_{\beta}^j \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) \left(\tilde{T}_{\alpha}^i \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j \right) \end{array} \right]} \quad \forall \tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j \in SP(\tilde{I}),$$

where \bar{p}, \bar{q} and \bar{r} are non-negative soft real numbers such that $\bar{p} + \bar{q} + \bar{r} < \bar{1}$ and $\xi : [\bar{1}, \infty) \rightarrow [\bar{1}, \infty)$ is a continuous function such that $\xi(\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j, \tilde{T}_{\gamma}^k, \tilde{T}_{\delta}^l) = \bar{1}$ iff one of $\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j, \tilde{T}_{\gamma}^k, \tilde{T}_{\delta}^l = \bar{1}$. Then, there exists a unique fixed point of (h, ψ) .

Proof. Let $\tilde{T}_{\alpha_0}^{i_0}$ be any soft point in $SP(\tilde{I})$. Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi) \tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi) \tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi) \tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\} \\ &= \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\} \right]^{\frac{\bar{p}}{2}} \\ &\leq \frac{\left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\} \right]^{\frac{\bar{q}}{2}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{r}}}{\xi \left[\begin{array}{l} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, (h, \psi) \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \end{array} \right]} \end{aligned}$$

Since ξ satisfies the given condition, thus

$$\xi \left[\begin{array}{l} \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \\ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \bar{1}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \end{array} \right] = \bar{1}.$$

Thus, we have

$$\begin{aligned} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1} - \frac{\bar{p} + \bar{q}}{2}} &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\bar{p} + \bar{q} + \bar{r}}{2}} \\ \Rightarrow \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\frac{\bar{p} + \bar{q} + \bar{r}}{2}}{\bar{1} - \frac{\bar{p} + \bar{q}}{2}}} \\ \Rightarrow \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\frac{\bar{p} + \bar{q} + \bar{r}}{2}}{\bar{1} - \frac{\bar{p} + \bar{q}}{2}} \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n-2}}^{i_{n-2}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{\eta}^2} \\ &\vdots \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n}. \end{aligned}$$

For any $m > n$, where $m, n \in \mathbb{N}$

$$\begin{aligned} \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \dots \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n+1}} \dots \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{m-1}} \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n(\bar{1} + \bar{\eta} + \bar{\eta}^2 + \dots + \bar{\eta}^{m-n-1})}} \\ &\lesssim \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\frac{\bar{\eta}^n}{\bar{1} - \bar{\eta}}}. \end{aligned}$$

Since $\bar{p} + \bar{q} + \bar{r} < \bar{1}$, therefore $\bar{\eta} < \bar{1}$ which indicates that $\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) \rightarrow \bar{1}$ as $m, n \rightarrow \infty$. So, the soft sequence $\{\tilde{T}_{\alpha_n}^{i_n}\}$ is soft multiplicative Cauchy sequence in \tilde{I} . Being the completeness of $(\tilde{I}, \tilde{\rho}^*, \Omega)$, there exists a soft point $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$ such that $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$ as $n \rightarrow \infty$.

Also,

$$\begin{aligned} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} &\lesssim \left[\tilde{\rho}^* \left\{ \left(\tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\} \left\{ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right), (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right\} \right\} \right] \\ &\quad \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\frac{\bar{p}}{2}} \\ &\lesssim \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \frac{\left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\} \right]^{\frac{\bar{q}}{2}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{r}}}{\xi \left[\begin{array}{l} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha_n}^{i_n} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \end{array} \right]} \end{aligned}$$

$$\begin{aligned}
 & \left[\tilde{\rho}^*(T_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\frac{\bar{p}}{2}} \\
 & \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right]^{\frac{\bar{q}}{2}} \\
 \lesssim & \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{r}}}{\left[\begin{array}{c} \tilde{\rho}^* \left(T_{\alpha_n}^{i_n}, T_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, T_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \end{array} \right]} \\
 & \left[\tilde{\rho}^* \left(T_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\frac{\bar{p}}{2}} \\
 & \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right]^{\frac{\bar{q}}{2}} \\
 \lesssim & \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{r}}}{\left[\begin{array}{c} \tilde{\rho}^* \left(T_{\alpha_n}^{i_n}, T_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, T_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \end{array} \right]} \\
 \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1} - \frac{\bar{p}}{2} - \frac{\bar{q}}{2}} \lesssim & \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \frac{\left\{ \tilde{\rho}^* \left(T_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\frac{\bar{p}}{2} + \frac{\bar{q}}{2} + \bar{r}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\frac{\bar{p}}{2} + \frac{\bar{q}}{2}}}{\left[\begin{array}{c} \tilde{\rho}^* \left(T_{\alpha_n}^{i_n}, T_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, T_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \end{array} \right]} \\
 \lesssim & \frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\frac{\bar{p}}{2} + \frac{\bar{q}}{2} + \bar{1}} \left[\tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right) \right]^{\frac{\bar{p}}{2} + \frac{\bar{q}}{2} + \bar{r}}}{\left[\begin{array}{c} \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right\}, \\ \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha_n}^{i_n} \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right) \end{array} \right]}
 \end{aligned}$$

Since $\bar{p} + \bar{q} + \bar{r} < \bar{1}$ and $\bar{r} > \bar{0}$, therefore $\bar{1} - \frac{\bar{p}}{2} - \frac{\bar{q}}{2} > \bar{0}$ and hence $\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$ as $n \rightarrow \infty$. This shows that $(h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right) = \tilde{T}_{\alpha^*}^{i^*}$ and hence $\tilde{T}_{\alpha^*}^{i^*}$ is a “soft fixed point” of (h, ψ) .

Now, if $\tilde{T}_{\alpha'}^{i'}$ be another “soft fixed point” of (h, ψ) . Then,

$$\begin{aligned}
 \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left(\tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left(\tilde{T}_{\alpha'}^{i'} \right) \right\} \\
 & \left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\} \right]^{\frac{\bar{p}}{2}} \\
 & \lesssim \frac{\left[\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\frac{\bar{q}}{2}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{r}}}{\left[\begin{array}{c} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\}, \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \end{array} \right]}
 \end{aligned}$$

$$\frac{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{r}}} = \frac{\xi \left[\begin{array}{l} \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right), \\ \tilde{\rho}^* \left(\tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \end{array} \right]}{\left\{ \tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{1}-\bar{q}-\bar{r}}} \lesssim \bar{1}$$

Since $\bar{p} + \bar{q} + \bar{r} < \bar{1}$ and $\bar{p} > \bar{0}$, therefore $\bar{1} - \bar{q} - \bar{r} > \bar{0}$ and hence $\tilde{\rho}^* \left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}$.

Hence, there is one and only one soft fixed point of (h, ψ) .

4 Conclusion

“Soft set theory” is a wide mathematical aid for handling vagueness and uncertainty. In this paper, some basic concepts of soft set and soft metric spaces are considered. We proved fixed point theorem for mappings satisfying generalized weak contractive conditions in soft multiplicative metric space.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Molodtsov DA. Soft set theory-first result. Computers and Mathematics with Applications. 1999; 37: 19-31.
- [2] Maji PK, Roy AR, Biswas R. An application of soft real number in a decision making problem. Computers and Mathematics with Applications. 2002; 42:1077-1083.
- [3] Maji PK, Biswas R, Roy AR. Soft set theory. Computers and Mathematics with Applications. 2003; 45:555-562.
- [4] Ali MI, Feng F, Liu X, Min WK, Shabir M. On some new operation in soft set theory. Computer and Mathematics with Applications. 2011; 62:351-358.
- [5] Wardowski D. On a soft mapping and its fixed points. Fixed Point Theory and Applications. 2013;182: 1-11.
- [6] Das S, Samanta SK. On soft metric spaces. Journal of Fuzzy Mathematics. 2013; 21:207-213.
- [7] Das S, Samanta SK. Soft real sets, soft real numbers and their properties. Journal of Fuzzy Mathematics. 2012; 20:551-576.
- [8] Bashirov AE, Kurpirnar EM, Ozgapiri A. Multiplicative calculus and its applications. Journal of Mathematical Analysis and Applications. 2008; 337:36-48.
- [9] Abodayeh K, Pitea A, Shatanavi W, Abdegawad T. Remarks on multiplicative metric spaces and related fixed points. 2015; arXiv:1512.03771v1:1-7.

- [10] Ozavsar M, Cevikel AC. Fixed Point of multiplicative contraction mappings on multiplicative metric space. 2012; arXiv:1205.5131v1,1-14.
- [11] Agrawal RP, Karapinar E, Samet B. An essential remark on fixed point results on multiplicative metric spaces. Fixed Point Theory and Applications. 2016; 21:1-3.
- [12] Dosenovic T, Pastolache M, Radenovic S. On multiplicative metric spaces: survey. Fixed Point Theory and Applications. 2016; 92:1-17.
- [13] Dosenovic T, Pastolache M, Radenovic S. Some essential remark on fixed point results on multiplicative metric space. Journal of Advanced Mathematical Studies. 2017;10(1):20-24.
- [14] He X, Song M, Chen D. Common fixed points for weak commutative mappings on a multiplicative metric space. Fixed Point Theory and Applications. 2014; 48:1-9.
- [15] Rathee S, Girdhar R, Dhingra K. Fixed point theorems in soft multiplicative metric spaces. Communications in Mathematics and Applications. 2020; 11:425-442.
- [16] Wadkar BR, Mishra LN, Bhardwaj R, Singh B. Fixed point results related to soft sets. Australian Journal of Basic and Applied Sciences. 2016;10(16):128-137.
- [17] Yazar MI, Gunduz C, Bayramov S. Fixed point theorems of soft contractive mappings. Filomat. 2016; 30:269-279.
- [18] Hosseinzadeh H. Fixed point theorem on soft metric spaces. Journal of Fixed Point Theory and Applications. 2017; 18(4):1-22.
- [19] Abbas M, Murtaza G. On the fixed point theory of soft metric spaces. Fixed Point Theory and Applications. 2016; 17:1-11.
- [20] Bhardwaj R, Singh S, Pandey S, Wadkar BR. Fixed point results in soft metric space. International Journal of Modern Agriculture. 2021;10(2):1792-1803.
- [21] Solanki S, Bhardwaj R, Singh BK, Sharma M. Fixed point theory of soft weak contraction on a complete soft metric space. International Journal of Advanced Engineering Science and Technological Research. 2017;41(1):425-432.

© 2023 Nagpal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/95326>