



Profile of Anxiety and Depression Symptoms Associated with Sleep Quality of Medical Students in the Eastern Amazon

**Vitor Maués Lopes ^{a*}, Ismaelino Mauro Nunes Magno ^a,
Narely Araújo Smith ^b, Valeria Francês Nascimento ^a
and Vitor Hugo Auzier Lima ^{c,d*}**

^a Department of Medicine, CESUPA - University Center of the State of Pará, Brazil.

^b Department of Medicine, UFPA - Federal University of Pará, Brazil.

^c Department of Biotechnology, UFPA - Federal University of Pará, Brazil.

^d Department of Physiotherapy, UEPA - State University of Pará, Pará, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. Author VML was the majority author of this manuscript, since he carried out the research and wrote in his majority. Author VHAL was the correspondent of the magazine, reviser of this work and updated its references. Author NAS was the third reviewer, translator and assistant in bibliographical research. Author IMNM was research supervisor on bachelor of medicine. Author VFN was scientific advisors and payers of this article. Authors IMNM, VML and VHAL managed the feasibility analyzes of the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/INDJ/2023/v19i1363

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96652>

Original Research Article

Received: 02/01/2023
Accepted: 28/01/2023
Published: 28/01/2023

*Corresponding author: E-mail: vitor.maués@yahoo.com, vitorauzier@hotmail.com;

ABSTRACT

Objective: An adequate sleep is essential for good mental health. Thus, this study associated the prevalence of the main symptoms of anxiety and depression with sleep quality in medical students at a Brazilian university center, a population frequently affected by these disorders.

Methods: This is a cross-sectional, descriptive and analytical study, with data obtained through online questionnaires answered by medical students, during the year 2022. To assess symptoms of depression and anxiety, the Hospital Anxiety and Depression Scale (HADS) was used, while, to assess the quality of sleep, the Pittsburgh Sleep Quality Index (PSQI).

Results: 229 students participated in the study. Of these, 51% had anxiety symptoms. As for depression, 27,5% had depression symptoms. With regard to sleep quality, 28% had poor sleep quality and 71% good quality. There was a significant association between sleep quality and anxiety and depression symptoms ($p < 0.001$): of the individuals with good sleep quality, 51% had possible and probably anxiety and 21,2% were classified as possible and probably depression. Furthermore, about the students with poor sleep quality: 80% were possible and probably anxious students and 43,8% were possible and probably depressive.

Conclusions: The study documents a statistically significant association between anxiety and depression symptoms and poor sleep quality. Further multicentric analyzes are needed to understand this scenario.

Keywords: Anxiety; depression; sleep; students; medical.

1. INTRODUCTION

Medical students have always been target of several studies that evaluate mental health, due to the great amount of content to which they are exposed and, consequently, to greater pressure and time dedicated to studies [1]. Thus, such students have a higher prevalence of mental disorders, such as depression and anxiety, when compared to university students from other areas. Their quality of life was also considered worse than that of individuals in the same age group [2].

Furthermore, sleep is one of the most relevant topics nowadays, as there is consistent evidence that deprivation influences metabolic and inflammatory processes with significant negative impacts on human health [3,4], and may also act as an early marker of anxiety and depression [5]. In general, medical students present susceptibility to unsatisfactory sleep, reducing their sleep time to support their academic obligations, corroborating that poor sleep has a higher prevalence in them compared to the general population [1,2,5].

Researches indicates that sleep disturbances tend to co-occur with the rise in mental health disorders in college students. Considering the increased prevalence of poor sleep quality in medical students, especially after the COVID-19 pandemic [6], little is known about the current

relationship between sleep quality and mental health in medical students [7].

Therefore, in view of this scenario and the increasing number of sleep disorders, it is necessary to increase the database for an effective understanding of the harmful relationship between these disorders and mental health symptoms today, as already pointed out by other studies [8]. Therefore, the present study aimed to establish the prevalence and relationship between anxiety and depression symptoms and sleep quality in medical students at a university center in northern Brazil.

2. METHODS

2.1 Study Design

This study is a cross-sectional, analytical and observational study to obtain data through the questionnaires answered by students enrolled in the Medicine Course of the University Center of the State of Pará-Brazil, in the year 2022. From this, it was analyzed data on the prevalence of anxiety and depression symptoms in medical students and compared them with sleep quality scores, seeking to identify possible relationships between these conditions.

Data collection through questionnaires was authorized by the center, through approval by the Institution's Research Ethics Committee. The

questionnaires had 3 sections, the first one with sociodemographic questions, the second with the Hospital Anxiety and Depression Scale (HADS) and the third with the Pittsburgh Sleep Quality Index (PSQI), which will be discussed later. In addition, data collection was authorized by the students, by signing the Informed Consent Form contained on the first page of the questionnaire.

2.2 Sample Selection

This analysis included all medical students from the university center, from the 1st to the 12th semester, who agreed to answer the questionnaires and who signed the Informed Consent Form. Exclusion criteria were not adopted.

The study's university medical center has about 50 students each semester. However, considering dropouts, failures and other variables, 48 students were considered per class. Thus, totaling 576 students enrolled and obtaining a calculated sample size of 231 students, with a confidence level of 95% and a margin of error of 5%.

In the second section of the questionnaire, there were sociodemographic questions, such as age, gender and stage in the medical school. To approach the participants, the calculated sample value of 231 students was considered, and a stratified random sample was chosen with a minimum objective of 19 students in each semester, who were approached during their academic activities by the researchers. However, given the difficulties in finding younger students, the collection included 17 students from the first 4 semesters, 18 from the 5th semester, 19 from the 6th, 7th and 8th semesters, in addition to 22 from the 9th semester and 21 from the 10th, 11th semesters and 12.

The students were considered per biennium, as the medical education in Brazil is divided into stages, with the first two years considered the "basic cycle", years 3 and 4 the "clinical cycle", and years 5 and 6 "internship", usually composed exclusively of internships and practices.

2.3 Data Collection and Analysis

To subjectively assess sleep quality, an excellent recognized tool is the "Pittsburgh Sleep Quality Index (PSQI)" of 1989 [9]. The questionnaire evaluates seven sleep components: subjective

quality of sleep, sleep latency, sleep duration, sleep efficiency, sleep disorders, medication use, and daytime dysfunction. The score can range from 0 to 3 for each component, with a maximum total score of 21 points. Scores below 5 indicate good quality of sleep and above that indicate poor quality of sleep. This index was validated for the Portuguese language [10] and is considered accessible to identify sleep quality, identifying whether individuals sleep well, poorly or have sleep disorders [11]. The PSQI showed strong reliability, with moderate structural validity in different samples, suggesting that the tool fulfills its intended utility [12].

To assess symptoms of anxiety and depression, the Hospital Anxiety and Depression Scale (HADS) [13] validated for the Portuguese language [14] was used. The scale evaluates subjective symptoms of anxiety and depression and does not confirm these diseases or even replace the evaluation of a specialist professional. The HADS consists of 14 items divided into two subscales, 7 questions to assess depression and 7 to assess anxiety. Each item is scored from 0 to 3, with a maximum score of 21 for anxiety or depression. The recommended scores according to Zigmond and Snaith [13] are 8 to 10 for possible cases and ≥ 11 for probable cases. Scores less than 7 indicate non-cases.

The choice to use the HADS scale was due to its easy completion, which allowed for quick and reliable data collection, and the great advantage of the absence of confusing symptoms common between clinical illnesses and depression or anxiety, such as fatigue, loss of appetite and sleep disturbances [15]. Furthermore, the scale proved to be potentially effective and applicable for screening and detection of mild anxiety and depression disorders [16,17,18].

Data were collected through online questionnaires after Institution's Research Ethics Committee approval, from June to August 2022. They were organized in Microsoft Excel 2010. Graphs and tables were built with the tools available in Microsoft Word, Excel and Bioestat 5.5 softwares.

All tests were performed using the Bioestat 5.5 software. Quantitative variables were described by frequency and percentage. 95% confidence intervals were calculated for the proportion to infer how the prevalences behave in relation to the population from which they were obtained. The independence or association between two

categorical variables was tested using the chi-square or Fisher's exact test, depending on the case, and significant associations were detailed using the analysis of standardized residues, to identify the categories that most contributed to the result. Results with $p \leq 0.05$ were considered statistically significant.

3. RESULTS

A total of 229 students were included in the study, 7 students refused to participate. Most of whom (142 or 62%) were female and 37.6% were male. Most (142 or 62%) were aged between 21 and 25 years and 25.3% of individuals were aged between 17 and 20 years (Table 1).

Considering the students per biennium, 37.6% of the individuals were from the 9th to the 12th semester, 75 individuals (32.8%) were from the 5th to the 8th semester and 86 (37.6%) were from the 9th to the 12th semester (Table 1).

As for the classification of anxiety and depression symptoms, 94 individuals (41%) had an improbable classification of anxiety and 37.6% of individuals had possible anxiety. Most (166 or 72.5%) had unlikely depression, 49 subjects (21.4%) had possible depression, and 6.1% (14 subjects) had probable depression. 64 (27.9%) had poor quality sleep. The narrower this interval is, the greater the certainty related to that prevalence in the population from which this sample was obtained (Table 2).

Table 1. Sociodemographic characteristics of the students, evaluated from June to August 2022, Belém-Pará

Variable	Frequency	Percentage
Sex		
Female	142	62.0
Masculine	86	37.6
Other	1	0.4
Age		
From 17 to 20 years old	58	25.3
From 21 to 25 years old	142	62.0
From 26 to 30 years old	20	8.7
Biennium in the medical school		
1st to 2nd year	68	29.7
3rd to 4th year	75	32.8
5th to 6th year	86	37.6

Percentages are relative to the total number of participants (n=229)

Table 2. Classification regarding anxiety and depression (HADS) and sleep quality (PSQI) of the students, evaluated from June to August 2022, Belém-Pará

Variable	Frequency	Percentage	CI95%
Anxiety Classification (HADS)			
Unlikely	94	41.0	34.7 - 47.7
Possible	86	37.6	31.3 - 44.2
Likely	49	21.4	16.4 - 27.4
Depression Classification (HADS)			
Unlikely	166	72.5	66.1 - 78.1
Possible	49	21.4	16.4 - 27.4
Likely	14	6.1	3.5 - 10.3
Sleep Quality (PSQI)			
Good sleep	165	72.1	65.7 - 77.7
bad sleep	64	27.9	22.3 - 34.3

Percentages are relative to the total number of participants (n=229). 95% CI: 95% confidence interval for the prevalence

Table 3 showed the association between sleep quality (PSQI), age, gender and stage in the medical school. In individuals with good sleep quality, most (104 or 63%) were aged 21 to 25 years and 26.1% of individuals were aged 17 to 20 years. In the group with poor sleep quality, more than half (38 or 59.4%) were aged 21 to 25 years and 23.4% of subjects were aged 17 to 20 years. However, these observed differences did not reach statistical significance ($p=0.353$). Sleep quality and sex were significantly associated ($p=0.007$): of the 64 individuals with poor sleep quality, 78.1% were female, in a higher proportion than expected; of the individuals with good sleep quality, 43.6% were male, a higher proportion than expected by the statistical test.

That is, the quality of sleep among women was significantly worse. Furthermore, according to the data obtained, sleep quality and stage in the medical school were not significantly associated ($p=0.064$).

Table 4 showed the association between sleep quality, anxiety classification and depression classification. There was a significant association between sleep quality and anxiety ($p<0.001$): of the individuals with good sleep quality, 81 (49.1%) had improbable anxiety, this proportion being higher than expected by the statistical test; of the 64 individuals with poor sleep quality, 24 (37.5%) were likely, this proportion being higher than expected by the statistical test.

Table 3. Association between age, gender and stage in the medical school with sleep quality of the students, evaluated from June to August 2022, Belém-Pará

Variable	Good Sleep (n=165)	Bad Sleep (n=64)	p-value
Age			0.353
From 17 to 20 years old	43 (26.1)	15 (23.4)	
From 21 to 25 years old	104 (63.0)	38 (59.4)	
From 26 to 30 years old	11 (6.7)	9 (14.1)	
From 31 to 39 years old	7 (4.2)	2 (3.1)	
Sex			0.007
Female	92 (55.8)*	50 (78.1)†	
Masculine	72 (43.6)†	14 (21.9)*	
Biennium in the medical school			0,064
1st to 2nd year	56 (33,9)	12 (18,8)	
3rd to 4th year	49 (29,7)	26 (40,6)	
5th to 6th year	60 (36,4)	26 (40,6)	

*Categorical variables are displayed as n (%). The percentages are relative to the total of each column. In all cases, the chi-square was used. *: this frequency was lower than what would be expected by chance. †: this frequency was higher than expected*

Table 4. Association between anxiety classification (HADS), depression classification (HADS) and sleep quality of the students, evaluated from June to August 2022, Belém-Pará

Variable	Good Sleep (n=165)	Bad Sleep (n=64)	p-value
Anxiety Classification (HADS)			<0.001
Unlikely	81 (49.1)†	13 (20.3)*	
Possible	59 (35.8)	27 (42.2)	
Likely	25 (15.2)*	24 (37.5)†	
Depression Classification (HADS)			<0.001
Unlikely	130 (78.8)†	36 (56.3)*	
Possible	30 (18.2)	19 (29.7)	
Likely	5 (3.0)*	9 (14.1)†	

*Categorical variables are displayed as n (%). The percentages are relative to the total of each column. In all cases, the chi-square was used. *: this frequency was lower than what would be expected by chance. †: this frequency was higher than expected*

There was also a significant association between sleep quality and depression ($p < 0.001$): of the individuals with good sleep quality, 130 (78.8%) were classified as unlikely depression, in a higher proportion than expected; in terms of poor sleep quality, 9 (14.1%) were likely to have depression, with this proportion being higher (†) than expected.

4. DISCUSSION

In this study, it was identified that 37% of students had possible anxiety, while 21% had probable anxiety [19]. As for depression, 21.4% of individuals had possible depression and 6.1% probable depression. A recent meta-analysis showed that 33.6% of students had depressive symptoms, while 39% had anxiety symptoms [20], being more compatible with the result of possible anxiety and depression. Furthermore, another study showed, using the HADS, that medical students had symptoms of anxiety and depression at a prevalence of 20% and 7% respectively [19,21], which is consistent with the "probable depression and anxiety" results found in the present study.

In a recent meta-analysis, it was identified an overall prevalence rate of anxiety among Brazilian undergraduate students of 37,7% and a prevalence of depression of 28,51% [22]. A later Brazilian meta-analysis has already shown similar compatibility in medical students, with 30.6% for depression and 32.9% for anxiety [23]. Thus, although the sample of the present study is small, of a unicentric nature, and there are different screening tools for these conditions, it was possible to identify similarity with the values obtained in relation to the classification of possible anxiety and depression, guaranteeing reliability in the screening of the symptoms of these conditions through HADS.

When considering the extensive workload demanded in the medical course, many students find the availability to study at night, thus reducing the number of hours reserved for sleep [6,24]. In addition, the impairment of the quality of this sleep due to the stressors of medical graduation, such as study overload, high competitiveness, responsibility and expectations associated with the physician's role, is analyzed [6,24,25,17]. Such contexts become alarming and are connected, since sleep disorders increase the incidence of psychiatric disorders, in addition to reducing the professional and academic performance of individuals [17].

Thus, anxiety symptoms are associated with more sleep disorders and the use of sleeping pills [18,26,27,28,29], with proven polysomnographic changes in these individuals [26]. In the present study, there was a significant association between anxiety symptoms and poor sleep quality ($p < 0.001$), in which 37.5% were probable and 42.2% possible anxious individuals with poor sleep quality. With that in mind, adding these values, about 80% of medical students with anxiety symptoms had poor sleep quality.

In other studies [27-29], poor sleep quality was associated with anxiety and depression. In the present study, which also showed a relevant association ($p < 0.001$), 29.7% had probable depression and 14.1% possible, totaling 43.8% of the total number of students with depressive symptoms and poor sleep quality.

In this sense, understanding that insomnia can behave as a predictor of anxiety [30] and depression [30-32] and that it can also be a symptom of both disorders - according to the DSM-5 [33] - it becomes difficult to establish a direct causal relationship. However, it is proven that cognitive-behavioral therapy for sleep disorders improves mental health and the severity of symptoms in patients with mental disorders, including depression [34]; making sleep treatment an important front not only for the prevention but also for the treatment of these disorders.

Although the study did not identify a significant association between sleep quality and stage in the course or age, sleep quality and gender were significantly associated ($p = 0.007$): of the 64 individuals with poor sleep quality, 78.1% were from the women. Other studies point to this same factor, in which they confirm that sleep disturbances, mainly insomnia, are more prevalent among women due to behavioral aspects and biological constitution [27,35-37].

In spite of the fact that medical students have naturally high rates of mental distress and sleep disturbances, this scenario has become more alarming recently with the COVID-19 pandemic [38-39]. The present study and the values found probably had an important impact from the pandemic, but it was not evaluated. One Brazilian study also showed concern about the negative impact of COVID-19 on medical education and the generation of medical jobs, which can also directly worsen sleep quality [40].

Other markers of mental suffering, such as burnout syndrome, also deserve attention in the medical field. This syndrome occurs not only in students, but also in medical residents, who perpetuate these disorders [41,42]. A recent Brazilian study with resident physicians exposed relevant prevalence of anxiety, depression and burnout, identifying similar patterns and individual and environmental common factors, such as high workloads, occurrence of psychological abuse, poor supervision of the faculty, unsatisfactory learning experience and conflicts between work and personal life [43].

As for the limitations of the study, the use of scales is listed, in which both the HADS - although it suggests, does not confirm anxiety or depression - and the PSQI are subjective tools and can be influenced by memory, inattention or even dishonesty of the participants. In addition, the different scales and other ways of evaluating these disorders used in the studies end up making a completely reliable comparative analysis difficult. It is also possible to point out the failure to establish causality and other confounders associated with sleep quality and/or symptoms of anxiety and depression. Finally, the unicentric character of the study may limit the interpretation of the results.

5. CONCLUSION

In summary, the data from this study contemplate what was outlined as an objective, significantly supporting the harmful relationship between mental health disorders, depression and anxiety, and poor sleep quality. With regard to sociodemographic characteristics, the study also showed a statistically significant association between poor sleep quality and female gender.

The data obtained are relevant for the definition of intervention models, applying them after recognizing the variables that interfere in the connection between mental health disorders and, consequently, bad sleep quality. Further multicenter analyzes are needed to understand this scenario in medical schools around the world.

CONSENT AND ETHICAL APPROVAL

This research was approved by the ethics and research committee of the Centro Universitário do Pará. It has approval opinion number 5.387.147. With respective contact email cep@cesupa.br.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Perotta B, Arantes-Costa FM, Enns SC, Figueiro-Filho EA, Paro H, Santos IS, Lorenzi-Filho G, Martins MA, Tempski PZ. Sleepiness, sleep deprivation, quality of life, mental symptoms and perception of academic environment in medical students. *BMC Med Educ.* 2021 Feb 17; 21(1):111. DOI: 10.1186/s12909-021-02544-8. PMID: 33596885; PMCID: PMC7890911.
2. Barbosa-Medeiros MR, Caldeira AP. Mental health of medical students: longitudinal study. *Rev. Bras. Educate Med.* 2021;45(3). DOI: 10.1590/1981-5271v45.3-20190285.
3. Barros MBA, Lima MG, Ceolim MF, Zancanella, E, Cardoso TAMO. Quality of sleep, health and well-being in a population-based study. *Rev. of Public Health.* 2019;53:82. DOI:10.11606/s1518-8787.2019053001067
4. Jin Q, Yang N, Dai J, Zhao Y, Zhang X, Yin J, Yan Y. Association of sleep duration with all-cause and cardiovascular mortality: A prospective cohort study. *Front Public Health.* 2022 Jul 15;10:880276. DOI: 10.3389/fpubh.2022.880276. PMID: 35910926; PMCID: PMC9334887
5. Rao WW, Li W, Qi H, Hong L, Chen C, Li CY, Ng CH, Ungvari GS, Xiang YT. Sleep quality in medical students: a comprehensive meta-analysis of observational studies. *Sleep Breath.* 2020 Sep;24(3):1151-1165. DOI: 10.1007/s11325-020-02020-5. Epub 2020 Feb 18. PMID: 32072469.
6. Saguem BN, Nakhli J, Romdhane I, Nasr SB. Predictors of sleep quality in medical students during COVID-19 confinement. *Encephale.* 2022 Feb;48(1):3-12. DOI: 10.1016/j.encep.2021.03.001. Epub 2021 Mar 18. PMID: 33863513; PMCID: PMC8828364.
7. Chen J, Tuersun Y, Yang J, Xiong M, Wang Y, Rao X, Jiang S. Association of

- depression symptoms and sleep quality with state-trait anxiety in medical university students in Anhui Province, China: a mediation analysis. *BMC Med Educ.* 2022 Aug 19;22(1):627.
DOI: 10.1186/s12909-022-03683-2.
PMID: 35982434; PMCID: PMC9388213.
8. Milojević HM, Lukowski AF. Sleep and mental health in undergraduate students with generally healthy sleep habits. *PLoS One.* 2016 Jun 9;11(6):e0156372.
DOI: 10.1371/journal.pone.0156372.
 9. Bugueño M, et al. Sleep quality and academic performance in high school students. *Rev. Med. from Chile.* 2017; 145(9):1106-1114.
 10. Bertolazi NA. Translation, cultural adaptation and validation of two sleep assessment instruments: EPWORTH Sleepiness Scale and Pittsburgh Sleep Quality Index. Porto Alegre. Federal University of Rio Grande do Sul; 2008.
 11. Escobar-Córdoba F, Eslava-Schmalbach J. Colombian Validation of the Pittsburgh Sleep Quality Index. *Rev de Neurol.* 2005; 40(3):150-155.
 12. Mollayeva T, Thurairajah P, Burton K, Mollayeva S, Shapiro CM, Colantonio A. The Pittsburgh sleep quality index as a screening tool for sleep dysfunction in clinical and non-clinical samples: A systematic review and meta-analysis. *Sleep Med Rev.* 2016;25:52-73.
DOI: 10.1016/j.smrv.2015.01.009
 13. Zigmund AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand.* 1983;67(6):361-370.
DOI: 10.1111/j.1600-0447.1983.tb09716.x
 14. Apostolo JLA, Mendes AC, Azeredo ZA. Adaptation to Portuguese of the Depression, Anxiety and Stress Scales (DASS). *Rev Lat Am Nursing;* 2006.
 15. Castro MMC, Quarantini L, Batista-Neves S, Kraychete DC, Daltro C, Miranda-Scippa Â. Validity of the hospital anxiety and depression scale in patients with chronic pain. *Rev Bras Anestesiol.* 2006; 56(5):470-477.
DOI: 10.1590/S0034-70942006000500005
 16. Botega NJ, Bio MR, Zomignani MA, Garcia Jr C, Pereira WAB. Mood disorders in a medical clinic ward and validation of an anxiety and depression measurement scale (HAD). *Rev Saude Publica.* 1995; 29(5):359-363.
DOI: 10.1590/S0034-89101995000500004
 17. Karagöl A. Levels of Depression, Anxiety and Quality of Life of Medical Students. *Psychiatr Danub.* 2021 Spring-Summer; 33(Suppl 4):732-737.
PMID: 34718310.
 18. Becker SP, et al. Sleep in a large, multi-university sample of college students: sleep problem prevalence, sex differences, and mental health correlates. *Sleep Health.* 2018;4(2):174-181.
 19. Faro A. Confirmatory Factor Analysis and Normalization of the Hospital Anxiety and Depression Scale (HADS). *Psicol Teor e Pesqui.* 2015;31(3):349-353.
DOI: 10.1590/0102-37722015032072349353
 20. Li W, Zhao Z, Chen D, Peng Y, Lu Z. Prevalence and associated factors of depression and anxiety symptoms among college students: a systematic review and meta-analysis. *J Child Psychol Psychiatry.* 2022 Nov;63(11):1222-1230.
DOI: 10.1111/jcpp.13606. Epub 2022 Mar 16. PMID: 35297041.
 21. Bertani DE, et al. Anxiety, depression and personality traits in Italian medical students. *Rev Psychiatr.* 2020;55(6):342-348.
DOI: 10.1708/3503.34892.
 22. Demenech LM, Oliveira AT, Neiva-Silva L, Dumith SC. Prevalence of anxiety, depression and suicidal behaviors among Brazilian undergraduate students: A systematic review and meta-analysis. *J Affect Disord.* 2021 Mar 1;282:147-159.
DOI: 10.1016/j.jad.2020.12.108.
Epub 2020 Dec 28. PMID: 33418361.
 23. Pacheco JP, Giacomini HT, Tam WW, Ribeiro TB, Arab C, Bezerra IM, Pinasco GC. Mental health problems among medical students in Brazil: A systematic review and meta-analysis. *Braz J Psychiatry.* 2017 Oct-Dec;39(4):369-378.
DOI: 10.1590/1516-4446-2017-2223. Epub 2017 Aug 31. PMID: 28876408; PMCID: PMC7111407.
 24. Almojali AI, Almalki SA, Allothman AS, Masuadi EM, Alaqeel MK. The prevalence and association of stress with sleep quality

- among medical students. *J Epidemiol Glob Health*. 2017 Sep;7(3):169-174.
DOI: 10.1016/j.jegh.2017.04.005.
Epub 2017 May 5.
25. Thomas M, Bigatti S. Perfectionism, impostor phenomenon, and mental health in medicine: A literature review. *Int J Med Educ*. 2020 Sep 28;11:201-213.
DOI: 10.5116/ijme.5f54.c8f8.
26. Baglioni C, Nanovska S, Regen W, et al. Sleep and mental disorders: A meta-analysis of polysomnographic research. *Psychol Bull*. 2016;142(9):969-990.
DOI: 10.1037/bul0000053.
27. Zhang L, Zheng H, Yi M, Zhang Y, Cai G, Li C, Zhao L. Prediction of sleep quality among university students after analyzing lifestyles, sports habits, and mental health. *Front Psychiatry*. 2022 Aug 4;13:927619.
DOI: 10.3389/fpsy.2022.927619.
PMID: 35990068; PMCID: PMC9385968.
28. Al-Khani AM, et al. A cross-sectional survey on sleep quality, mental health, and academic performance among medical students in Saudi Arabia. *BMC Research Notes*. 2019;12(1):1-5.
29. Joo HJ, Kwon KA, Shin J, Park S, Jang SI. Association between sleep quality and depressive symptoms. *J Affect Disord*. 2022;3(10):258-265.
DOI:10.1016/j.jad.2022.05.004
30. Hertenstein E, Feige B, Gmeiner T, et al. Insomnia as a predictor of mental disorders: A systematic review and meta-analysis. *Sleep Med Rev*. 2019;4(3):96-105.
DOI:10.1016/j.smrv.2018.10.006
31. Tubbs AS, Gallagher R, Perlis ML, Hale L, Branas C, Barrett M, Gehrels JA, Alfonso-Miller P, Grandner MA. Relationship between insomnia and depression in a community sample depends on habitual sleep duration. *Sleep Biol Rhythms*. 2020 Apr;18(2):143-153.
DOI: 10.1007/s41105-020-00255-z. Epub 2020 Feb 6. PMID: 34305449; PMCID: PMC8296753.
32. Riemann D, Krone LB, Wulff K, Nissen C. Sleep, insomnia, and depression. *Neuropsychopharmacology*. 2020 Jan; 45(1):74-89.
DOI: 10.1038/s41386-019-0411-y. Epub 2019 May 9. PMID: 31071719; PMCID: PMC6879516.
33. Nascimento MIC, et al. Manual Diagnóstico e Estatístico de Transtornos Mentais: DSM-5. 5ª ed. Porto Alegre. Artmed; 2014. Accessed on: 10 Jan 2023. Available: <http://www.institutopebioetica.com.br/documentos/manual-diagnostico-e-estatistico-de-transtornos-mentais-dsm-5.pdf>.
34. Hertenstein E, Trinca E, Wunderlin M, et al. Cognitive behavioral therapy for insomnia in patients with mental disorders and comorbid insomnia: A systematic review and meta-analysis. *Sleep Med Rev*. 2022;62:101-597.
DOI:10.1016/j.smrv.2022.101597
35. Hur S, Oh B, Kim H, Kwon O. Associations of Diet Quality and Sleep Quality with Obesity. *Nutrients*. 2021;13(9):3181.
DOI: 10.3390/nu13093181
36. Mehta N, Shafi F, Bhat A. Unique Aspects of Sleep in Women. *Mo Med*. 2015;112(6): 430-434.
37. Tang J, Liao Y, Kelly BC, Xie L, Xiang YT, Qi C, Pan C, Hao W, Liu T, Zhang F, Chen X. Gender and Regional Differences in Sleep Quality and Insomnia: A General Population-based Study in Hunan Province of China. *Sci Rep*. 2017 Mar 6;7:43690.
DOI: 10.1038/srep43690.
PMID: 28262807; PMCID: PMC5337959.
38. Eleftheriou A, Rokou A, Arvaniti A, Nena E, Steiropoulos P. Sleep Quality and Mental Health of Medical Students in Greece During the COVID-19 Pandemic. *Front Public Health*. 2021 Nov 19;9: 775374.
DOI: 10.3389/fpubh.2021.775374.
PMID: 34869189; PMCID: PMC8639533.
39. Chang J, Yuan Y, Wang D. [Mental health status and its influencing factors among college students during the epidemic of COVID-19]. *Nan Fang Yi Ke Da Xue Xue Bao*. 2020 Feb 29;40(2):171-176. Chinese.
DOI: 10.12122/j.issn.1673-4254.2020.02.06. PMID: 32376528; PMCID: PMC7086131.
40. Fernandes ACA, Padilha DMM, de Moura ACMA, de Aquino CEF, Lima IBA, Mota-Rolim SA. COVID-19 pandemic decreased sleep quality of medical students. *Sleep Sci*. 2022 Oct-Dec;15(4): 436-440.
DOI: 10.5935/1984-0063.20220075.
PMID: 36419809; PMCID: PMC9670770.

41. Gil-Calderón J, Alonso-Molero J, Dierssen-Sotos T, Gómez-Acebo I, Llorca J. Burnout syndrome in Spanish medical students. *BMC Med Educ.* 2021 Apr 22;21(1):231.
DOI: 10.1186/s12909-021-02661-4.
42. Bhugra D, Molodynski A, Ventriglio A. Well-being and burnout in medical students. *Ind Psychiatry J.* 2021 Jul-Dec; 30(2):193-197.
DOI: 10.4103/ipj.ipj_224_21.
Epub 2021 Nov 23.
43. De Mélo SJML, Valença MM, Rocha-Filho PAS. Individual and residency program factors related to depression, anxiety and burnout in physician residents - a Brazilian survey. *BMC Psychiatry.* 2022 Apr 19; 22(1):272.
DOI: 10.1186/s12888-022-03916-0.

© 2023 Lopes et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96652>*