



Spécies Diversity and Physiological Characterization of *Fusarium* spp. Isolates Causing Potato (*Solanum tuberosum* L.) Fusariosis (Far-North, Cameroon)

Boydoul Frédéric Ulrich ^{a*}, Sobda Gonné ^b,
Fankou Dougouan Merline Yoyo ^b, Abdoul Madjerembe ^a,
Rapmo Kononer Serge ^c, Eloa Seusseмага ^a,
Metsena Pierre ^c, Ngoh Dooh Jules Patrice ^a
and Philippe Kosma ^c

^a Département of Biological Sciences, Faculty of Science, University of Maroua, P.O. Box 814 Maroua, Cameroon.

^b Agricultural Reaserch Institue for Developpment (IRAD) Maroua, Region BP: 33 Maroua, Cameroon.

^c Higher National Polytecnic School of Maroua, University of Maroua, P.O.Box 1450 Maroua, Cameroon.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPR/2023/v12i1215

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96113>

Original Research Article

Received: 21/10/2022

Accepted: 29/12/2022

Published: 27/02/2023

*Corresponding author: Email: boydoulfrederic@gmail.com;

ABSTRACT

Introduction: Potato production in the Far North Region, Cameroon is hampered by a wide range of fungi of the genus *Fusarium*.

Aims: The aim of this work was to isolate and conduct physiological characterization of *Fusarium* spp. isolates responsible for potato *Fusarium* disease in the Far North Region, Cameroon.

Methodology: Isolates were obtained from diseased organs (leaves) from fifteen villages in the district of Mogodé, Mokolo and Koza, Far North Cameroon. The species responsible for fusariosis were identified using several identification Keys on a base of symptoms and morphological spores characteristics (macroconidia, microconidia and chlamyospores). *Fusarium* incidence and rainfall were assessed. Radial growth, sporulation and pathogenicity of the isolates were use to evaluated the parameters of physiological characterization.

Results: The overall average incidence of *Fusarium* head blight in the Far North Region, Cameroon is 29.36%. Furthermore, out of the 20 isolates obtained, six *Fusarium* species (*Fusarium oxysporium*, *F. solani*, *F. equiseti*, *F. avenacearum*, *F. colmorum* and *F. sambicum*) were identified. The highest growth rate (8.1 cm), the highest spore production (8×10^5) and the highest severity index (3%) were respectively obtained with the isolates FUROM 2 and FUTEK 3, FUMOG 1, FUMOG 2, FUROM 1 and FURAF from Mogodé District on PCA medium.

Conclusion: Potato production in the Far North Region, Cameroon is confronted by various *Fusarium* species which have physiological characteristics that vary according to the area of origin of the isolates and the growing media.

Keywords: *Solanum tuberosum* L.; *Fusarium*; incidence; isolates; sporulation; pathogenicity.

1. INTRODUCTION

Potato (*Solanum tuberosum* L.) is a tuberous herbaceous plant native to Peru and belonging to the Solanaceae family [1]. It is a vegetatively propagated species that produces more nutritious food on less cultivated area and in the harshest of climates than any other crop [2,3]. It has become one of the staple foods of humanity and thus occupies a prominent place in the diet of many countries in terms of the land it occupies, the jobs it provides and the production volumes it generates. World production was estimated at more than 376 million tonnes on 19.4 million hectares spread over 151 producing countries in 2014 [4]. Which makes the potato the fourth most important non-cereal food crop in the world after wheat, rice and maize [4].

In Cameroon, it is grown in high altitude areas (1000 to 3000 m) [5,6] and extensively in six of the ten regions of Cameroon (North-West, South-West, West, Adamaoua, Littoral and Far-North), mostly by rural people and mainly by women [7]. It is a staple food for the populations of these regions (annual consumption of 4 to 10 kg per capita/year) [5,6] and an important source of income for producers. Yields are generally very low and range from 3 to 11 tonnes per hectares in the Far North Region of Cameroon, while those in European countries average 25 tonnes per hectares and reach 60 t/ha [7,8,9]. The low

yields observed in this production zone are associated with poor farming practices such as the absence of crop rotation, the semi-open system, the poor choice of varietal material and chemical fertilisation methods [7,10,6] pests and especially diseases that hamper potato development in all potato production areas in the world [11,8,6].

In almost all potato production areas, *Fusarium* head blight appears to be the most important foliar and post-harvest disease causing the highest yield losses estimated at more than 25% per year [12,13,14]. Various studies have also shown that the severity of diseases caused by *Fusarium* varies from 6 to 25% and even 60% in some cases [12,14]. In Great Britain, the national average incidence of affected tubers could be as high as 1.4% and 50-100% of tubers are attacked [13]. More than 50% of tubers are infected with *Fusarium* in Michigan [15]. Other studies have shown that the severity of diseases caused by *Fusarium* phytopathogens are among the most common fungi found on potato crops in the Far North Region, Cameroon [6].

It is estimated that there are thirteen species of *Fusarium* causing potato fusariosis depending on agro-ecological zones [16] edaphic factors, environmental factors [17,18], climatic factors [18,19] and cultivars. Nevertheless, *Fusarium sambicum* is the most widespread fungus in

North America, China and parts of Europe [20-23]. *Fusarium coeruleum* is the most widespread agent in the United Kingdom [22]. In addition, *F. oxysporium*, *F. avenaceum*, *F. acuminatum*, *F. equiseti*, *F. sulphureum* and *F. solani* have been recorded on potato [17,24]. In Africa, *Fusarium sambicum*, *F. oxysporium*, *F. verticillioides* and *F. incarnatum* predominate in Egypt [14]. To the best of our knowledge, no study has revealed the specific diversity of fungi of the genus *Fusarium* in Cameroon in general and in the Far North Region in particular. However, the knowledge of the diversity and aggressivity of these species constitute the diagnostic gateways to master the biology, the epidemiological mechanisms and especially the setting up of biological control methods. The present review provides an update on the status of the genus *Fusarium* of potato in the Far North Region, Cameroon, and opens up avenues of investigation. The aim of this work is to carry out a physiological characterization of *Fusarium* isolates and to evaluate the degree of aggressiveness of the species.

2. MATERIALS AND METHODS

2.1 Area of Study

The study was carried out in two Departments : the Department of Mayo-Tsanaga with geographical coordinates 10°24'23" N and 13°49'17" E, and the Department of Diamaré with coordinates between 10°44'33" N and 14°16'81" E [25], all located in the Far North Region, Cameroon.

2.2 Plant Materials

The plant material used for this study consisted of two potato varieties, namely the resistant local variety called Dosa and the improved susceptible variety called Spunta.

2.3 Sample Collection, Incidence Assessment and Rainfall

Samples potato leaves were collected over a period of two years (June 2020 to October 2022) in fifteen villages spread over three district namely Mogodé, Mokolo and Koza all located in the Mayo-Tsanaga Department, Far North Region of Cameroon. The samples collected were coded and then sent to the IRAD plant pathology laboratory in Maroua for isolation and specific tests. The incidence was evaluated by counting the number of diseased plants out of the total number of plants present in the fields

surveyed per village and per District. Rainfall data were collected from the stations of the Agricultural Delegations of each District surveyed [26]. Incidence of diseases were evaluated using the following formula [26]:

$$I(\%) = \frac{np}{N} \times 100.$$

Where, I is incidence; np number of plants showing symptoms per plot; and N total number of plant in plot.

2.4 Culture Media and Obtention of Isolates

Obtaining isolates consisted firstly of preparing the culture media, then culturing the samples, purifying the strains and finally isolating them.

The culture of the samples consisted first of the preparation of the culture media PDA (200g potato, 20g D-Glucose and 15g Agar), PPA (70 grams of Petit Pois and 15g agar), PSA (200g potato, 20g sucrose and 20g Agar) and PCA (40g potato, 40g carrot and 20g Agar) all dissolved in 1L distilled water as defined by [27,28,29].

The disinfected organs were dried on a white hygienic cloth for 30 minutes to remove or absorb the water on the samples. Finally, the parts of the samples showing symptoms of the disease were cut into squares using a bustori blade. The samples were taken with the help of a coin lifter and aseptically and individually placed in petri dishes containing culture media previously prepared in the fume hood. The petri dishes were closed, sealed with parafilm and incubated in a climate chamber at 22-27 °C under a 12/12h photoperiod. The mycelium developed from the explant and after 2-4 days reached sufficient growth to proceed to purification [27].

The purification of the strains consisted of three successive subcultures on the PDA medium, by taking fragments of mycelial discs of 08 mm in diameter at the growth front of the 10-day-old cultures to be sown in the centre of each petri dish containing the medium. Isolates were obtained by taking a mycelial disc at the growth front from each pure culture obtained after three successive purifications. The isolates thus obtained were directly used for species identification and partially preserved in sterile distilled water at 4°C for further use [27].

2.5 Species Identification

The identification of the species was based on the identification keys as defined by [30,12,14,6] this was done by comparing the symptoms on different organs (leaves and tubers) in the field and the microscopic structures in the laboratory. The identification of the different species in the laboratory was done by determining the presence or absence of filamentous structures such as macroconidia, microconidia and chlamyospores characteristic of each *Fusarium* species on 20-day-old culture media.

2.6 Physiological Characterization of Isolates

Four culture media, namely PDA (Potato Dextrose Agar), PCA (Potato Carrot Agar), PSA (Potato Sucrose Agar), and PPA (Pea Agar) were used to evaluate the parameters of physiological characterisation, including cultural characteristics, radial growth, sporulation and pathogenicity.

2.6.1 Evaluation of the cultural characteristics of the isolates

The parameters of morphological characterisation. the colour, shape, appearance and relief of the colonies grown on PPA, PCA, PSA and PDA culture media were evaluated 7 days after subculturing, by macroscopic observation in order to determine the specific cultural characteristics of each isolate according to the culture media and the origin of the strains. The whole set was incubated at a photoperiod of 12H/12 and at the laboratory's ambient temperature, which varied between 25° and 30°C.

2.6.2 Measurement of radial growth of isolates

Radial growth was assessed by measuring the perpendicular diameters previously traced on the back of the petri dishes and inside which a fragment of mycelium disc of 08 mm diameter of each isolate was placed. Measurements were made every two days and at the same time by simple measurement with a 30 cm ruler. The measurement was done after 2, 4, 6 and 6 days (day that one of the colonies had filled the petri dish) of culture to evaluate the growth rate of the different strains under the influence of the different culture media. Three replicates were made for each isolate and culture medium. The

Petri dishes containing the culture media were incubated at a photoperiod of 12/12h and at room temperature of the laboratory, ranging between 25° and 30°. The formula of [31] opposite was used to calculate the radial growth of each isolate.

$$D = \frac{d1 + d2}{2} - d_0$$

D = diameter of growth

d₀ = diameter of mycelial disc

d1 and d2 = two perpendicular diameters

2.6.3 Sporulation measurement of isolates

Spores production was evaluated on the 20-day-old cultures, incubated at a 12H/12 photoperiod and at room temperature. For this purpose, four 08 mm mycelial discs of each isolate were collected and placed in tubes containing 1 mL of sterile distilled water. The solutions were filtered through muslin cloths to remove unwanted fragments. Next, 0.1mL of solution from each isolate was withdrawn with an insulin syringe and then calibrated with a malassez cell or hematimeter for spore enumeration. Each operation was repeated three times for each isolate [32].

2.6.4 Pathogenicity test of isolates

The detached leaf disc test was used to assess the pathogenicity of the different isolates. For this purpose, young healthy leaves of two varieties, namely Dosa, a resistant local variety, and Spunta, a susceptible introduced variety, one month old, were collected very early in the morning in the field at about 6 a.m. and individually placed in white plastic bags and transported to the laboratory for conditioning and testing.

2.6.4.1 Conditioning of detached leaves and incubation

Conditioning consisted of carefully washing the leaves in running tap water, rinsing them with 70° alcohol for one minute, and finally drying them on white absorbent paper for 30 minutes with their petioles inserted in the cotton wool to maintain moisture [33]. Then white toilet paper of 110 mm length and width dimension was cut with the scissors and placed in sterile petri dishes of 150 mm diameter. Four millilitres (4ml) of sterile distilled water were sprayed into each petri dish using a sterile propette and the leaves were

transferred into each dish. Finally, spore suspensions obtained from the 20-day-old pure cultures were calibrated at $3-4 \times 10^5$ spores/ml using the Malassez cell. The suspensions were spread on the four cardinal points of the upper surface of the leaf using a micropipette. Three replicates were made for each isolate and according to the potato varieties. The media were regularly humidified every two days to maintain the relative humidity of the medium. Incubation was done at a 12H/12 photoperiod and at room temperature [33].

2.7 Data Analysis

The data obtained were subjected to an analysis of variance using SPSS 25.0 software. The DUNCAN test was used to compare the means of the measured parameters at the 5% threshold.

3. RESULTS

3.1 Evolution of the Incidence of Fusariosis in the Study Area

The incidence of *Fusarium* wilt varied considerably between villages and study Districts. The average incidence of fusariosis in the Far North Region, Cameroon is 29.36%. However, it is in the district of Mogodé that the highest average incidence was obtained, 51.08%, contrary to the district of Koza where the lowest incidence was obtained, 6% (Table 1). The intermediate average incidence was

obtained in the Borough of Mokolo, 19.5% (Table 1).

3.2 Rainfall Trends in the Study Area

Rainfall varies over time and space depending on the weeks, months and Study Districts. The highest average rainfall, the average or intermediate rainfall, and the lowest average rainfall were respectively obtained in the district of Mogodé, Koza, and Mokolo: 1,988 mm, 1,024 mm and 828.5 mm (Fig. 1).

3.3 Physiological Characteristics of the Isolates

3.3.1 Isolates obtained

Twenty isolates were obtained in the district of Mogodé, Mokolo and Koza. Of the Twenty isolates, fifteen isolates, 75% (FUMOG 1, FUMOG 2, FUROM 1, FUROM 2, FUROM 3, FUNDG 5, FUTEK 3, FUZIM 1, FUSRA 1, FUSRA 2, FURAF 12, FUKAR 5, FUSIK 8, FUGOU 1 and FUMOV) were obtained respectively in the villages (Mogodé, Rhoumzou, Ndegvaya, Teki, Zimi, Sirakoti, Raffa, Karanti, Rhumsiki, Gouria and Mouvou) of Mogodé District; three isolates, 15% (FUKOS 1, FUKOS 2 and FUGAW 4) were obtained in two villages (GAWAR and KOSEHONNE) in the of Mokolo and two isolates, 10% (FUKOZ 1 and FUKOZ 2) were obtained in the village ZILER in the district of Koza.

Table 1. Evolution of the incidence of fusariosis according to the villages and districts surveyed in two years

Areas	Means Incidence 2020-2021			Means Incidence 2021-2022			Means 2020-2022
	Villages	Incidence%	Means	Villages	Incidence%	Means	
Mogode	Mogodé	45%	39,58%	Mogodé	75%	62,58%	51,08%
	Rhomuzou	35%		Rhoumzou	45%		
	Ndegvaya	67%		Ndegvaya	57%		
	Teki	25%		Teki	28%		
	Zimi	51%		Zimi	53%		
	Sirakoti	20%		Sirakoti	40%		
	Raffa	49%		Raffa	69%		
	Karanti	32%		Karanti	42%		
	Rhumsiki	23%		Rhumsiki	33%		
	Gouria	15%		Gouria	35%		
	Mouvou	32%		Mouvou	62%		
Mokolo	Migi	34%	Migi	24%	24,5%	19,5%	
	Kosehone	20%	Kosehone	25%			
Koza	Gawar	15%	Gawar	24%	7%	6%	
	Ziler	5%	Ziler	7%			
Overall average incidence in the study area: 29.36%							




Fig. 1. Changes in rainfall in the districts of Mogodé, Mokolo and Koza

3.3.2 Species identified

In general, 6 species were identified from the samples collected in the three study Districts (Mogodé, Mokolo and Koza): *Fusarium oxysporium*, *Fusarium solani*, *Fusarium equiseti*, *Fusarium avenacearum*, *Fusarium colmorum* and *Fusarium sambicum*. Clamydospores, macroconidia and microconidia characteristic of the genus *Fusarium* were also obtained (Fig. 2). All the 6 species inventoried are present in Mogodé District unlike Mokolo and Koza Districts where only *Fusarium oxysporium* and *Fusarium solani* were specifically identified.

3.3.3 Cultural characteristics of the isolates obtained

The results of the morphological characterization of the colonies grown on PPA, PCA, PSA and PDA media revealed considerable variation according to the above media and the areas of origin of the isolates (Fig 3). In general, the isolates obtained in the three study Districts all show a white colour as a common morphological feature, as opposed to the bulging relief feature not observed in Koza District with isolates (FUKOZ 1 and FUKOZ 2).

3.3.4 Radial growth diameters of isolates

Four statistical groups were obtained from the evaluation of the growth diameter at 6 days after transplantation. Statistical analyses ($P = 0.000$) showed that growth diameters varied considerably between isolates and culture media.

Nevertheless, the highest growth diameters (8.1 cm) were obtained mainly in Mogodé District with isolates FUROM 2, FUTEK 3 and FUKOS1, on PCA medium, in contrast to isolates from Mokolo and Koza Districts (Table 2).

3.3.5 Sporulation rate of isolates

Statistical analysis revealed a highly significant difference ($P=0.000$) in spore production. Isolates from Mogodé were the most abundant in spore production compared to isolates from the other district. (Table 2). Spores were most abundant in PCA and PSA media respectively with isolates FUMOG 1 and FUMOG 2 at 8×10^5 in Mogodé District as opposed to isolate FUKOZ 1 from Koza District at 1×10^5 spores (Table 2).

3.3.6 Pathogenicity of isolates

Statistical analysis of the severity index carried out by artificial inoculation of potato leaves of the Dosa and Spunta varieties from *Fusarium* spore suspensions revealed a significant difference ($P=0.000$) in the degree of aggressivity of the different isolates (Fig. 4). The severity indices of the isolates in Mogodé District ranged from 0.92% to 2% for the Dosa variety and from 1% to 3% for the Spunta variety. However, in Mokolo district, the index varied from 1.5% to 2% for Dosa and from 1% to 2.5% for Spunta. No variation in the severity index was observed across the two potato varieties in Koza. However, the index was 1.5% for the Dosa variety and 2% for the Spunta variety. Overall, the isolates with the highest severity index were

obtained in Mogodé district with isolates FUROM 1 and FURAF 12, 3%, 2.5%, in contrast to Koza district, 1.5% obtained with isolate FUGAW 4 (Fig. 5). The highest severity index (3%) was obtained with isolate FUROM 1 on the variety Spunta, which confers the status of susceptible variety on Spunta and resistant variety on Dosa.

4. DISCUSSION

The results of the survey work showed that the overall average incidence of *Fusarium* spp. is 29.36%. This percentage shows that *Fusarium* head blight is a cosmopolitan fungal disease that significantly affects the potato crop in the Far North Region, Cameroon. These results are in agreement with the work of [13,14] who reported that in Great Britain, the national average incidence of affected tubers was as high as 1.4% and 50-100%. Furthermore, [15] also showed that more than 50% of tubers are infected by *Fusarium* in Michigan; it is 43.7% in the Oued Tafna region of Algeria [34]. The random distribution of *Fusarium* incidence in Mogodé (51.08%), Mokolo (19.5%) and Koza (6%) correspond to the rainiest months respectively obtained during September in Mogodé and August in Mokolo and Koza: this shows that the

intensity of the disease varies according to the rainfall prevailing in a given geographical area. These findings are in agreement with those of [17,18,35] who showed that *Fusarium* head blight is a disease that adapts rapidly to climatic conditions to develop on the host plant.

Regarding the isolation itself, it should be noted that twenty (20) isolates were obtained from samples collected from potato fields in the Far North Region, Cameroon. This result is in line with those reported by [36]. Indeed, the latter was able to isolate 120 isolates of *Fusarium* responsible for yield losses on cowpea in the Far North Region, Cameroon. Various of *Fusarium* (*Fusarium oxysporium*, *Fusarium solani*, *Fusarium equiseti*, *Fusarium avenaceum*, *Fusarium colmorum* and *Fusarium sambicum*) were identified as being responsible for potato fusariosis in the Far North Region. These results are in agreement with those identified in North America, China and parts of Europe by [22,23]. These authors reported the presence of *Fusarium sambicum* as the most prevalent fungus in these regions. In addition, *F. oxysporium*, *F. avenaceum*, *F. equiseti*, and *F. solani* have been recorded on potato by [17, 24]. In Africa, *Fusarium sambicum*, *F. oxysporium*, predominate in Egypt [14].

Table 2. Growth diameter and spore production of isolates

Areas	Isolates	Number of spores×10 ⁵ ml				Isolats	Growth diameter (cm)			
		Culture media					Culture media			
		PCA	PSA	PPA	PDA		PCA	PSA	PPA	PDA
Mogode	FUMOG1	8	7	5	2	FUMOG1	8	8	7,5	4,2
	FUMOG2	8	8	5	3	FUMOG2	8	8	8,1	6,9
	FUROM1	5	6	4	2	FUROM1	6,2	5,9	7	5,3
	FUROM2	5	6	5	4	FUROM2	8,1	7,6	8	6,1
	FUROM3	4	7	4	3	FUROM3	8	8	8	8
	FUNDG5	3	6	3	3	FUNDG5	6,6	6	7,9	6,5
	FUTEK3	6	4	4	4	FUTEK3	8,1	8	8	8
	FUZIM1	5	4	5	2	FUZIM1	8	8	8	8
	FUSRA1	4	4	4	2	FUSRA1	7,6	7,4	7,8	6
	FUSRA2	5	6	4	3	FUSRA2	6,2	5,5	7,5	6
	FUKAR5	7	6	8	3	FUKAR5	6,8	7,9	8	6,5
	FURAF12	5	5	5	2	FURAF12	7,5	7,8	7,1	5,1
	FUSIK8	5	4	5	2	FUSIK8	7,3	8	7	6
	FUGOU1	6	6	5	4	FUGOU1	6,5	5,4	6,8	6
	FUMOV1	4	6	3	2	FUMOV1	7,6	4,2	7	5,5
Mokolo	FUKOS1	6	5	4	2	FUKOS1	8,1	7,9	8	7
	FUKOS2	6	4	4	3	FUKOS2	6,9	6,6	8	7
	FUGAW4	5	4	5	2	FUGAW4	7,7	5,8	7,7	6
Koza	FUKOZ1	4	5	2	1	FUKOZ1	7,8	7,6	7,8	6,4
	FUKOZ2	4	5	3	3	FUKOZ2	7,7	7,7	7,5	6,7
Probability P=0,000						Probability P=0,001				




Fig. 2. *Fusarium* species diversity in the are study




Fig. 3. Cultural characteristics of some *Fusarium* isolates on culture media




Fig. 4. Leaf appearance 10 days after inoculation




Fig. 5. Severity index of isolates according to varieties.

The results of the morphological characterization of the colonies of the isolates grown on PPA, PCA, PSA and PDA culture media revealed that the majority of the colonies had a white colour, a regular growth, a cottony aspect on the different culture media with a relief varying from Flat to Domed. The results obtained corroborate with those of [36]. Who showed that the white colour of the colonies is widely observable on most of the PDA culture media. The large variability observed in the growth rate and spore production

of the isolates is due to the fact that the isolates tested came from three different districts, all of which were subject to the influence of climatic, soil and environmental factors prevailing in each district. Furthermore, the nature of the isolates, the nutritional elements contained in the culture media, the temperature of the reaction medium, and the humidity of the air that accompanied the experiment would be at the origin. This same finding was reported by [34,37,36]. For them, the growth rate varies according to the culture

media. However, [38] reported that PDA medium is not conducive to the production of *Fusarium* spores isolated from cowpea in the Far North Region, Cameroon in contrast to potato isolates of *Fusarium* that produced spores in low quantities. The variation in the degree of aggressiveness of the isolates is thought to be due to the fact that the isolates do not all come from the same area and that the two potato varieties used do not have the same genetic make-up or ability to respond to phytopathogenic stresses. This was also reported by [20] who showed that the aggressiveness of the isolates varies between potato cultivars.

5. CONCLUSION

It was found that six different species of *Fusarium* (*Fusarium oxysporium*, *F. solani*, *F. equiseti*, *F. avenacearum*, *F. colmorum* and *F. sambicum*) cause potato fusariosis in the Far North Region, Cameroon but also cause an estimated 29.39% damage. Furthermore, isolates from Mogodé District have a higher radial growth, sporulation rate and degree of aggressiveness than the preventive isolates from Mokolo and Koza Districts. The Dosa variety is the one that is resistant to *Fusarium* fungi, while the Spunta variety is not.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

RÉFÉRENCES

1. Spooner DM, McLean K, Ramsay GA. Single domestication for potato based on multilocus amplified fragment length polymorphism genotyping. Proceedings of the National Academy of Sciences of the United States of America. 2005;102:14694-14699.
2. Titsop V. Adaptability and effect of fertilizers on the production of 5 varieties of potato: case of the departments of Diamare and Mayo-Tsanaga- Far North Cameroon, end of study dissertation with a view to obtaining the design engineering degree in agronomy, plant production option, Department of Agriculture Livestock and Derived Products, University of Maroua. 2018:42-52.
3. Jean-Pierre W. the potato saga, Paris, Edition Cercle d'Art. 2008:160.
4. FAO. Food and Agriculture Organization of the United Nations. Rome, Italy, Production Year Book; 2019.
5. IRAD. (Institute of Agricultural Research for Development), Sustainable improvement of the productivity and competitiveness of the plantain sector in Cameroon through the use of innovative technologies. 2012:4-8.
6. Ngoh Dooh JP, Boydoul FU, Abdoul Tchoupou TDB, Bouba D, Hawaou A, Philippe Kosma, Ambang Z. Inventory of the potato diseases and impact on growth and yield traits in far North Cameroon. International Journal of Biological and Chemical Sciences. 2020a;14(8):2826-2836.
7. Fonten DA, Demo P, Njuaem DK. Status of potato production, marketing and utilization in Cameroon. Paper presented at the 9th Triennial Symposium of the International Society of Tropical Root Crops - Africa Branch; 2005.
8. Diop P, Sylla ES, Diatte M, Labou B, Diarra K. Effect of cut seed tubers and pregermination on potato tuber yield. International Journal of Biological and Chemical Sciences. 2019;(7):31573163. Available:https://dx.doi.org/10.4314/ijbcs.v13i5
9. Ngoh Dooh JP, Nsimi Mva A, Boydoul FU, Philippe Kosma, Ambang Z. X influence of fertilizers on incidence and severity of viral and bacterial potato (*Solanum tuberosum*) diseases under field condition. International Journal of Environment, Agriculture and Biotechnology. 2020b:8.
10. Ngoyi AN, Masanga GK, Bila HM, Yashima AY, Milambo MM, Ndjibu LN, Baboy LL. Effect of organic amendments on the growth and yield of potato (*Solanum tuberosum*) grown on degraded soil in the Kabinda region, Democratic Republic of Congo. International Journal of Biological and Chemical Sciences. 2020;14(5):1812-1819. Available:https://doi.org/10.4314/ijbcs.v14i5.24
11. Son D, Somda I, Legreve A, Schiffers B. Effect of plant diversification on pest abundance and tomato yields in two cropping systems in Burkina Faso: Farmer practices and integrated pest management. International Journal of Biological and Chemical Sciences. 2018; 12(1):101-119.

- Available:<http://doi.org/10.4314/acsj.v9i1.27638>
12. Mecteau MR, Arul J, Tweddell RJ. Effect of different salts on the development of *Fusarium solani* var *coeruleum*, a causal agent of potato dry rot. *Phytoprotection*. 2008;89(1):1-6. Available: <https://doi.org/10.7202/000377ar>
 13. Estrada JR, Gudmestad NC, Rivera VV, Secor GA. *Fusarium graminearum* as a dry rot pathogen of potato in the USA: Prevalence, comparison of host isolate aggressiveness and factors affecting etiology. *Plant Pathology*. 2010;59:1114-1120.
 14. Yikilmazsoy, Tosun. Characterization of *Fusarium sambucinum* isolates associated with potato dry rot and evaluation of cultivar susceptibility and fungicides. *Turkish Journal of Agriculture and Forestry*. 2021;45:222-233. DOI: 10.3906/tar-2006-100
 15. Kirk W, Wharton P. *Fusarium* dry rot posing problems in potatoes. In *Vegetable Crop Advisory Team Alert Michigan Potato Diseases*, Michigan State University; 2008. Available: <http://tinyurl.com/3jzmx4l>.
 16. Cullen DW, Toth IK, Pitkin Y, Boonham N, Walsh K. Use of quantitative molecular diagnostic assays to investigate *Fusarium* dry rot in potato stocks and soil. *Phytopathology*. 2005;95:1462-1471.
 17. Xu XM, Nicholson P, Thomsett MA, Simpson D, Cooke BM, Doohan FM, Brennan J, Monaghan S, Moretti A, Mule G, Hornok L, Beki E, Tatnell J, Ritieni A, Edwards SG. Relationship between the fungal complex causing *Fusarium* head blight of wheat and environmental conditions. *Phytopathology*. 2008b;100:763-773. DOI: 10.1094/PHYTO-98-1-0069.
 18. Ferrigo D, Raiola A, Causin R. *Fusarium* toxins in cereals: Occurrence, legislation, factors promoting the appearance and their management. *Molecules*. 2016;21(5):627. Available:<https://doi.org/10.3390/molecules21050627>
 19. Rossi V, Ravanetti A, Patteri E, Giosuè S. Influence of temperature and humidity on the infection of wheat spikes by some fungi causing *Fusarium* headblight. *Journal of Plant Pathology*. 2001;83(3):189-198. Available:<http://dx.doi.org/10.4454/jpp.v83i3.1128>
 20. Doohan FM, Brennan J, Cooke BM. Influence of climatic factors on *Fusarium* species pathogenic to cereals. *European Journal of Plant Pathology*. 2003;109(7):755-768. DOI: 10.1023/a:1026090626994
 21. Bojanowski A, Avis TJ, Pelletier S, Tweddell RJ. Management of potato dry rot. *Postharvest Biology and Technology*. 2013;84:99-109.
 22. Heltoft P, Brurberg MB, Skogen M, Le VH, Razzaghian J, Hermansen A. *Fusarium* spp. causing dry rot on potatoes in Norway and development of a real-time PCR method for detection of *Fusarium coeruleum*. *Potato Research*. 2016;59(1):67-80.
 23. Patil VU, Sagar VGV, Chakrabarti SK. Draft genome sequence of potato dry rot pathogen *Fusarium sambucinum* Fckl. F-4. *American Journal of Potato Research*. 2017;94(3):266-269.
 24. Yin Y, Li YC, Bi Y, Chen SJ, Li YC, Yuan L, Wang Y, Wang D. Postharvest treatment with β -aminobutyric acid induces resistance against dry rot caused by *Fusarium sulphureum* potato tuber. *Agricultural Science in China*. 2010;9:1372-1380.
 25. Raunet M. Quelques clés morphologiques de sol du Nord Cameroun à l'usage des agronomes. 2017:18.
 26. MINADER. Ministère de l'Agriculture et du Développement Rural. Données pluviométriques 2020-2022: Cas des Délégations d'Agriculture des Arrondissements de Mogodé, Mokolo et Koza; 2022.
 27. Ngoh Dooh JP, Ambang Z, TihEwola A, Heu A, Kosma P, MahoYalen EJ, GhogomuTih R. Screening and the effect of extracts of *Thevetia peruviana* on the development of *Colletotrichum gloeosporioides*, causal agent of cassava anthracnose disease. *E3 Journal of Agricultural Research and Development*. 2014a;4(4):054-065.
 28. Samson RA, Hoekstra ES, Frisvad JC, Filtenborg O. Introduction to food and air borne fungi. 6th.ed. Utrecht Central Bureau Voor Schimmel Cultures. 2022:379.
 29. Simmons EG. *Alternaria* and identification manuel. CBS Biodiversity Series N° 6 CBS. Fungal Biodiversity Centre Utrecht, The Netherlands. 2007:775.

30. Lepoivre P. Phytopathologie. De loece, les presses agronomiques de Gembloux. 2003;413.
31. Singh SR, Jackai LEN, Thottappilly G, Cardewell KF, et Myers GO. Status of research on constraints to cowpea production. In: Biotechnology: Enchancing Research on Tropical Crops in Africa, éd., IITA. 1999:21-26.
32. Nyassé S, Cilas C, Hérail C, Blaha G, Leaf inoculation as an early screening test of cocoa (*Theobroma cacao* L.) resistance to Phytophthora black pod disease: Crop Protection. 1995;(14):657-663.
33. Ganie SA, Ghany MY, Nissar Q, Jabeen N, Anjum Q, Ahangar Ayaz AF. Status and symptomatology of early blight (*Alternaria solani*) of potato (*Solanum tuberosum*) in Kashmir valley. African Journal of Agricultural Rechearch. 2013;8(41):5104-5115.
34. Bessadat M. Isolement, identification et caractérisation des *Alternaria* sp. Responsables de la détérioration des plantes maraichères par des systèmes enzymatiques et moléculaire. Mémoire Doctorat. Université d'Oran; 2014.
35. Gordon JL, Lefeuvre P, Escalon A, Barbe V, Cruveiller S, Gagnevin L, Pruvost O. Comparative genomics of 43 strains of *Xanthomonas citri* pv. *citri* reveals the evolutionary events giving rise to patotyoes with different host ranges. BMC Genomics. 2015:16.
36. Mechta N. Azouaoui-at KT, Rahmatou F. *Fusarium oxysporum* f. sp. *Albedinis*: Effets du milieu de culture sur la croissance mycelienne, la sporulation et la production de l'acide fusarique. Algerian Journal of Arid Environnement. 2015:9.
37. Ferrocino I, Chitarra W, Pugliese M, Gilardi G, Gullino ML, Garibaldi A. Effect of elevated atmospheric CO₂ and temperature on disease severity of *Fusarium oxysporum* f.sp. *lactucae* on lettuce plants. Applied Soil Ecology. 2013; 72:1-6. Available:<http://dx.doi.org/10.1016/j.apsoil.2013.05.015>
38. Metsena P, Sobda G, Kosma P, Fankou Dougouan MY. Identification of *Fusarium oxysporum* sf *tracheiphilum* strains responsible of cowpea wilt in Far-north region of Cameroon. Journal of Applied Biosciences. 2021;164. Available:www.m.elewa.org/journals/ Available:<https://doi.org/10.35759/JABs.164.7>

© 2023 Ulrich et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96113>