

Changes in Intraocular Pressure and Anterior Chamber Depth after Phacoemulsification in Non-Glaucomatous Patients

Mona Abdallah^{1*}, Omar M. Ali¹, Abd El-Nasser A. Mohammad¹ and Wael Soliman¹

¹*Department of Ophthalmology, Assiut University Hospital, Assiut, 71516, Egypt.*

Authors' contributions

This work was carried out in collaboration among all authors. Author OMA designed the study. Author MA collected the data, wrote the protocol and managed the literature searches. Authors WS and AENAM performed the statistical analysis and wrote the first draft of the manuscript. All authors read, revise and approve the final manuscript.

Article Information

DOI: 10.9734/OR/2019/v10i130098

Editor(s):

(1) Dr. Tatsuya Mimura, Associate Professor, Department of Ophthalmology, Tokyo Women's Medical University Medical Center East, Nishiogu, Arakawa-ku, Tokyo, Japan.

Reviewers:

- (1) Gabor Nemeth Borsod-Abaúj-Zemplén, University Teaching Hospital, Miskolc, Hungary.
(2) Umezurike Benedict Chidozie, Government House Clinic, Umuahia, Nigeria.
(3) Adnan Aksoy, Sütçüimam University Faculty of Medicine, Turkey.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/47792>

Received 17 December 2018

Accepted 05 March 2019

Published 18 March 2019

Original Research Article

ABSTRACT

Aims: To evaluate changes in intraocular pressure (IOP) and anterior chamber depth (ACD) after uneventful phacoemulsification with intraocular lens (IOL) implantation.

Study Design: Prospective interventional comparative study.

Place and Duration of Study: Department of Ophthalmology, Assiut University Hospital, Assiut, Egypt between September 2016 and October 2017.

Methodology: 100 Patients with visually significant cataract (39 males (39%) and 61 females (61%). All patients underwent uneventful phacoemulsification. Intraocular pressure and ACD were measured preoperatively by using Goldmann applanation tonometer and Ultrasonography respectively, and at 1 week, 1 month, and 3 months postoperatively.

Results: The postoperative ACD was higher than the preoperative value by 0.26 ± 0.06 mm at 1 week, 0.45 ± 0.06 mm at 1 month, and 0.59 ± 0.08 mm at 3 months postoperatively ($P = .000$). $P < 0.05$ was considered to indicate a statistically significant difference. The IOP at 1 week, 1 month,

*Corresponding author: E-mail: monaabdallah51290@aun.edu.eg;

and 3 months postoperatively was lower than the preoperative value by 1.03 ± 0.3 mmHg, 1.52 ± 0.46 mmHg, and 2.14 ± 0.62 mmHg, respectively ($P=0.001$).

Conclusion: This study revealed that there was a decrease in IOP and increase in ACD after cataract extraction by phacoemulsification. Patients with ocular hypertension, open-angle glaucoma, and narrow-angle glaucoma may benefit from cataract extraction.

Keywords: Cataract; phacoemulsification; intraocular pressure; anterior chamber depth.

ABBREVIATIONS

IOP : Intraocular Pressure

ACD : Anterior Chamber Depth

AS-OCT : Anterior Segment Optical Coherence Tomography

1. INTRODUCTION

Cataract surgery is one of the most common surgeries performed in the world. In addition to its significant impact on improvement of visual acuity, phacoemulsification also decreases the intraocular pressure (IOP) [1]. Cataract and glaucoma are ranked as the leading causes of blindness worldwide [2]. There are anatomical and physiological factors that influence IOP reduction after phacoemulsification. Pre-operative angle configuration is considered one of the main factors contributing to this variability. After lens removal, even eyes without glaucoma experience anatomical changes in the anterior chamber, and many biometrical factors are modified [3,4,5]. IOP reduction depends on anatomical factors, especially in patients with a narrow angle [6,7]. Different mechanisms that decrease IOP are as follows:

1. Lens removal allows the posterior capsule to move posteriorly, dislodging the zonula over the ciliary body with a consequent widening of the Schlemm's canal and increased aqueous humor drainage [8,9].
2. The ultrasound radiation used in the phacoemulsification procedure is responsible for a sudden rise in the anterior chamber pressure, producing inflammatory cytokines (mostly IL-1) that stimulate metalloproteinase production and trabecular meshwork remodeling, improving aqueous humor drainage [10,11].
3. Inflammation induced by cataract surgery leads to ciliary body shut down and reduction in IOP. Anterior capsular shrinkage after capsulorhexis leads to

traction on the ciliary body and decreases aqueous humor production [12].

2. MATERIALS AND METHODS

One hundred patients with visually significant cataract who visited the outpatient clinic in the Department of Ophthalmology, Assuit University Hospital, Assuit, Egypt between September 2016 and October 2017 were included in this prospective comparative study. All patients underwent cataract extraction by uneventful phacoemulsification.

2.1 Inclusion Criterion

Significant cataract affecting vision.

2.2 Exclusion Criteria

A history of trauma, glaucoma, or previous ocular surgery; complicated cataract; use of steroids as topical or systemic medications; and corneal opacity.

2.3 Ethical Aspects

Consent for participation in the study was obtained from the patients and their confidentiality was maintained. Refusal to participate in the study by any patient did not affect the quality of his/her treatment.

2.4 Pre-operative Evaluation

All the patients included in this study underwent the following clinical evaluations 1 day before the surgery:

1. Measurement of uncorrected and best corrected visual acuity by using Snellen chart.
2. Detailed ocular examination including anterior segment examination by Haag-Streit 900 series slit lamp and fundus examination by VOLK 90.

3. Measurement of IOP using the Goldmann applanation tonometer AT900, [Haag-Streit 900 series slit lamp].
4. Measurement of anterior chamber depth (ACD) by A-scan (EZ SCAN™ AB5500+ SONOMED, USA) which is one of the contact methods.
5. Medical fitness for all patients: Complete blood count, Blood pressure, Random blood sugar.

2.5 Surgical Technique

Cataract removal by phacoemulsification was performed under peribulbar anesthesia (lignocaine 2%). Draping of the eyelids and eyelashes was done to isolate these areas from the surgical field. Two small incisions and one main incision were made through the clear cornea and viscoelastic (Healon) was injected into the anterior chamber. Capsulorhexis was performed, which was continuous, curvilinear, and central. Hydro-dissection was performed to separate the nucleus and cortex from the capsule, which simplified the rotation of the nucleus.

The phaco-probe was introduced into the anterior chamber of the eye. Removal of the nucleus was performed by the divide and conquer, phaco chop (vertical or horizontal chop), or stop and chop technique. The lens cortex was removed by aspiration. Before the insertion of the intraocular lens, the capsule was filled with viscoelastic. The intraocular lens was then inserted in the capsular bag with a special injector. Viscoelastic was removed from the anterior chamber by aspiration. The incisions were sealed by stromal hydration. Finally, a subconjunctival injection of steroids and antibiotics was given to prevent infection.

2.6 Postoperative Evaluation

The patients were followed up at 1 week, 1 month, and 3 months postoperatively to measure

the IOP and ACD, and to identify any postoperative complications.

3. RESULTS AND DISCUSSION

The study included 100 eyes (52 left and 48 right eyes) of 100 patients (39 males and 61 females). Seventeen patients were younger than 50 years, 29 were between 50 and 60 years, 31 were between 60 and 70 years, and 23 patients were older than 70 years. The mean age of the patients was 60.70±10.04 years.

Preoperative and postoperative variation of ACD:

Preoperatively, the ACD was 3.19±0.39 mm. It was 3.45±0.45 mm at 1 week postoperatively, 3.64±0.45 mm at 1 month postoperatively, and 3.64±0.45 mm at 3 months postoperatively.

The ACD increased by 0.26±0.06 mm in the 1st postoperative week, by 0.45±0.06 mm in the 1st postoperative month, and by 0.59±0.08 mm by the 3rd postoperative month (Table 1).

Difference between preoperative and postoperative IOP:

The mean preoperative IOP was 15.28±3.12 mmHg; the corresponding values at 1 week, 1 month, and 3 months postoperatively were 14.28±2.82 mmHg, 13.76±2.66 mmHg, and 13.14±2.50 mmHg, respectively. The IOP decreased by 1.03±0.3 mmHg, 1.52 ± 0.46 mmHg, and 2.14±0.62 mmHg in 1 week, 1 month, and 3 months postoperatively (Table 2).

Correlation between IOP and ACD:

There was a statistically significant negative correlation between ACD and IOP at 1 week postoperatively ($r = -0.222$; $P = .027$) (Fig. 1) and 3 months postoperatively (Fig. 2) ($r = -.222$; $P = .026$); however, there was no statistically significant correlation between preoperative ACD and IOP and those at 1 month postoperatively (Table 3).

Table 1. Comparison between preoperative and postoperative ACD

ACD (mm)	Preoperative (n = 100)	After 1 week (n = 100)	After 1 month (n = 100)	After 3 months (n = 100)
Mean ± SD	3.19 ± 0.39	3.45 ± 0.45	3.64 ± 0.45	3.78 ± 0.47
Range	2.5 - 4.7	2.6 - 4.7	3.0 - 4.9	3.0 - 4.9
P-value		0.000*	0.000*	0.000*

Table 2. Comparison between preoperative and postoperative IOP

IOP (mmHg)	Preoperative (n= 100)	After 1 week (n= 100)	After 1 month (n= 100)	After 3 months (n= 100)
Mean ± SD	15.28 ± 3.12 mmHg	14.28 ± 2.82 mmHg	13.76 ± 2.66 mmHg	13.14 ± 2.50 mmHg
Range	13.0 - 22.0	12.0 - 19.0	11.0 - 20.0	11.0 - 19.0
P-value		0.001*	0.001*	0.000*

Table 3. Correlation between ACD and IOP

		Preoperative (mm)	1 week (mm)	1 month (mm)	3 months (mm)
IOP	Preoperative (mmHg)	r-value -0.160			
		P-value 0.111			
	1 week (mmHg)	r-value	-0.222		
		P-value	0.027*		
	1 month (mmHg)	r-value		-0.099	
		P-value		0.327	
	3 months (mmHg)	r-value			-0.222
		P-value			0.026*




Fig. 1. Correlation between ACD and IOP after one week

3.1 Discussion

The present study revealed that ACD was significantly increased at 1 week, 1 month, and 3 months after uneventful phacoemulsification cataract surgery; the respective increases in ACD were 0.1 mm, 0.2-0.5 mm, and 0.2-0.5 mm, respectively. The mean ACD was 3.19±0.39 mm preoperatively, 3.45±0.45 mm at 1 week postoperatively, 3.64±0.45 mm at 1 month postoperatively, and 3.78±0.47 mm at 3 months

postoperatively. The results of the study conducted by Mustafa Kamal Junejo, and associates [13] were similar to those of our study; they showed that there was a statistically significant increase in the mean ACD, from 3.02±0.43 mm preoperatively to 3.81±0.46 mm 1 month after the surgery (p<.0001)—an increase of 0.73±0.58 mm. However, when mean ACD 1 month after the surgery was compared between male (3.83 ± 0.49 mm) and female (3.77 ± 0.43 mm) groups, no statistically significant difference




Fig. 2. Correlation between ACD and IOP after three months

was observed ($p = .42$). In our study there are statistically significant differences of ACD between male and female groups at one week, one month, and three months postoperatively ($p = .037$, $p = .015$, and $p = .013$ respectively). In Mustafa Kamal Junejo, et al.'s study, 74 patients (74 eyes in 42 males and 32 females) underwent Ultrasonography A-scan to evaluate anterior chamber configuration before, one day after, one week after and one month postoperatively; however, in our study, we evaluated patients for longer period i.e., up to three months postoperatively. We also measured central ACD. Preoperative and postoperative data was compared by using paired tests [13]. Another study that is comparable to our study is by S Moghim et al. [14] which comprised 99 eyes of 99 consecutive patients who underwent cataract surgery, of which 14 eyes were excluded because of the inability to detect the scleral spur in anterior segment optical coherence tomography (AS-OCT) images. IOP and biometric parameters were measured by AS-OCT preoperatively and three months after surgery, unlike our study in which, IOP was measured by Goldman applanation tonometer and biometric parameters by A-scan. Out of the 85 patients included in the analysis, 35 were male and 50 were female with an average age of 62.2 ± 8.9 (37–81) years. Forty-six patients had

narrow angle glaucoma and 39 patients had open angle glaucoma; however, in our study, we excluded any glaucomatous patients. Preoperatively, the average IOP was 17.12 ± 2.47 mm Hg, which dropped to 12.20 ± 2.69 mm Hg at three months with an average change of 4.95 ± 2.26 mm Hg ($P < .001$) [14].

Another study that is in concordance with our study [15] showed that clear corneal phacoemulsification (CCP) was associated with a statistically significant reduction in IOP. The study was done on 273 normal patients selected for cataract extraction by phacoemulsification, however, our study included only 100 patients. IOP was measured by Goldman applanation tonometer by the same examiner preoperatively, on the 15th day, and subsequently one, two, three and six months after surgery. Patients with the history of ocular surgery, trauma, preoperative IOP greater than 21 mmHg, on ocular medication and who developed postoperative complication were excluded from the study [15]. The study showed that the mean IOP before surgery was 14.18 ± 3.4 mmHg. Patients showed a mean decrease in IOP of 2.25 mmHg (16%) compared to preoperative values. The postoperative IOP was inversely related to

preoperative ACD ($P=.012$). Changes of IOP was significantly correlated with change in ACD ($P = .002$) [10]. However, in our study, there is significant correlation between IOP and ACD measured at one week ($r= -.222$; $P\text{-value}= .03$) and three months postoperatively ($r= .222$; $P\text{-value}= .03$), and no statistically significant difference between IOP and ACD measured preoperatively ($r= -.16$;) and one month postoperative ($r= -.099$) [15].

One of the studies that had comparable results to our study [16] reported that there was increase in ACD, and widening of irideo corneal angle (ICA) in non-glaucomatous eyes after uneventful phacoemulsification. The changes were statistically significant over 6 months. The study was performed on Fifty-three eyes of 49 patients and they were evaluated for 6 months postoperatively [16].

The mean preoperative ICA grade of 2.97 ± 0.72 increased to 3.55 ± 0.48 at 1 week and 3.68 ± 0.45 at 1 month ($P=.03$). The mean preoperative ACD of 3.06 ± 0.49 mm increased to 3.57 ± 0.47 mm at 4 weeks, 3.69 ± 0.32 mm at 1 month, and 3.70 ± 0.36 mm at 3 months ($P=.012$) [16].

On the other hand, there were several studies that had results different from our results. These studies showed that there were no statistically significant differences between preoperative IOP and postoperative IOP. One of these studies [17] reported that no statistically significant difference in terms of changes in IOP could be attributed to phacoemulsification with posterior chamber intraocular lens type [17]. The study was performed on 103 eyes of 103 patients without ocular co-morbidity, all of whom underwent phacoemulsification. ACD and IOP were recorded 1-2 weeks preoperatively, 8-9 weeks postoperatively and 4 weeks after stopping steroid therapy. Mean preoperative and postoperative IOP was 15.23 (2.47) mmHg and 12.68 (1.65) mmHg respectively, and this represented a mean drop of 2.55 (1.78) mmHg. The extent of IOP reduction postoperatively was directly related to preoperative IOP ($r=.745$; $r^2=56\%$; $P=.01$) [17].

4. CONCLUSION

This study proves that there was a decrease in IOP and increase in ACD after cataract extraction by phacoemulsification with intraocular

lens implantation which help in aqueous drainage by pushing back iris lens diaphragm and widening anterior chamber angle. Low sample size and short term follow up are considered as main limitation of our study.

CONSENT

A written informed consent was obtained from all patients participated in the study after explaining the whole procedure and complications.

ETHICAL APPROVAL

All authors hereby declare that approval was obtained from the medical research ethics committee of the scientific research, faculty of medicine, Assiut University that adhered to the tents of the declaration of Helsinki.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. *Br J Ophthalmol.* 2006;90:262–267.
2. Pascolini D, Mariotti SP: Global estimates of visual impairment. *Br J Ophthalmol.* 2012;96:614–618.
3. Maloney WF, Dillman DM, Nichamin LD: Supracapsular phacoemulsification: A capsule-free posterior chamber approach. *J Cataract Refract Surg.* 1997;23:323–328.
4. Huang G, Gonzalez E, Lee R, Chen Y-C, He M, Lin SC. Association of biometric factors with anterior chamber angle widening and intraocular pressure reduction after uneventful phacoemulsification for cataract. *J Cataract Refract Surg.* 2012;38:108–116.
5. Shingleton BJ, Gamell LS, O'Donoghue MW, Baylus SL, King R. Long-term changes in intraocular pressure after clear corneal phacoemulsification: Normal patients versus glaucoma suspects and glaucoma patients. *J Cataract Refract Surg.* 2005;25:885–890.
6. Kanski JJ. *Clinical ophthalmology* (Sixth edition) Edinburgh: Elsevier Butterworth-Heinemann; 2007.

7. Yang HS, Lee J, Choi S. Ocular biometric parameters associated with intraocular pressure reduction after cataract surgery in normal eyes. *Am J Ophthalmol.* 2013;156: 89–94.
8. Shrivastava A, Singh K: The effect of cataract extraction on intraocular pressure. *Curr Opin Ophthalmol.* 2010;21:118–122.
9. Pohjalainen T, Vesti E, Uusitalo RJ, Laatikainen L. Intraocular pressure after phacoemulsification and intraocular lens implantation in nonglaucomatous eyes with and without exfoliation. *J Cataract Refract Surg.* 2001;27:426–431.
10. Mansberger SL, Gordon MO, Jampel HD, Borade A, Brandt JD, Wilson B, et al. Reduction in intraocular pressure after cataract extraction: The Ocular Hypertension Treatment Study. *Ophthalmology.* 2012;119:1826–1831.
11. Shin HC, Subrayan V, Tajunisah I. Changes in anterior chamber depth and intraocular pressure after phacoemulsification in eyes with occludable angles. *J Cataract Refract Surg.* 2010;36:1289–1295.
12. Takmaz T, Edinburgh P. Elsevier Butterworth-Heinemann clear corneal phacoemulsification: Normal patients versus glaucoma suspects and glaucoma patients. *J Sci.* 2013;43:289–293.
13. Junejo MK, Chaudhry TA. Anterior chamber depth changes after uneventful phacoemulsification. *Pak J Ophthalmol.* 2016;32(1):30-34.
14. Moghimi S, Abdi F, Latifi G, Fakhraie G, Ramezani F, He M, Lin SC. Lens parameters as predictors of intraocular pressure changes after phacoemulsification. *J Eye (Lond).* 2015;29(11):1469–1476.
15. Bhallil S, Andalloussi IB, Chraibi F, Daoudi K, Tahri H. Changes in intraocular pressure after clear corneal phacoemulsification in normal patients. *Oman J Ophthalmol.* 2009;2(3):111–113.
16. Altan C , Bayraktar S, Altan T, Eren H, Yilmaz OF. Anterior chamber depth, iridocorneal angle width, and intraocular pressure changes after uneventful phacoemulsification in eyes without glaucoma and with open iridocorneal angles. *Journal of Cataract & Refractive Surgery.* 2004;30(4):832-838.
17. Issa SA, Pacheco J, Mahmood U, Nolan J, Beatty S. A novel index for predicting intraocular pressure reduction following cataract surgery. *Br J Ophthalmol.* 2005; 89:543–546.

© 2019 Abdallah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47792>