

## Effect of intrauterine insemination treatment on sexual function and quality of life for infertile women

Emre Sinan Gungor<sup>1</sup>, Olcay Seval<sup>2</sup>,  
Gulşah Ilhan<sup>3</sup>, Fatma Ferda Verit<sup>4</sup>

### ABSTRACT

**Objectives:** To evaluate the effect of intrauterine insemination (IUI) on sexual functioning, quality of life and psychological well-being.

**Methods:** One hundred and thirty four infertile women going to IUI treatment as study group and 134 women who do not report any infertility complaint attending to gynecology clinic for routine control as control group were enrolled. Demographic data of the patients were collected. Patients were asked to complete Female Sexual Functioning Index (FSFI), Beck Depression Inventory (BDI) and SF 36 form.

**Results:** Total FSFI score (mean±SD) for study group was 23.4±4.1 and 24.8±3.4 for control group (p<0.05). This means a lower sexual function for patients going to IUI. There were also statistically significant differences according to subscales of FSFI scores for sexual desire, arousal and satisfaction. Mean±SD scores for Beck Depression Inventory analysis was 18.6±9.8 for study group and 18.5±7.1 for control group. According to SF-36 scores, there were statistically significant differences between the groups for four subscales: Role physical, bodily pain, general health and vitality.

**Conclusions:** Going to an IUI treatment has negative effects on FSFI scores and some of SF 36 scores but we did not find a significant negative effect on BDI scores.

**KEYWORDS:** Intrauterine insemination, Infertile women, Sexual function, Quality of life, depression.

doi: <https://doi.org/10.12669/pjms.344.14789>

### How to cite this:

Gungor ES, Seval O, Ilhan G, Verit FF. Effect of intrauterine insemination treatment on sexual function and quality of life for infertile women. *Pak J Med Sci.* 2018;34(4):891-896. doi: <https://doi.org/10.12669/pjms.344.14789>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### INTRODUCTION

Infertility is a disease characterized by the failure to establish a clinical pregnancy after 12 months of regular, unprotected sexual intercourse.<sup>1</sup> About

15% of the worldwide population is infertile and this condition expected to rise.<sup>2</sup>

Some infertile couples have some psychological difficulties, including; lack of marital satisfaction, impairment of relationships, lack of sexual satisfaction, forced timing of intercourse, loss of confidence in relation to sex, decreased libido and negative emotional effects.<sup>3-5</sup> There are many studies investigating the impact of infertility on female sexual function. Some of them reported higher prevalence of sexual dysfunction in infertile females,<sup>6,7</sup> whereas some others revealed no significant difference between fertile and infertile females.<sup>8,9</sup>

Infertility may also have a negative effect on the Quality Of Life (QOL) for the woman and maybe a cause of developing depression. QOL is defined as 'an individual's perception of their position in life in the context of the culture and value systems in which they live and concerns incorporating

1. Emre Sinan Gungor, M.D.
2. Olcay Seval, M.D.
3. Gulşah Ilhan, M.D.
4. Prof. Dr. Fatma Ferda Verit,
- 1-4: Suleymaniye Maternity Research and Training Hospital, Department of Obstetrics and Gynecology, Istanbul, Turkey.

Correspondence:

Dr. Emre Sinan Gungor, M.D.  
Suleymaniye Maternity Research and Training Hospital,  
Department of Obstetrics and Gynecology,  
Istanbul, Turkey.  
E-mail: doksinangungor@hotmail.com

- \* Received for Publication: February 6, 2018
- \* Revision Received: May 25, 2018
- \* Revision Accepted: June 5, 2018

physical health, psychological state, social relations and personal beliefs.<sup>10</sup> SF-36 is one of the best well defined test for assessing measure of self-reported health.

Depression may be another problem which a female may experience if she has infertility. In the literature, there are some articles concluding that infertility may cause depression and some articles deny such a relation.<sup>11-13</sup>

Intra Uterine Insemination (IUI) is a procedure performed at earlier stages of infertility treatment compared to In vitro Fertilization (IVF) and IVF is considered to be the end point treatment stage for infertility by the patients. So when compared with IVF procedure, starting an IUI cycle may be less stressful for the couple. To our knowledge, there is no study investigating the sexual problems and quality of life of females who are going to IUI treatment.

The aim of our study was to evaluate the effect of starting an IUI cycle on sexual functioning, quality of life and psychological well-being for females in comparison with a group of presumed fertile women.

## METHODS

Women aged between 20-40 years attending to infertility clinic of Suleymaniye Maternity Research and Training Hospital for IUI treatment were included in this prospective study as study group (Group 1) between January 2016 and February 2017. Women attending the gynecology clinic for general control without any complaints were taken as control group (Group 2). A total of 268 patients (134 study group and 134 control group) entered the study. One hundred forty two patients were invited to answer the questionnaires for the study group but six of them refused to fill the FSFI questionnaire and two of them answered the questionnaires but there were missing data; so these eight women were excluded from the study. One hundred forty five patients were invited to answer the questionnaires for the control group but six patients did not accept to fill the questionnaires and five patients' demographic data were missing. Non-pregnant and non-lactating, in a sexual relationship during the last 4 weeks were prospectively included in the study. Women with known psychiatric diseases, using antipsychotic drugs, genitourinary infection, genital tract abnormality, physical disabilities, and having chronic medical conditions associated with sexual dysfunction were excluded. Females with partners that had erectile or ejaculation disturbances associated with sexual dysfunction were also excluded.

All participants from the control group had at least one living child. None of the patients in the control group did not have any complaint while attending to hospital. They were just attending to gynaecology clinic for routine control. Infertile women included all those who were experiencing primary or secondary infertility. Participants who had never been able to conceive were diagnosed as primary infertility, whereas those with a previous history of pregnancy were diagnosed as secondary infertility.

Ethical committee approval was taken before starting the data collection. All participants gave informed consent after which demographic data and reproductive history of the patients were collected. Patients were asked to complete three self-administered questionnaires. Study group patients filled the questionnaires for the month just before starting to use the drugs related the IUI practice. All patients underwent stimulated IUI cycles, but administration of hormonal treatment is not a confounding variable because all our patients answered the questionnaires before starting any treatment. For the control group, patients were requested to answer the questionnaires after their physical examination evaluating the previous month. three self-administered questionnaires were as follows:

1. Short Form Health Survey SF-36: The SF-36 is a 36-item self-administered instrument that measures health status. It consists of eight subscales including Physical Functioning (PF), Role- Physical (RP) (limitations, difficulties due to one's physical problems), Bodily Pain (BP) (magnitude of pain), General Health (GH), Vitality (VT) (state of being strong and active), Social Functioning (SF), Role-Emotional (RE) (role limitations due to emotional problems) and Mental Health (MH). These eight subscales are scored from 0 to 100 where 100 indicates the highest level of health status.<sup>14</sup> SF 36 was adapted and validated for the Turkish population by Pinar et al.<sup>15</sup> SF36 can also be scored as two summary measures: Mental Component Summary (MCS) and Physical Component Summary (PCS). PF, RP, BP, GH are evaluated in PCS; and VT, SF, RE and MH are evaluated in MCS.
2. The Female Sexual Functioning Index (FSFI) is a 19-item multidimensional survey measuring six domains including sexual desire, arousal (both subjective and physiological), lubrication, orgasm, satisfaction, and pain. Each domain

was scored on a scale of 0 to 6, with higher scores indicating better sexual function for each domain.<sup>16</sup> It was validated for Turkish population and was used to assess female sexual function among women.<sup>17</sup> The individual domain scores were totaled and multiplied by a predetermined factor to weigh each domain equally.<sup>16</sup> A score  $\leq 26.5$  was defined as female sexual dysfunction.<sup>18</sup>

3. Beck Depression Inventory (BDI) was used to assess depression for the study and control groups. It consisted of 21 questions on particular aspects of depression related symptoms. Total score ranged from 0 to 63. A score of  $\geq 17$  was chosen as clinically significant symptoms of depression.<sup>18</sup> Turkish version of the BDI has been tested and validated in the Turkish population.<sup>19</sup>

**Statistical Analyses:** Mean, standard deviation, median, minimum, maximum value frequency and percentage were used for descriptive statistics. The distribution of variables was checked with kolmogorov-simironov test. Mann-whitney U test was used for the comparison of quantitative data. Chi-Square test was used for the comparison of the qualitative data. SPSS 22.0 was used for statistical analyses.  $p < 0.05$  was considered as statistically significant.

## RESULTS

The mean age was  $30.1 \pm 4.5$  for study group and  $29.5 \pm 4.5$  for control group. Also there was not a statistically significant difference for BMI

rates between the groups ( $26.2 \pm 3.6$  vs  $25.5 \pm 3.1$  respectively for study and control group). 97/134 (72.3%) of patients were primary infertile in study group. Mean duration of infertility was  $19.3 \pm 4.6$  months for study group. All patients in the control group had at least one child. Demographic characteristics of the patients are shown in Table-I.

Forty two out of 134 patients (31.3%) had an FSFI score more than 26.5 and 49/134 (36.5%) of our control group had an FSFI score more than 26.5. But when we analysed the overall FSFI score (mean  $\pm$  SD) for study group was  $23.4 \pm 4.1$  and  $24.8 \pm 3.4$  for control group ( $p < 0.05$ ). This means a lower sexual function in infertile women. There were also statistically significant differences between groups according to subscale of FSFI scores for sexual desire ( $3.6 \pm 1$  vs  $4.5 \pm 0.9$ , respectively  $p < 0.05$ ), arousal ( $4.1 \pm 1.1$  vs  $4.6 \pm 1$ , respectively,  $p < 0.05$ ) and satisfaction ( $3.7 \pm 1$  vs  $4.1 \pm 1$ , respectively,  $p < 0.05$ ). There were no statistically significant differences for other subscales of FSFI between groups.

The mean  $\pm$  SD scores for Beck Depression Inventory analysis was  $18.6 \pm 9.8$  for study group and  $18.5 \pm 7.1$  for control group and there was not a statistically significant difference. Twenty out of 134 (14.8%) of infertile patients experienced severe depression and 12/134 (8.9%) of control group had severe depression and this was not statistically significant too.

When the groups were analysed according to SF-36 scores, there were statistically significant differences between the groups for four subscales:

Table-I: Demographic characteristics of the patients.

		Study group		Control group		p
		Mean $\pm$ s.d.	Median	Mean $\pm$ s.d.	Median	
Age in years (mean $\pm$ SD)		$30.1 \pm 4.5$	30.0	$29.5 \pm 4.5$	29.0	0.167
Body Mass Index (kg/m <sup>2</sup> )		$26.2 \pm 3.6$	26.0	$25.5 \pm 3.1$	25.2	0.148
Gravidy		$0.1 \pm 0.3$	0.0	$1.0 \pm 0.8$	1.0	0.000
Parity		$0.3 \pm 0.5$	0.0	$1.4 \pm 0.9$	1.0	0.000
Educational Status	Primary School	$83 \pm 61.5\%$		$67 \pm 49.6\%$		0.057
	High School	$45 \pm 33.3\%$		$52 \pm 38.5\%$		
	University	$7 \pm 5.2\%$		$16 \pm 11.9\%$		
Family status	Nuclear	$131 \pm 97.0\%$		$130 \pm 96.3\%$		0.735
	Extended	$4 \pm 3.0\%$		$5 \pm 3.7\%$		
Occupational status	Employee	$36 \pm 26.7\%$		$43 \pm 31.9\%$		0.349
	Housewife	$99 \pm 73.3\%$		$92 \pm 68.1\%$		
Marital Status	Married	$135 \pm 100.0\%$		$135 \pm 100.0\%$		-
	Single	$0 \pm 0.0\%$		$0 \pm 0.0\%$		
Use of antidepressants		$15 \pm 11.1\%$		$12 \pm 8.9\%$		0.543
Cigarette smoking		$40 \pm 29.6\%$		$36 \pm 26.7\%$		0.588
Use of alcohol		$13 \pm 9.6\%$		$11 \pm 8.1\%$		0.669
Doing exercise		$19 \pm 14.1\%$		$31 \pm 23.0\%$		0.060

s.d. Standard Deviation.

Table-II: Mean BDI, FSFI, SF 36 scores of study and control group.

	Study Group		Control Group		p
	Mean±s.d.	Median	Mean±s.d.	Median	
Beck Depression Score	18.6±9.8	18.0	18.5±7.1	18.0	0.798
Normal	64±47.4%		57±42.2%		
Mild depression	51±37.8%		66±48.9%		
Severe depression	20±14.8%		12±8.9%		
<b>FSFI</b>					
Total FSFI	23.4±4.1	23.6	24.8±3.4	24.9	0.001
Desire	3.6±1.0	3.6	4.5±0.9	4.2	0.000
Arousal	4.1±1.1	4.2	4.6±1.0	4.8	0.000
Lubrication	4.1±1.0	4.2	4.1±0.9	3.9	0.215
Orgasm	3.8±1.2	4.0	3.9±0.9	4.0	0.641
Satisfaction	3.7±1.0	3.6	4.1±1.0	4.0	0.015
Dyspareunia	4.1±1.0	4.0	4.3±4.6	3.6	0.096
<b>SF-36</b>					
Physical function	66.1±19.6	70.0	65.5±18.4	65.0	0.728
Role Physical	63.5±28.0	75.0	71.9±26.9	75.0	0.018
Bodily pain	58.4±20.3	57.5	63.8±23.0	67.5	0.017
General health	51.6±12.0	50.0	56.9±16.4	55.0	0.002
Vitality	50.3±11.7	50.0	55.4±13.0	55.0	0.000
Social functioning	60.2±19.4	62.5	58.1±21.9	50.0	0.340
Role emotional	50.4±33.1	66.7	53.1±29.2	66.7	0.531
Mental health	53.5±12.7	52.0	56.6±15.2	56.0	0.109

BDI: Beck Depression Inventory, FSFI: Female Sexual Functioning Index, SF-36: Short Form Health Survey.

Role physical (63.5±28 vs 71.9±26.9 respectively,  $p<0.05$ ), bodily pain (58.4±20.3 vs 63.8±23 respectively,  $p<0.05$ ), general health (51.6±12 vs 56.9±16.4 respectively,  $p<0.05$ ) and vitality (50.3±11.7 vs 55.4±13 respectively,  $p<0.05$ ). There were no statistically significant differences for other subscales of SF-36 between the groups. Scores for all patients and for all questionnaires are shown in Table-II.

There was a positive correlation between total FSFI scores and SF36 physical function score. There was also a positive correlation between FSFI dyspareunia score and SF 36 physical function scores ( $p<0.05$ ). We could not find any other correlation between FSFI scores and SF 36 scores (Table-III).

## DISCUSSION

Sexual function is one of the important components of health and overall quality of life. Infertility has a major negative impact on women's and also couples quality of life and sexual life. It is associated with feelings of loss of control, diminished self-esteem, anxiety and depression.<sup>7</sup>

As we found lower FSFI scores for infertile patients, similar to us Oddens et al and Drosdzol et al reported lower sexual life satisfaction for infertile patients when compared with healthy controls.<sup>11,20</sup>

Oddens et al found significant difference for desire and satisfaction subgroups; Drosdzol et al found difference for arousal, orgasm and pain subgroups. We found significant difference between groups for sexual desire, arousal and satisfaction subgroups of FSFI. Also Millheiser et al and Hentschel et al reported lower FSFI scores for sexual desire and arousal subdomains.<sup>6,21</sup> From these results it may be concluded that, sexual desire and arousal are the most problematic subdomains in infertile patients. However problems in satisfaction subgroup is revealed to be common in Turkish women.<sup>17</sup>

Prior studies have indicated that infertile women report more depressive symptoms than controls. In a study of 338 infertile women and 39 healthy controls, women who had a duration of infertility between two to three years reported a higher prevalence of depression compared with the control group.<sup>22</sup> In a study conducted in the Netherlands, 281 women seeking infertility treatment reported higher rates of depressive symptoms compared to the 289 healthy controls.<sup>11</sup> However, another recent study concluded that depression is uncommon in women seeking infertility treatment.<sup>12</sup> Although it may be anticipated that infertility may cause depression, like Brasile et al we also could not find different BDI scores between infertile patients and control group. The relationship between female infertility

Table-III: Correlation between FSFI and SF 36 scores.

		<i>Physical function</i>	<i>Role Physical</i>	<i>Bodily Pain</i>	<i>General health</i>	<i>Vitality</i>	<i>Social functioning</i>	<i>Role emotional</i>	<i>Mental health</i>
Total FSFI	r	0.162	0.036	0.041	0.062	-0.011	-0.055	-0.052	0.065
	p	0.008	0.555	0.502	0.311	0.857	0.366	0.395	0.288
Desire	r	-0.047	0.065	0.002	0.085	0.077	-0.011	0.020	0.000
	p	0.442	0.287	0.969	0.163	0.209	0.855	0.741	0.999
Arousal	r	0.079	0.100	0.031	0.079	-0.017	0.039	0.038	0.052
	p	0.197	0.101	0.615	0.195	0.781	0.519	0.538	0.390
Lubrication	r	0.077	-0.089	-0.055	0.041	-0.084	-0.091	-0.138	0.085
	p	0.206	0.143	0.366	0.504	0.166	0.137	0.052	0.163
Orgasm	r	-0.050	-0.043	-0.058	-0.055	-0.025	-0.073	-0.059	-0.090
	p	0.413	0.480	0.339	0.368	0.683	0.232	0.330	0.140
Satisfaction	r	0.116	-0.043	0.101	0.077	0.078	-0.015	-0.048	0.074
	p	0.057	0.478	0.098	0.207	0.199	0.808	0.431	0.228
Dyspareunia	r	0.225	0.066	0.081	-0.030	0.050	-0.009	-0.044	0.116
	p	0.000	0.279	0.185	0.618	0.412	0.877	0.469	0.057

and depression remains somewhat unclear. As 99% of people living in our country, Turkey, is Muslim; our population may accept infertility as a destiny. As a result of this acceptance, although infertility affects sexual function and some of SF 36 scores, its effect on BDI scores is limited.

Drosdzol et al and Souter et al reported lower QoL among infertile women but different from us Souter et al reported lower scores in all SF-36 categories as we found lower scores for role physical, bodily pain, general health and vitality subgroups.<sup>20,23</sup> These results are comparable to ours. We could find limited correlation between FSFI and SF 36 scores. This correlation was only on physical function subdomain.

Although İstanbul is the biggest and most well-known city of Turkey, our hospital's patients are generally from lower income group and come from eastern cities of Turkey. Different from Western cultures where marriages are thought to be result of love, for such a population like ours, marriage is based on family constitution. Therefore low marital satisfaction especially among women may be a common problem. Higher BDI scores, lower quality of life and higher frequencies of sexual problems may be more common for both our infertile and also fertile group of patients.

**Limitations of the study:** The sample size might be larger to generalize the findings. Another thing we did not evaluate was the husband's sexual function

index. The etiology of infertility in these couples was not assessed.

Our SF 36 analyzes also shows similar results that generally physical component of the questionnaire is negatively effected for infertile patients (role physical, bodily pain and general health). Only vitality subgroup of SF 36 questionnaire which is related with mental component summary is worser in infertile patients. The second finding of our study is that there are no group differences in symptoms of depression, is a good news that adds to existing evidence that, although infertility is distressing, psychopathology is no more common among infertile than fertile women. Infertile patients may perceive IVF treatment as the last chance of having a baby and a more stressful process but our study demonstrated that planning an IUI treatment also have unfavourable effects. Although there are several studies reporting investigating the QOL and sexual problems of infertile women going to IVF or ICSI treatment,<sup>24</sup> to our knowledge it is the first study to date investigating FSD and quality of life for infertile women going to IUI treatment.

## CONCLUSIONS

We investigated the effect of going to an IUI treatment on sexual functioning and quality of life on infertile women and found lower FSFI and SF 36 scores but we did not find a significant negative effect on BDI scores. Worse sexual functioning and

quality of life probably indicates that anticipating intrauterine insemination treatment is a stressful life event. We may help them by normalizing their feelings and by explaining that it is common to feel less interested in sex leading up to treatment and make sure that they may seek help from a counsellor if this persists.

**Grant Support & Financial Disclosures:** None.

**Disclosure statement:** The authors report no conflicts of interest.

## REFERENCES

- Zegers-Hochschild F, Adamson GD, Dyer S, Racowsky C, de Mouzon J, Sokol R, et al. The International Glossary on Infertility and Fertility Care, 2017. *Hum Reprod.* 2017;32:1786-1801. doi: 10.1093/humrep/dex234.
- Thoma ME, McLain AC, Louis JF, King RB, Trumble AC, Sundaram R, et al. Prevalence of infertility in the United States as estimated by the current duration approach and a traditional constructed approach. *Fertil Steril.* 2013;99(5):1324-1331. doi: 10.1016/j.fertnstert.2012.11.037.
- Ashraf DM, Ali D, Azadeh DM. Effect of Infertility on Sexual Function: A Cross-Sectional Study. *J Clin Diag Res.* 2015;9:1-3. doi: 10.7860/JCDR/2015/11148.5934.
- Damti OB, Sarid O, Sheiner E, Zilberstein T, Cwikel J. Stress and distress in infertility among women. *Harefuah.* 2008;147(3):256-260, 276.
- Cousineau T, Domar A. Psychological impact of infertility. *Best Pract Res Clin Obstet Gynaeco.* 2007;21(2):293-308. doi: 10.1016/j.bpobgyn.2006.12.003
- Millheiser LS, Helmer AE, Quintero RB, Westphal LM, Milki AA, Lathi RB. Is infertility a risk factor for female sexual dysfunction? A case-control study. *Fertil Steril.* 2010;94(6):2022-2025. doi: 10.1016/j.fertnstert.2010.01.037.
- Pakpour AH, Yekaninejad MS, Zeidi IM, Burri A. Prevalence and risk factors of the female sexual dysfunction in a sample of infertile Iranian women. *Arch Gynecol Obstet.* 2012;286:1589-1596. doi: 10.1007/s00404-012-2489-x
- Furukawa AP, Patton PE, Amato P, Li H, Leclair CM. Dyspareunia and sexual dysfunction in women seeking fertility treatment. *Fertil Steril.* 2012;98:1544-1548. doi: 10.1016/j.fertnstert.2012.08.011.
- Iris A, Aydogan Kirmizi D, Taner CE. Effects of infertility and infertility duration on female sexual functions. *Arch Gynecol Obstet.* 2013;287(4):809-812.: 1589-1596.
- WHOQOL-Group. The World Health Organization Quality of Life Assessment (WHOQOL): position paper from the World Health Organization. *Social Sci Med* 1995;41:1403-1409.
- Oddens BJ, den Tonkelaar I, Nieuwenhuyse H. Psychosocial experiences in women facing fertility problems-A comparative survey. *Hum Reprod.* 1999;14:255-261.
- Brasile D, Katsoff B, Check JH. Moderate or severe depression is uncommon in women seeking infertility therapy according to the beck depression inventory. *Clin Exp Obstet Gynecol.* 2006;33:16-8.
- Joelsson SL, Tyden T, Wanggren K, Georgakis MK, Stern J, Berglund A, et al. Anxiety and depression symptoms among sub-fertile women, women pregnant after infertility treatment, and naturally pregnant women. *Eur Psychiatry.* 2017;45:212-219. doi: 10.1016/j.eurpsy.2017.07.004.
- Moghadam S, Zeinalzadeh A, Salavati M, Almasi S, Negahban H. Adaptation, reliability and validity testing of a Persian version of the Health Assessment Questionnaire-Disability Index in Iranian patients with rheumatoid arthritis. *J Bodyw Mov Ther.* 2017;21(1):133-140. doi: 10.1016/j.jbmt.2016.07.001.
- Pınar R. A new concept in health investigations: Quality of life; test of validity and reliability of a quality of life scale for patients with chronic diseases. *Nursing Bulletin.* 1995;9:85-95.
- Rosen RC, Brown C, Heiman J, Leiblum S, Meston C, Shabsigh R, et al. The Female Sexual Function Index (FSFI): A multidimensional self-report instrument for the assessment of female sexual function. *J Sex Marital Therapy.* 2000;26:191-208. doi: 10.1080/009262300278597
- Oksuz E, Malhan S. Reliability and validity of the Female Sexual Function Index in Turkish population. *Sendrom.* 2005;17(7):54-60.
- Kucur KS, İlay G, Aysenur A, Kerem HG, Eda UU, Pasa U, et al. Effects of Infertility Etiology and Depression on Female Sexual Function. *J Sex Marital Ther.* 2016;42:27-35. doi: 10.1080/0092623X.2015.1010673
- Hisli NS. Validity and reliability of Beck Depression Inventory between university students. *Psychology J.* 1989;7:3-13.
- Drosdzol A, Skrzypulec V. Quality of life and sexual functioning of Polish infertile couples. *Eur J Contracept Reprod Health Care.* 2008;13:271-281. doi: 10.1080/13625180802049187.
- Hentschel H, Alberton DL, Sawdy RJ, Capp E, Goldim JR, Passos EP. Sexual function in women from infertile couples and in women seeking surgicalsterilization. *J Sex Marital Ther.* 2008;34:107-114. doi: 10.1080/00926230701636171.
- Domar AD, Broome A, Zuttermeister PC, Seibel M, Friedman R. The prevalence and predictability of depression in infertile women. *Fertil Steril.* 1992;58:1158-1163.
- Souter VL, Hopton JL, Penney GC, Templeton AA. Survey of psychological health in women with infertility. *J Psychosom Obstet Gynecol.* 2002;23:41-49.
- Smith NK, Madeira J, Millard HR. Sexual Function and Fertility Quality of Life in Women Using In Vitro Fertilization. *J Sex Med.* 2015;12:985-993. doi: 10.1111/jsm.12824.

## Authors' Contribution:

**ESG:** Conceived, designed and editing of manuscript.

**OS, GI:** Did data collection and manuscript writing.

**FFV:** Did review and final approval of manuscript.