



Effect of Organic Manures and Potassium Levels on Growth and Yield Attributes of Pearl Millet (*Pennisetum glaucum* L.)

Tiravaturu Madhan^{a++*} and C. Umesh^{a#}

^a Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj- 211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102750

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104798>

Original Research Article

Received: 09/06/2023

Accepted: 13/08/2023

Published: 24/08/2023

ABSTRACT

The field experiment was conducted during *rabi* 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj (U.P.) India. To study the Response of Organic manures and potassium on growth and yield of Pearl millet The treatment combinations were T1: FYM 5t/ha + 30 kg/ha potassium, T2: FYM 5t/ha + 40 kg/ha potassium, T3: FYM 5t/ha + 50 kg/ha potassium, T4: Vermicompost 5t/ha + 30 kg/ha potassium, T5: Vermicompost 5t/ha + 40kg/ha potassium, T6: Vermicompost 5t/ha + 50kg/ha potassium, T7: Poultry Manure 4t/ha + 30kg/ha potassium, T8: Poultry Manure 4t/ha + 40kg/ha potassium, T9: Poultry Manure 4t/ha + 50kg/ha potassium, and control plot T10:(RDF80:40:40 NPK kg/ha) . Results obtained that the plant height (168.06 cm), plant dry weight(48.87 g/plant),crop growth rate (24.0 g/m²/day), ear head length (25.07cm) number

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

^{*}Corresponding author: E-mail: tmadhan02@gmail.com;

of grains/ear head (2372.69), test weight (10.92 gm), seed yield (36.90 q/ha) and stover yield (68.03 q/ha) were significantly influenced with application of Poultry Manure 4t/ha + 50 kg/ha potassium. Higher gross return (INR 86707.17/ha), higher net return (INR 58590.17/ha) and highest B:C ratio (2.08) were also recorded in treatment-9 (Poultry Manure 4t/ha + 50 kg/ha potassium).

Keywords: Economics; growth parameters; organic manures; potassium; farm.

1. INTRODUCTION

“Pearl millet [*Pennisetum glaucum* (L.)] is one of the important millet crops of India. Among the coarse cereals, it stands next to sorghum in area and production. Rajasthan ranks first in area and production of pearl millet. Besides being a staple diet of approximately 10 per cent population of our country, it is also used as important fodder crop. It is the only cereal crop that is able to produce reliable yield under the marginal environments and simultaneously responds to excessive management conditions. Its nutritious grain forms the important component of human diet and stover forms the principal maintenance ration for ruminant livestock during the dry season. In addition, pearl millet grain is increasingly being used as feed for livestock and poultry” [1].

“Vermicompost is the product of decomposition process utilizing earthworms to turn the organic waste into high-quality compost that consists mainly of worm cast in addition to decayed organic matter” [1]. “Vermicomposting helps to convert the organic wastes (agro-wastes, animal manure and domestic refuse) into highly nutrient fertilizers for plant and soil” [2].

2. MATERIALS AND METHODS

The experiment was conducted in Randomized Block Design (RBD) which includes ten treatments that are replicated thrice and was laid out with the different treatments allocated randomly in each replication. “The soil of the experimental field was sandy loam in texture, slightly alkaline reaction (pH 7.1) with low level of organic carbon (0.48%), available N (225 Kg/ha), P (13.6 kg/ha) and higher level of K (215.4 kg/ha) and the treatment combinations are as follows T1: FYM 5t/ha + 30kg/ha potassium, T2: FYM 5t/ha + 40 kg/ha potassium, T3: FYM 5t/ha + 50kg/ha potassium, T4: Vermicompost 5t/ha + 30 kg/ha potassium, T5: Vermicompost 5t/ha + 40kg/ha potassium, T6: Vermicompost 5t/ha + 50kg/ha potassium, T7: Poultry Manure 4t/ha + 30kg/ha potassium, T8: Poultry Manure 4t/ha + 40kg/ha potassium, T9: Poultry Manure 4t/ha +

50kg/ha potassium, and control plot T10:(RDF 80:40:40 NPK kg/ha) are used” [3].

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant height

The Growth parameters were recorded at 20, 40, 60 and at 80 DAS.

The significantly higher plant height (196.73cm) was observed in treatment-9 (Poultry Manure 4 t/ha + Potassium 50 Kg/ha). it might be due to with application of poultry manure, “the beneficial response of poultry manure to yield attributes might also be attributed to the availability of sufficient amounts of easily utilizable from of plant nutrients throughout the growth period and especially at critical growth periods of crop resulting in better uptake, plant vigour and superior yield attributes” [4].

“Further increase in plant height is due to with the application of Potassium in higher level play's crucial role in meristematic growth through its effects on the synthesis of Phytohormones, among various plant hormones, cytokinin plays an important role in growth of the plant” [5].

3.1.2 Dry weight

The Plant dry weight of pearl millet was recorded at 20, 40, 60, 80 DAS at Harvest differed significantly as influenced by Organic manures and Potassium levels. At 80 DAS, the higher plant dry weight (14.08 gm/plant) was observed in treatment-9 (Poultry Manure 4 t/ha + Potassium 50 kg/ha). However, treatment-8 (Poultry Manure 4 t/ha + Potassium 40 kg/ha) was found to be statistically at par with treatment- 9 (Poultry Manure 4 t/ha + Potassium 50 kg/ha). “The significantly higher plant dry weight (14.08 gm/plant) was observed with the application of Poultry manure, the increase in the total dry matter production may be due to better source and sink capacity developed due to better

dry matter production and its accumulation in assimilatory surface area and increase in the photosynthetic efficiency and thus increased the production of photosynthesis reflected in better growth and ultimately in higher dry accumulation” [6].

Where as increase in dry matter might be due to with the application of potassium it involved in number of physiological processes, protein synthesis and activation of enzymes. In the present investigation, the crop responded up to 50 kg K₂O/ha. Potassium aggregating agent which is known to have positive effect on soil physical properties such as plant height, healthy growth etc., and subsequently crop yields.

3.2 Yield Parameters

3.2.1 Ear head length

The significant and higher ear head length (25.07 cm) was observed in treatment-9 with (Poultry Manure 4 t/ha + Potassium 50 kg/ha), which was significantly superior over rest of the treatments. However, treatment-8 (Poultry Manure 4 t/ha + Potassium 40 kg/ha), was found to be statistically at par with treatment-9 (Poultry Manure 4 t/ha + Potassium 50 kg/ha).

“Potassium aggregating agent which is known to have positive effect on soil physical properties such as plant height, healthy growth etc and subsequently crop yields” [7].

3.2.2 Number of grains/ear head

The significant and higher number of grains/ear head (2372.69) was observed in treatment-9 with (Poultry Manure 4 t/ha + Potassium 50 kg/ha), which was significantly superior over rest of the treatments. However, treatment-8 (Poultry Manure 4 t/ha + Potassium 40 kg/ha), was found to be statistically at par with treatment-9 (Poultry Manure 4 t/ha + Potassium 50 kg/ha).

Further increase in number of grains with the significant effect of potassium, Application of potassium improved the number of grains per head which might be due to the favorable effects of potassium on nutrient uptake, photosynthetic activity, improving its mobilization reported by Yadav et al. [8].

3.2.3 Grain yield (q/ha)

The significant and higher grain yield (36.90) was observed in treatment-9 with (Poultry Manure 4

t/ha + Potassium 50 kg/ha), which was significantly superior over rest of the treatments. However, treatment-8 (Poultry Manure 4 t/ha + Potassium 40 kg/ha), was found to be statistically at par with treatment-9 (Poultry Manure 4 t/ha + Potassium 50 kg/ha).

These findings are conformity with Jagadeesha et al. [9]. Further, increase in grain yield with application of Potassium at 50 kg/ha it stimulates the cumulative effect improvement in yield attributes viz., ear head length, number of grains/ear head, test weight and enhances the development of strong cell walls and therefore stiffer straw which might be resulted into profuse tillering [10].

3.2.4 Stover yield (q/ha)

The significant and higher grain yield (68.03) was observed in treatment-9 with (Poultry Manure 4 t/ha + Potassium 50 kg/ha), which was significantly superior over rest of the treatments. However, treatment-8 (Poultry Manure 4 t/ha + Potassium 40 kg/ha), was found to be statistically at par with treatment-9 (Poultry Manure 4 t/ha + Potassium 50 kg/ha). And also, an increase in uptake of plant nutrients empowered the plant to manufacture more quantity of photosynthates resulting in more stover yield. Similar results were reported by Thumar et al. [11].

3.3 Economic Analysis

Observations regarding economics of different treatments of pearl millet are given in Table 3.

3.3.1 Gross returns (INR/ha)

The maximum gross return (86,707.17 INR/ha) was found in treatment-9 (Poultry Manure 4t/ha + Potassium 50 kg/ha).

3.3.2 Net returns (INR/ha)

The maximum net return (58,590.17INR/ha) was found in treatment-9 (Poultry Manure 4t/ha + Potassium 50 kg/ha).

3.3.3 Benefit cost ratio (B:C)

The maximum benefit Cost ratio (2.08) was found in treatment-9 with Poultry Manure 4t/ha + Potassium 50 kg/ha [12].

Table 1. Effect of organic manures and potassium on yield attributes of pearl millet

S.No.	Treatments	80 DAS		60-80 DAS	
		Plant height (cm)	Plant dry Weight (g)	Crop growth rate (g/m ² /day)	Relative growth rate (g/g/day)
1.	FYM 5t/ha + Potassium 30 kg/ha	163.71	37.97	16.1	0.015
2.	FYM 5t/ha + Potassium 40 kg/ha	165.80	39.61	18.4	0.016
3.	FYM 5t/ha + Potassium 50 kg/ha	168.96	41.66	20.5	0.018
4.	Vermicompost 5t/ha + Potassium 30 kg/ha	169.71	42.55	20.4	0.017
5.	Vermicompost 5t/ha + Potassium 40 kg/ha	171.87	43.84	21.2	0.017
6.	Vermicompost 5t/ha + Potassium 50 kg/ha	173.96	45.68	21.9	0.017
7.	Poultry Manure 4t/ha + Potassium 30 kg/ha	172.27	44.96	19.3	0.015
8.	Poultry Manure 4t/ha + Potassium 40 kg/ha	175.05	47.60	22.6	0.017
9.	Poultry Manure 4t/ha + Potassium 50 kg/ha	178.06	48.87	24.0	0.018
10.	Control(RDF 80:40:40 NPK kg/ha)	165.23	38.64	16.0	0.014
	S Em(±)	1.95	0.61	1.06	0.001
	CD (P=0.05)	5.79	1.82	3.14	-

Table 2. Effect of organic manures and potassium on yield attributes of pearl millet

S. No.	Treatments	Ear head length (cm)	Grains /Ear head	Test weight (g)	Seed yield (q/ha)	Stover yield (q/ha)	Harvest index (%)
1.	FYM 5t/ha + 30kg/ha potassium	17.07	2023.79	8.50	24.26	54.56	30.75
2.	FYM 5t/ha + 40kg/ha potassium	17.87	2079.36	9.07	26.74	57.87	31.61
3.	FYM 5t/ha + 50kg/ha potassium	18.91	2184.02	9.21	28.25	59.11	32.40
4.	Vermicompost 5t/ha + 30kg/ha potassium	20.61	2084.47	9.32	27.60	58.65	32.04
5.	Vermicompost 5t/ha + 40kg/ha potassium	21.42	2146.27	10.17	31.20	62.70	33.26
6.	Vermicompost 5t/ha + 50kg/ha potassium	22.02	2148.41	10.41	32.60	64.69	33.50
7.	Poultry Manure 4t/ha + 30kg/ha potassium	23.39	2188.61	8.87	28.16	61.00	31.57
8.	Poultry Manure 4t/ha + 40kg/ha potassium	24.14	2246.42	10.21	33.43	63.99	34.20
9.	Poultry Manure 4t/ha + 50kg/ha potassium	25.07	2372.69	10.92	36.90	68.03	35.18
10.	Control(80:40:40 NPK kg/ha)	19.30	2030.02	8.85	25.69	55.12	31.80
	F-Test	S	S	NS	S	S	NS
	SEm±	0.46	32.27	0.58	0.85	1.97	0.91
	CD (P=0.05)	1.35	96.37	--	3.53	5.85	--

Table 3. Effect of organic manures and potassium on economic analysis of pearl millet

S. No.	Treatments	Cost of cultivation (INR/ha)	Gross return (INR/ha)	Net return (INR/ha)	B:C Ratio
1.	FYM 5t/ha + 30kg/ha potassium	27450.00	57018.83	29568.83	1.08
2.	FYM 5t/ha + 40kg/ha potassium	27783.00	62839.56	35056.00	1.26
3.	FYM 5t/ha + 50kg/ha potassium	28117.00	66379.67	38262.67	1.36
4.	Vermicompost 5t/ha + 30kg/ha potassium	29950.00	64852.17	34902.17	1.17
5.	Vermicompost 5t/ha + 40kg/ha potassium	30283.00	73320.45	43037.00	1.42
6.	Vermicompost 5t/ha + 50kg/ha potassium	30617.00	76610.83	45993.00	1.50
7.	Poultry Manure 4t/ha + 30kg/ha potassium	27450.00	66183.83	38733.83	1.41
8.	Poultry Manure 4t/ha + 40kg/ha potassium	27783.00	78153.17	50370.17	1.81
9.	Poultry Manure 4t/ha + 50kg/ha potassium	28117.00	86707.17	58590.17	2.08
10.	Control(80:40:40 NPK kg/ha)	23450.00	60379.33	36929.33	1.57

4. CONCLUSION

It is concluded that in pearl millet with the combination of poultry manure (4 t/ha) along with Potassium (50 kg/ha) (Treatment-9) was observed highest grain yield and benefit cost ratio.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Devi J, Prakash M. Microbial population dynamics during vermicomposting of three different substrates amended with cow dung. *International Journal of Current Microbiology and Applied Sciences*. 2015;4(2):1086-1092.
- Gajalakshmi S, Abbasi SA. Earthworms and vermicomposting. *Indian Journal of Biotechnology*. 2003;3:486-494.
- Yadav BB, Singh R, Indu T, Pradhan A. Effect of Spacing and Potassium Levels on Growth and Yield of Foxtail Millet (*Setaria italica* L.). *International Journal of Plant & Soil Science*. 2023;35(5):46-52.
- Fazily T, Hanshul CS. Effect of organic manures on yield and economics of late sown wheat (*Triticum aestivum*). *International Journal of Research & Review*. 2019;6(1):168-171.
- Chauhan NN, Chaudhary FK, Patel HN. Bio-efficacy of different insecticides against pearl millet earhead worm, *Helicoverpa armigera* (Hub.) in summer

- pearl millet: *Pennisetum glaucm* (L). *Trends in Biosciences*. 2017;10(45):9270-9274.
- Bhattacharyya P, Chakraborty A, Bhattacharya B, Chakrabarti K. Evaluation of MSW compost as a component of integrated nutrient management in wetland rice. *Compost Science & Utilization*. 2003;11(4):343-350.
 - Srinivasa DK, Chikkaramappa T, Basavaraja PK, Sukanya TS, Murali K, Chamegowda TC. Status of different forms of potassium under foxtail millet crops influenced by graded levels of potassium in alfisols of Chikkaballapura region, Karnataka. *International Journal of Chemical Studies*. 2019;7(3): 3435-3441.
 - Yadav S, Irfan M, Ahmad A, Hayat S. Causes of salinity and plant manifestations to salt stress: A review. *Journal of Environmental Biology*. 2011;32(5):667.
 - Jagadish SVK, Cairns J, Lafitte R, Wheeler TR, Price AH, Craufurd PQ. Genetic analysis of heat tolerance at anthesis in rice. *Crop Science*. 2010;50(5):1633-1641.
 - Kacha DJ, Khafi HR, Mehta AC, Shekh MA, Jadav RP. Effect of potassium and zinc on yield and quality of rabi pearl millet (*Pennisetum glaucum*). *Crop Research*. 2011;41(1, 2 & 3):31-34.
 - Thumar NJ, Wei QH, Hu WH. Recent advances in asymmetric metal-catalyzed carbene transfer from diazo compounds toward molecular complexity. *Advances in Organometallic Chemistry*. 2016;66: 33-9.

12. Roopashree DH, Nagaraju YM, Ramesha T Bhagyalakshmi, Raghavendra S. Effect of integrated nutrient management on growth and yield of baby corn (*Zea mays* L.). *International Journal of Current Microbiology and Applied Sciences*. 2019;8(6):766-772.

© 2023 Madhan and Umesh; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104798>