



## **Sensory Evaluation of Different Value Added Products of Aonla (*Emblica officinalis* G.) during Storage**

**Ashwini Uikey<sup>a\*</sup>, P. K. S. Gurjar<sup>b#</sup> and Shubham Rathour<sup>a\*</sup>**

<sup>a</sup> *Department of Horticulture, College of Agriculture, RVSKVV, Gwalior, India.*

<sup>b</sup> *Krishi Vigyan Kendra, RVSKVV, Gwalior-474002, Madhya Pradesh, India.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJECC/2022/v12i111377

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92533>

**Original Research Article**

**Received 06 August 2022**

**Accepted 11 October 2022**

**Published 19 October 2022**

## **ABSTRACT**

A research experiment was conducted during the year 2019-20 and 2020-21 in the Fruit and Vegetable Processing Unit Laboratory, Department of Horticulture, College of Agriculture, Gwalior with seven different varieties of aonla viz., NA-4, NA-5, NA-6, NA-7, NA-10, Laxmi and Chakaiya. It was studied about the preparation of various value added products such as aonla toffee, with beverages like nectar and squash. Observations were recorded up to storage period of six months at ambient conditions. This study is analyzed in a completely randomised design with three replications. Laxmi and Chakaiya varieties were found superior with respect to colour, appearance, taste and aroma as well as flavour. It has been noticed that the overall acceptability of processed products have been found significant with the storage intervals such as 30, 60, 90 and 120 days. As far as sensory evaluation of different aonla products was concerned, the Laxmi and Chakaiya varieties are found to be superior for the purpose of the processing industry.

*Keywords: Aonla; processing industry; storage; value addition.*

<sup>o</sup> *Ph.D Research Scholar;*

<sup>#</sup> *Scientist (Horticulture);*

<sup>\*</sup>*Corresponding author: E-mail: ashwiniuikey@gmail.com;*

## 1. INTRODUCTION

“Aonla is one of the oldest minor fruit and considered to be a wonder for human health. India ranks 1<sup>st</sup> in aonla area and production all over the world. It belongs to the family Euphorbiaceae and comprises about 350 pieces” [1]. The fruit has high indigenous medicinal value such as an anti- ascorbic, laxative and antibiotic. Aonla is used in Ayurvedic systems of Indian medicines [2]. “Area and production of aonla has increased from 67,000 hectares to 1,08,000 hectares according to the world wide record. In India, aonla is grown in area of about 50,000 hectares with a production of around 200000 metric tonnes. The area under aonla has been expanding rapidly in the last couple of years” [3].

“The main cultivated varieties are Banarasi (Drying), Bansi Red, Chakaiya (Pickle, Candy and Syrup), Desi, Krishna/NA-5(Candy and jam), Kanchan/NA-4 (Candy and Jam), Franchis (Hathijhool), NA-6, NA-7 (Candy and Jam), NA-8, NA-9, NA-10 and Anand-7 (Pickle)” [3]. Besides, it is not consumed as fresh or in the raw state as it is acidic and astringent. Therefore, it is not so much popular as a table fruit. The excellent nutritive and therapeutic value offers [4-7]. great potential for processing it into several quality products like preserve, squash, candy, jelly, jam, syrup, pickle, chutney, powder, barfi, segments in sugar syrup, supari, blended beverage, carbonated drinks and ready-to-serve [8-11].

Due to its perishable nature and glut after harvest, which lowers the market value of fruit, aonla experiences postharvest losses that range from 30 to 40% [12-14]. Therefore, the only practical method for future economic usage of expanded aonla output would be value addition through processing. Traditional procedures were time-consuming and unhygienic [15-21]. Due to this, the nutritive losses are seen in higher amounts. Therefore an attempt is made to find out the suitability of some varieties for processing as well as stability of different aonla products.

## 2. MATERIALS AND METHODS

### 2.1 Plant Material

Fully matured aonla fruits of different varieties such as NA-4, NA-5, NA-6, NA-7, NA-10, Laxmi and Chakaiya. etc are harvested from the main experimental station of Department of Horticulture, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior during the year 2019-20 and 2020- 21.

### 2.2 Chemicals and Additives

All chemicals used in this investigation were of analytical grade. Cane sugar, skimmed milk powder, glucose, butter, citric acid, hot spices, vegetables and salt were obtained from a local market and used as ingredients for the preparation of different products such as aonla toffee, nectar and squash.

### 2.3 Extraction of Pulp

Aonla fruits that were fully developed, had a firm texture, and were all the same size were blanched and utilised in the experiment. Using a small-scale pulping equipment, the fruits were processed to extract pulp, yielding a fine pulp. The final pulp was prepared with an addition of 2% potassium bisulphite solution for 15-20 minutes.

### 2.4 Standardization of Recipes

#### 1. Aonla toffee

Aonla toffees were prepared from seven different varieties of aonla with the standardized recipe. The other ingredients, such as sugar, skimmed milk powder, glucose, butter and salt were kept constant.

#### Preparation of aonla toffee recipe

Aonla pulp	1 kg
Sugar	750 gm
Glucose	100 g
Skimmed milk powder	150 g
Butter	50 g

#### 2. Aonla nectar

Aonla nectar was prepared from seven different varieties of aonla with the standardized recipe. The other ingredients, such as sugar, citric acid and water were kept constant as per required for the preparation of nectar.

#### Preparation of aonla nectar Recipe

Aonla pulp	20 per cent
Total soluble solids	15 per cent
Acidity	0.3 per cent
SO <sub>2</sub>	70 ppm

#### 3. Aonla squash

Aonla squash were prepared from seven different varieties of aonla with the standardized recipe of that beverage. The other ingredients, such as sugar, citric acid and water were kept

constant as per required for the preparation of squash.

Aonla squash type of fruit beverage, which contain at least 25 per cent juice and 45 per cent total soluble solids. It also contains about 1 per cent acidity and 350 ppm sulphur dioxide and 600 ppm sodium benzoate (in coloured juices). It is diluted before being served.

### Recipe

Aonla pulp	25 per cent
Sugar	50 per cent
Acid	1.2 per cent
Sodium metabisulphite	350 ppm

### 2.5 Preparation of Value Added Products

Following standardization, five value added products of aonla were prepared using the standardized recipe of toffee, nectar and squash.

1. Aonla toffee
2. Aonla nectar
3. Aonla squash

### 2.6 Organoleptic Analysis


“The organoleptic evaluation of the various processed products for assessing the colour, flavour, texture, appearance etc., were done by a panel of 6 judges who allotted marks on the basis of 9.0 Point Hedonic Scale” [22].


### 2.7 Statistical Analysis

The data were analysed according to a factorial completely randomized design (FCRD) with three replications for statistical significance, as specified by Panse and Sukhatme [23].


## 3. RESULTS AND DISCUSSION

The recovery of aonla pulp was found to be 950 g/kg of fruit without straining and the various value added products such as toffee, nectar and squash were prepared from seven different varieties NA-4, NA-5, NA-6, NA- 7, NA-10, Laxmi and Chakaiya.






**Fig. 1. Flow sheet for the preparation of aonla toffee**



**Fig. 2. Flow sheet of preparation of aonla nectar**



**Fig. 3. Flowsheet on preparation of aonla squash**

### 3.1 Colour and Appearance

The effect of treatments on Colour and Appearance of Aonla Toffee is depicted in Table 1. Variety Chakaiya was (mean score 7.71) ranked highest for colour and appearance followed by variety Laxmi. Similarly, the effect of treatments on Colour and Appearance of Aonla Nectar was observed in Table 2. Variety NA-7 was (mean score 6.83) ranked highest for colour and appearance followed by variety Chakaiya. The effect of treatments on Colour and Appearance of Aonla Squash is depicted in Table 3, Chakaiya Variety was highest rank (meanscore 7.18) was recorded.

### 3.2 Taste

In organoleptic evaluation, taste is very important factor after colour and appearance. Statistical analysis revealed a significant effect of treatment on taste of various value added products of aonla. Variety Chakaiya (7.39 score) got higher scores of taste in Aonla Toffee was followed by (7.57 score) in variety Laxmi (Table 1), while the variety Chakaiya (8.48 score) got higher scores of taste in Aonla Nectar was followed by (8.41 score) in Variety NA-10 (Table 2).

Similarly, variety Chakaiya (7.39 score) got higher scores of taste in Aonla Squash was followed by (7.33 score) in variety Laxmi (Table 3).

### 3.3 Flavour

The varieties had pronounced the effect on flavour of fruits. The varieties significantly affected the flavour value on days of storage. In case of aonla toffee, maximum flavour taste

(7.91) was in NA-10 variety as compared to minimum (6.91) in NA-4 variety (Table 1). Similarly it has been reported that, maximum flavour taste (6.84) was in Chakaiya variety as compared to minimum (6.20) in NA-4 variety in aonla nectar (Table 2). The Chakaiya variety showed higher flavour value (7.96) at the end of storage period as compared to lower value (5.00) in NA-4 variety in aonla squash (Table 3).

### 3.4 Aroma

Aroma scores of the value added products increased with the varieties. The maximum aroma value (7.57) was observed in Chakaiya variety followed by Laxmi (7.30) and NA-10 (7.22) in aonla toffee (Table 1). In case of aonla nectar, the maximum aroma was observed (7.41) in Chakaiya variety as compared to minimum aroma score (6.91) in NA-4 variety (Table 2). In case of aonla squash, the maximum aroma (6.31) was recorded in Chakaiya variety while minimum aroma (5.87) was recorded in NA-4 variety (Table 3). The interaction effect of various varieties and products on flavour was found to be significant at the beginning of the storage.

### 3.5 Overall Acceptability

Overall acceptability was influenced with the varieties. It decides the consumer's preference to the product. Higher level of value added products could not produce top acceptability due to deviation from standard colour, Taste, Flavour and Aroma. Though, the best result was recorded (7.60 score) in Laxmi variety, it is closely followed by (7.36 score) in Chakaiya variety. No certain pattern was observed with overall acceptability with the varieties concerned.

**Table 1. Effect of varieties on organoleptic properties of aonla toffee**

Aonla toffee	Sensory score				
	Colour and appearance	Taste	Aroma	Flavour	Overall acceptability
NA-4	6.88	6.16	6.34	6.91	6.67
NA-5	6.90	6.33	6.46	7.21	6.70
NA-6	7.38	6.67	6.67	7.41	6.83
NA-7	7.06	6.68	6.92	7.25	6.92
NA-10	7.41	7.33	7.22	7.91	7.10
Laxmi	7.52	7.57	7.30	7.60	7.25
Chakaiya	7.71	7.39	7.57	7.41	7.33
SeM±	0.702	0.653	1.062	0.758	0.543
CD at 5% (1 10)	0.332	0.308	0.501	0.357	0.256

**Table 2. Effect of treatments on organoleptic properties of aonla nectar**

Aonla Nectar	Sensory score				
	Colour and appearance	Taste	Aroma	Flavour	Overall acceptability
NA-4	6.50	8.01	6.24	6.20	6.55
NA-5	6.00	8.06	6.39	6.50	6.67
NA-6	6.67	8.07	6.44	6.50	6.77
NA-7	6.83	8.26	6.76	6.70	6.80
NA-10	6.67	8.41	6.83	6.71	6.92
Laxmi	6.75	8.28	7.25	6.80	7.00
Chakaiya	6.79	8.48	7.41	6.84	7.10
SeM±	0.628	0.758	0.940	0.854	0.152
CD at 5% (1-10)	0.296	0.357	0.443	0.403	0.072

**Table 3. Effect of treatments on organoleptic properties of aonla squash**

Aonla Squash	Sensory score				
	Colour and appearance	Taste	Aroma	Flavour	Overall acceptability
NA-4	5.97	6.58	5.87	5.00	6.80
NA-5	6.30	6.61	5.89	5.50	6.87
NA-6	6.64	6.70	5.92	6.00	6.99
NA-7	6.50	7.01	5.97	6.00	7.05
NA-10	6.81	7.20	5.99	7.50	7.21
Laxmi	6.92	7.33	6.02	7.80	7.35
Chakaiya	7.18	7.39	6.31	7.96	7.43
SeM±	0.103	0.114	0.540	0.153	0.553
CD at 5% (1-10)	0.581	0.054	0.255	0.544	0.261

#### 4. CONCLUSION

It can be concluded from this study that, the products can be effectively used as alternative source of value addition in toffee, nectar and Squash. On the basis of results obtained it may be concluded that the varieties Laxmi and Chakaiya can be used in commercialization of different value added products. Even after a few days of storage, all of the sensory parameters continue to be within acceptable limits despite a modest reduction during storage.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Hooker JD. The flora of British India. Vol. VI. 1st Indian Reprint, Periodical Experts, Delhi; 1973.
- Singh Om, Singh Richa. Development and evaluation of aonla based blended nectar drink from different fruits using stevia for low calorie. Plant Archives. 2014;14 (1):115–9.
- Seema Shekhawat, Rathore, N.S. and Kaushik, R.A. (2014). “Advances in processing and Product development of aonla (*Emblica Officinalis*) in Indian context- A review. International Journal of Food and Nutritional Sciences. e – ISSN; 3(6):242-247.
- Nath V, Sharma RK. Screening of aonla (*Emblica officnailis* Gaertn,) cultivars for processing. Progressive Horticulture. 1998;30(1 & 2):76-77.
- Pareek S, Rathore NS, Kaushik RA. Aonla (*Emblica officnailis* Gaertn.): Post harvest handling and processing technology. Rajasthan College of Agriculture, Techn. Bull. No. 1, Udaipur, India. 2011;45.
- Priya MD, Khatkar BS. Effect of processing methods on keeping quality of Aonla (*Emblica officinalis* G.) preserve. International Food Research Journal. 2013;20(2):617-622.
- Priyanka Nayak, Tandon, Dileep Kumar, Bhatt Devendra Kumar. Study on changes of nutritional and organoleptic quality of flavoured candy prepared from aonla (*Emblica officinalis* G.) during storage.

- International Journal of Nutrition and metabolism. 2012;4(7):100-106.
8. Anonymous. Annual Report, AICRP on arid fruits, Tech. Doc. 1988;18.
  9. Balaji Vikram, Purnima Singh Sikarwar. Studies on preparation of value added herbal kinnow–aonla beverages (RTS and Squash) during storage. Int. J. Pure App. Biosci. 2017;6(1):758-765. ISSN; 2320-7051
  10. Jain, Akansha, Gehlot Rakesh, Siddiqui Saleem, Mukesh R. Jangra. Optimization of recipe for development of aonla squash –A response surface methodology approach. Journal of Applied and Natural Science. 2016;8(3):1341–1348.
  11. Jat ML, Shrivam JS, Jat RK, Pragati. Commercial products of aonla fruits, increasing the value addition. Journal of Pharmacognosy and Phytochemistry. 2021;10(1):1331-1337.
  12. Kalra CL, The Aonla. Indian Food Packer. 1988;42:67-83.
  13. Khushbu, Mahendra Bairwa, Vijay Bahadur, Manisha K Rolaniya, Jayvanth Kumar. Study on changes in organoleptic quality of value added blended herbal ready-to-serve (RTS) of aonla (*Emblica officinalis* Gaertn.) CV. NA-6 during storage period. Journal of Pharmacognosy and Phytochemistry. 2018;708-711.
  14. Madhu Goyal, Kavita Sharma, Vrinda Kiradoo. New vistas of value addition to utilize Amla (*Emblica officinalis*) and Ber (*Ziziphus mauritiana*) fruits. Journal of Dairying, Foods & H.S. 2008;27(2): 145-147.
  15. Rajendra Kumar. Studies on process standardization and storage behaviour of ready to serve (RTS) beverage prepared from aonla cultivars. Journal of Pharmacognosy and Phytochemistry. 2018;7(6):74-77.
  16. Rakesh, Arya SS, Moond SK. Processed products of aonla. Processed Food Industry. 2004;7(4):20-23.
  17. Singh IS, Pathak RK. Evaluation of aonla (*Emblica Officinalis* Gaertn.) varieties for processing. Acta Horticulturae. 1987;208: 173-177.
  18. Singh V. Studies on processing and storage stability of aonla (*Emblica Officinalis* Gaertn.) products. Ph. D. Thesis, Narendra Dev University of Agriculture and Technology, Kumarganj, Faizabad (U.P.); 1999.
  19. Sumana Devi, Ena Gupta, Neelesh Kumar Maurya. Development of a value added Amla Product. Int. Arch. App, Sci. Techno; 2020.
  20. Visen, Gaurav Singh, Pathak, Sanjay, Mishra, Krishna Kumar. Studies on physico-chemical composition of Aonla (*Emblica officinalis* G.) fruit and evaluation of ginger extract concentration for preparation of aonla candy cv. Chakaiya. Int. J. Curr. Microbiol. App. Sci. 6 2017; (11):3536- 3540.
  21. Yadav, Prem Kant, Vishwakarma Govind, and Yadav DK. Studies on determination of storage stability of Aonla products. Res. Environ. Life Sci. 2015;8(3):425-43.
  22. Amerine MA, Pangborn RM, Rossler EB. Principles of sensory evaluation of Food. Academic Press. New York. 1965; 283-289.
  23. Panse VG. Sukhatme PV. Statistical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi. 1967;381.

© 2022 Uikey et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/92533>