



Coriander Yield Characteristics as Influenced by Varied Date of Sowing and Planting Geometry under Chhattisgarh Plain Zone

**Sandeep Kumar Painkara^{a*}, D. K. Chandrakar^a,
P. M. Paraye^a, A. K. Verma^b, Golmei Langangmeilu^b
and Anjali Patel^b**

^a *College of Agriculture and Research Station, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh, 492012, India.*

^b *College of Agriculture, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh, 492012, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i23936

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112401>

Original Research Article

Received: 26/11/2023

Accepted: 31/01/2024

Published: 03/02/2024

ABSTRACT

A field experiment was carried out during the rabi season of 2020-21 and 2021-22 at Instructional farm, College of Agriculture and Research Station, IGKV, Raigarh, Chhattisgarh to find out the influence of different sowing dates and planting geometry on coriander. The experiment was laid out in split plot design with sixteen treatment combinations of main and sub plots, replicated thrice. Four Dates of sowing viz. D1: 25th October, D2: 10th November, D3: 25th November and D4: 10th December were arranged in main plot and four planting geometries viz., S1: 30 x 5 cm, S2: 30 x 7.5 cm, S3: 30 x 10 cm and S4: 30 x 12.5 cm were taken as subplot treatments. Results revealed that coriander sown on 25th October registered significantly yield attributing characters viz., number of

*Corresponding author: E-mail: skpainkara15@gmail.com;

umbels plant⁻¹, number of umbellets umbel⁻¹, number of umbellets plant⁻¹, number of seeds umbel⁻¹, number of seeds umbellet⁻¹, length of umbel, diameter of umbel, weight of umbels, test weight, seed yield plant⁻¹ and seed yield ha⁻¹. Among planting geometry, sowing of coriander at 30 x 12.5 cm spacing produced significantly higher values of aforesaid characters except for seed yield ha⁻¹, which was maximum under 30 x 10 cm spacing. The interaction effect of 25th October coupled with the spacing 30 x 12.5 cm produced significant maximum seed yield plant⁻¹.

Keywords: Coriander; date of sowing; planting geometry; yield attributes; umbel.

1. INTRODUCTION

Coriander (*Coriandrum sativum* L.) is a significant annual spice and medicinal herb belonging to the Apiaceae family, native to Eastern Europe and Asia. Cultivated for its fragrant seeds, commonly known as "Dhania," it possesses carminative, diuretic, stomachic, and aphrodisiac properties when dried. The essential oil extracted from coriander seeds during stem distillation comprises major components such as linalool (67.7%), 1-pipen (10.5%), 1-terpinin (9.0%), geranyl acetate (4%), and geraniol (1.9%).

Successful coriander production is influenced by genetic, weather, and agronomic factors Szemplinski and Nowak, [1] Optimal fruit and essential oil yields are achieved when the right combination of these factors is provided [2,3]. The date of sowing plays a crucial role in coriander management, affecting the photoperiodic response, yields, and qualities [4]. Changes in sowing time impact weather microclimates and subsequently crop performance. The physical environment, including temperature, humidity, and rainfall, can limit plant growth and production individually or collectively.

Establishing suitable spacing for maintaining the optimum plant population per unit area is essential for maximizing yields in any crop. Adequate planting geometry supports better crop growth and development, ultimately leading to higher production. Enhancing the productivity of both local and improved coriander varieties through the adoption of recommended sowing dates and planting geometries is crucial in contemporary agriculture. Therefore, an effort was made to examine the effect of date of sowing and planting geometry on productivity of coriander under loamy sandy soil of Chhattisgarh plains.

2. MATERIALS AND METHODS

The study was conducted over two consecutive *Rabi* seasons, namely 2020-21 and 2021-22, at

the Instructional Farm of the College of Agriculture and Research Station, Raigarh, IGKV, Chhattisgarh. The location is positioned at 21.9' North latitude, 83.4' East longitude, and has an altitude of 215 meters above mean sea level. The experimental soil at the site is characterized as loamy sand, slightly acidic, low in organic carbon, and deficient in available nitrogen, phosphorus, sulfur, and zinc. However, it exhibits medium availability of potassium and has low moisture retention capacity.

The experimental design employed a split-plot layout with sixteen treatment combinations in main and sub-plots, replicated thrice. The main plots involved four different sowing dates: 25th October, 10th November, 25th November, and 10th December. The subplots consisted of four planting geometries: 30 x 5 cm, 30 x 7.5 cm, 30 x 10 cm, and 30 x 12.5 cm. The recommended fertilizer dose of 80:60:40 kg N: P₂O₅: K₂O ha⁻¹ was applied. The impact of these treatments on coriander cultivation's yield characteristics was examined by collecting data on number of umbels plant⁻¹, number of umbellets umbel⁻¹, number of umbellets plant⁻¹, number of seeds umbel⁻¹, number of seeds umbellet⁻¹, umbel length, umbel diameter, weight of umbels, test weight, seed yield plant⁻¹, and seed yield ha⁻¹, and analysed statistically, using the F-test following Gomez and Gomez's [5] procedure, was performed. Significance of mean differences among treatments was determined using Critical Difference (CD) values at P=0.05.

3. RESULTS AND DISCUSSION

3.1 Effect of Sowing Dates

Coriander planted on October 25th exhibited notably higher yield-related characteristics, including the number of umbels plant⁻¹, number of umbellets umbel⁻¹, number of umbellets plant⁻¹, number of seeds umbel⁻¹, number of seeds umbellet⁻¹, umbel length, umbel diameter, weight of umbels, test weight, seed yield plant⁻¹, and seed yield ha⁻¹ compared to other sowing dates (Table 1 and 2). This trend was observed

Table 1. Coriander yield characteristics as influenced by different date of sowing and planting geometry

Treatment	No. of umbels plant ⁻¹			No. of umbellets umbel ⁻¹			No. of umbellets plant ⁻¹			No. of seeds umbel ⁻¹			No. of seeds umbellet ⁻¹			Length of Umbel (cm)			Diameter of umbel (cm)			Weight of umbel (g)		
	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean
Date of sowing																								
25 th October	21.33	23.16	22.25	5.03	5.49	5.26	109.38	129.24	119.31	32.06	34.86	33.46	6.65	7.11	6.88	10.66	11.31	10.98	8.32	9.19	8.75	0.34	0.37	0.35
10 th November	18.07	19.37	18.72	4.65	5.04	4.84	86.51	100.15	93.33	29.68	31.39	30.53	6.46	6.83	6.64	10.37	10.97	10.67	7.97	8.82	8.39	0.30	0.34	0.32
25 th November	16.31	17.28	16.79	4.16	4.52	4.34	68.66	79.70	74.18	24.60	26.02	25.31	5.97	6.32	6.15	9.47	9.89	9.68	7.84	8.53	8.19	0.28	0.31	0.29
10 th December	13.53	14.32	13.92	4.00	4.28	4.14	55.20	62.75	58.97	21.29	22.42	21.86	5.36	5.61	5.48	9.18	9.51	9.34	7.14	7.72	7.43	0.25	0.28	0.26
SEm±	0.62	0.58	0.46	0.15	0.13	0.14	4.59	3.62	3.51	0.63	0.82	0.69	0.20	0.19	0.18	0.20	0.20	0.20	0.17	0.19	0.18	0.01	0.01	0.01
CD (P=0.05)	2.15	2.01	1.61	0.51	0.46	0.44	15.89	12.53	12.16	2.17	2.85	2.40	0.62	0.65	0.61	0.69	0.69	0.69	0.60	0.66	0.63	0.03	0.03	0.03
Planting geometry																								
30 cm x 5 cm	11.86	12.72	12.29	3.97	4.29	4.13	47.87	55.75	51.81	20.23	21.58	20.91	5.12	5.33	5.22	8.99	9.34	9.16	7.23	7.89	7.56	0.22	0.24	0.23
30 cm x 7.5 cm	14.95	16.13	15.54	4.26	4.61	4.44	64.73	75.49	70.11	25.61	27.34	26.47	6.04	6.33	6.18	9.61	10.08	9.85	7.73	8.45	8.09	0.27	0.31	0.29
30 cm x 10 cm	20.25	21.72	20.98	4.62	5.01	4.81	94.87	110.95	102.91	29.88	31.78	30.83	6.54	7.07	6.80	10.45	11.02	10.73	8.00	8.80	8.40	0.33	0.36	0.34
30 cm x 12.5 cm	22.17	23.57	22.87	4.98	5.43	5.20	112.28	129.64	120.96	31.92	33.99	32.96	6.74	7.15	6.95	10.63	11.23	10.93	8.30	9.12	8.71	0.35	0.39	0.36
SEm±	0.44	0.48	0.36	0.10	0.13	0.11	2.65	3.62	2.31	0.51	0.72	0.58	0.16	0.21	0.16	0.14	0.14	0.14	0.10	0.11	0.11	0.01	0.01	0.01
CD (P=0.05)	1.27	1.41	1.04	0.28	0.39	0.32	7.73	10.56	6.74	1.50	2.11	1.70	0.46	0.61	0.47	0.40	0.40	0.40	0.30	0.34	0.31	0.02	0.03	0.02
Interaction	-	-	-	-	-	-	*	*	*	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

* Significant at 0.05% probability (Table 3)

Table 2. Growth of coriander as influenced by different date of sowing and planting geometry

Treatment	Test weight (g)			Seed yield Plant ⁻¹ (g)			Seed yield (q ha ⁻¹)		
	20-21	21-22	Mean	20-21	21-22	Mean	20-21	21-22	Mean
Date of sowing									
25 th October	12.98	13.82	13.40	5.27	5.84	5.55	18.16	20.35	19.26
10 th November	12.12	12.94	12.53	4.48	4.91	4.70	14.86	16.65	15.76
25 th November	11.03	11.66	11.34	3.86	3.98	3.92	12.40	13.14	12.77
10 th December	10.84	11.24	11.04	3.13	3.35	3.24	9.72	10.80	10.26
SEm±	0.16	0.21	0.12	0.13	0.15	0.12	0.59	0.68	0.45
CD (P=0.05)	0.55	0.72	0.43	0.45	0.51	0.42	2.06	2.34	1.53
Planting geometry									
30 cm x 5 cm	10.52	10.99	10.75	1.81	2.02	1.91	10.79	12.49	11.64
30 cm x 7.5 cm	11.52	12.21	11.87	3.28	3.60	3.44	13.28	14.74	14.01
30 cm x 10 cm	12.31	13.05	12.68	5.39	5.75	5.57	16.01	17.39	16.70
30 cm x 12.5 cm	12.61	13.41	13.01	6.26	6.71	6.48	15.06	16.30	15.68
SEm±	0.18	0.17	0.14	0.12	0.14	0.12	0.63	0.74	0.62
CD (P=0.05)	0.53	0.50	0.40	0.36	0.41	0.35	1.84	2.16	1.80
Interaction	-	-	-	*	*	*	-	-	-

* Significant at 0.05% probability (Table 3)

Table 3. Interaction effects between different sowing dates and planting geometry on number of umbellets plant⁻¹ and seed yield plant⁻¹

Number of umbellets plant ⁻¹															
2020-21					2020-21					Mean					
Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	
Spacing					Spacing					Spacing					
S ₁	69.47	44.98	37.33	26.38	S ₁	86.36	52.54	43.45	30.04	S ₁	77.91	48.76	40.39	28.21	
S ₂	93.57	64.01	52.04	35.45	S ₂	109.76	82.15	60.79	44.95	S ₂	101.66	73.08	56.42	40.20	
S ₃	126.24	109.44	72.03	57.43	S ₃	151.99	122.85	89.30	63.92	S ₃	139.12	116.14	80.67	60.68	
S ₄	147.91	127.91	88.30	69.33	S ₄	170.70	144.34	108.53	81.28	S ₄	159.30	136.13	98.41	75.31	
SEM±	4.31				SEM±	3.608				SEM±	2.663				
CD (P=0.05)	12.57				CD (P=0.05)	10.530				CD (P=0.05)	7.772				
Seed yield plant ⁻¹															
Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	Sowing date	D ₁	D ₂	D ₃	D ₄	
Spacing					Spacing					Spacing					
S ₁	2.45	1.59	0.86	0.56	S ₁	2.87	1.87	1.27	0.80	S ₁	2.66	1.73	1.06	0.68	
S ₂	4.14	3.29	2.46	1.62	S ₂	4.71	3.60	2.92	1.86	S ₂	4.43	3.45	2.69	1.74	
S ₃	6.77	5.50	4.51	2.81	S ₃	7.46	6.04	4.98	3.03	S ₃	7.11	5.77	4.74	2.92	
S ₄	7.98	6.34	5.26	3.38	S ₄	8.74	7.27	5.52	3.75	S ₄	8.36	6.81	5.39	3.57	
SEM±	0.161				SEM±	0.193				SEM±	0.162				
CD (P=0.05)	0.470				CD (P=0.05)	0.563				CD (P=0.05)	0.474				

D₁: 25th October, D₂: 10th November, D₃: 25th November, D₄: 10th December; S₁: 30 cm x 5 cm, S₂: 30 cm x 7.5 cm, S₃: 30 cm x 10 cm, S₄: 30 cm x 12.5 cm

consistently across both years and in the pooled analysis. Early sowing dates contribute to ample moisture availability, improved nutrition, maximal interception of photosynthetically active radiation (PAR), and increased synthesis of photosynthates. This leads to luxurious plant growth, resulting in enhanced growth and yield characteristics and ultimately an increase in seed yield. Similar findings were reported by Khoja [6] in fenugreek, Anitha et al. [7] in fenugreek, and Kiran [8] in black cumin.

3.2 Effect of Planting Geometry

Regarding planting geometry, the use of 30 x 12.5 cm spacing resulted in significantly higher values for the number of umbels plant⁻¹, number of umbellets umbel⁻¹, number of umbellets plant⁻¹, number of seeds umbel⁻¹, number of seeds umbellet⁻¹, umbel length, umbel diameter, weight of umbels, test weight, and seed yield plant⁻¹ over the two years and on average, whereas the seed yield ha⁻¹ was recorded significantly maximum under 30 x 10 cm spacing (Table 1 and 2). The observed outcome can be attributed to the heightened growth and yield characteristics in the 30 cm x 12.5 cm planting configuration. This success is a consequence of sufficient moisture availability, improved nutrition, and optimal interception of photosynthetically active radiation (PAR). The resulting profuse branching and higher biomass accumulation per plant contribute to increased growth and yield characteristics, ultimately leading to a higher seed yield. The abundant branching appears to stimulate increased flowering initiation, and the augmented supply of metabolites, facilitated by the higher biomass plant⁻¹, likely aids in flower retention. This, in turn, promotes greater seed formation and seed growth. While, under wider spacing i.e. 30 x 12.5 cm, reduced plant population per unit area though, improved overall growth of crop, but due to less plants per units area of crop it failed to record highest yield per hectare. Similar findings were documented by Ajay et al. [9], Diwan et al. [10], and Nethravathi [11].

3.3 Interaction Effects

The interaction between sowing date and planting geometry did not show significance in various aspects of coriander yields, however, there were notable variations in number of umbellets plant⁻¹ and seed yield plant⁻¹ due to these interactions. Specifically, the combined effect of sowing on October 25th along with a

spacing of 30 x 12.5 cm resulted in significantly higher number of umbellets plant⁻¹ and seed yield plant⁻¹. This performance was comparable to the combination of October 25th sowing with 30 x 10 cm spacing across both years and on average (Table 3). Kiran (2018) also reported comparable findings in black cumin, supporting similar results.

4. CONCLUSION

The findings from the two-year experiments lead to the conclusion that coriander crops sown on October 25th and utilizing a spacing of 30 x 12.5 cm exhibited significantly superior yield characteristics and the per hectare yield was maximum under 30 x 10 cm spacing. Hence, for yield maximization of coriander, it should be sown with 30 x 10 cm spacing in 2nd fortnight of October.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Szemplinski W, Nowak J. Nitrogen fertilization versus the yield and quality of coriander fruit (*Coriander sativum* L.). Acta Scientiarum Polonorum Hortorum Cultus. 2015;14(3): 37-50.
2. Rangappa M, Bhardwaj HL, Showhda M, Hamama AA. Cilantro (*Coriandrum sativum* L.) response to nitrogen fertilizer rates. Journal of Herbs, Spices and Medicinal Plants. 1997;5(1):63-68.
3. Gill BS, Randhawa GS, Saini SS. Effect of sowing dates and herb cutting management on growth and yield of fenugreek (*Trigonella foenum graecum*). Indian Journal of Agronomy. 2001;46(2):364-367.
4. Kuri B, Jat N, Shivran A, Puniya M.. Productivity and profitability of coriander varieties influence by sowing time and plant growth regulators. Annals of Agricultural Research. New Series. 2015; 36(2):204-211.
5. Gomez KA, Gomez AA. Statistical procedure for Agricultural Research., edn 2, John Wiley & Sons, New York. 1984;241-271.
6. Khoja JR. Effect of sowing time and sources of nitrogen on growth, thermal

- requirement, yield and quality of coriander (*Coriandrum sativum* L.). Ph.D. thesis. Rajashan Agricultural University. Bikaner; 2005.
7. Anitha B, Reddy MLN, Rao AVDD, Kiran Patro TSJK, Salomi Suneetha DR. Effect of sowing date on yield and quality of fenugreek. *Plant Archives*. 2016;16(1):479-484.
 8. Kiran MR. Effect of sowing time and plant geometry on growth, yield and quality of black cumin (*Nigella sativa* L.) under Malwa plateau condition. M.Sc (Ag.) Thesis; 2018.
 9. Ajay, S., Naruka, S. and Shaktawat, N. 2016. Effect of row spacing and nitrogen on growth and yield of coriander (*Coriandrum sativum* L.). *Journal of Krishi Vigyan*, 5(1): 49-53.
 10. Diwan G, Bisen BP, Maida P. Effect of nitrogen doses and row spacing on growth and seed yield of coriander (*Coriandrum sativum* L.). *International Journal of Chemical Studies*. 2018;6(4):2768-2772.
 11. Nethravathi R. Influence of planting geometry and fertilizer levels on the seed yield and quality of coriander (*Coriandrum sativum* L.). M.Sc. (Ag.) Thesis; 2019.

© 2024 Painkara et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112401>