



Influence of Foliar Spray of Brassinosteroids (BR), Salicylic Acid (SA) and Gibberellic Acid (GA₃) on Vegetative Growth and Flowering Parameters of Cucumber (*Cucumis sativus* L) cv. Arpit.

Netra Pal Yadav^{a++*}, Vijay Bahadur^{a#}, Gyanendra Singh^{a++}
and Nikhil Vikram Singh^{a++}

^a Department of Horticulture, NAI, SHUATS, Prayagraj, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i121498

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94777>

Original Research Article

Received: 06/10/2022

Accepted: 12/12/2022

Published: 14/12/2022

ABSTRACT

An investigation entitled "Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative and flowering parameters of Cucumber (*Cucumis sativus* L) cv. Arpit" was carried out at the Vegetable Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology & Science, Prayagraj, (U.P.) - 211007 during the

⁺⁺ Ph.D. Scholar;

[#] Associate Professor and Head;

*Corresponding author: E-mail: rajyadav1358@gmail.com;

year 2018-19 and 2019-2020. The experiment was laid out in a Randomized Block Design with 15 treatments and 3 replications and each replicated thrice. Source of variables were Brassinosteroids (BR) 0.05 PPM, Brassinosteroids (BR) 0.10 PPM, Brassinosteroids (BR) 0.5 PPM, Brassinosteroids (BR) 1.0 PPM, Brassinosteroids (BR) 2.0 PPM, Salicylic acid (SA) 0.05 mM, Salicylic acid (SA) 0.10mM, Salicylic acid (SA) 0.2mM, Salicylic acid (SA) 0.5mM, Salicylic acid (SA) 1.0mM, Gibberellic acid (GA3) 25 PPM, Gibberellic acid(GA3) 50 PPM, Gibberellic acid(GA3) 100 PPM, Gibberellic acid (GA3) 150 PPM and combination of them with control treatment taken as 15 treatments. As far as the vegetative growth and flowering is concerned, application of different treatment of plant growth regulators significantly enhanced the Length of main vine (cm), Number of branches per plant, Number of leaves per plant, Diameter of main stem (cm), Inter-nodal distance (cm), Leaf area (cm²), Days to first flower bud initiate (DAS), Days taken to 50% flowering, Number of male flower per plant, Number of female flower per plant at all successive stages. The maximum vine length (143.17 and 146.04cm), number of leaves per plant (93.74 and 94.74), number of branches per plant (19.49 and 19.82), diameter of main stem (14.70 and 16.02cm), leaf area (277.42 and 280.90 cm²), internodal distance (7.27 and 6.97cm) and as per flowering parameters lowest days taken to first flower bud initiation (36.66 and 35.73 days), days taken to 50% flowering (42.40 and 422.73 days), number of male flower per plant (91.71 and 92.05) and number of female flower per plant (47.59 and 46.92) were recorded in both successive year with T₅ (Brassinosteroids (BR) 2.0 PPM) and minimum growth and flowering were observed in T₀ (control).

Keywords: *Cucumber; vegetative growth; flowering; brassinosteroids; salicylic acid; gibberellic acid.*

1. INTRODUCTION

Cucumber (*Cucumis sativus* L.) is a cross pollinated and widely grown vegetable crop in the Cucurbitaceae family, with chromosome number 2n=14. It is an indigenous vegetable to India [1]. It is often a monoecious, annual, trailing or climbing vine with hirsute or scabrous stems and triangular ovate leaves with shallow and sharp sinuses [2]. At the leaf axils, unbranched lateral tendrils appeared. Flower clusters form in leaf axils as the lateral branches grow [3]. It is a warm-season crop with minimal to no frost resistance. Cucumber growth and development are aided by temperatures over 20-30°C.

It is widely farmed in the Indian states of Madhya Pradesh, Tamil Nadu, Uttar Pradesh, Andhra Pradesh, Kerala, and Maharashtra. It covers an area of 41 million ha in India, produces 641 MT, and has a productivity of 15.63 t/ha (Handbook of Horticulture Statistics 2019-20).

Depending on the cultivar, area, and soil climate, the plant begins flowering early and produces marketable fruits within two or three months. Flowering is a critical stage in the development of cucurbits since it determines fruiting and yield. Cucumber is a monoecious plant, which means that the first flowers that develop near the base of a cucumber plant are male. A week after the male flower initiation, the female flowers develop, with the little cucumber fruit at the base [4]. Cucumbers have a high water content while

being low in calories, fat, cholesterol, and salt. The fruit is an elongated, round triangular fake berry or pepo. Its size, shape, and colour differ according to the cultivar. Fruits are beneficial to persons who have jaundice, constipation, or indigestion. It is eaten raw with salt and pepper, or as part of a salad with pickles. The fruit pulp is used to make mash cakes.

Cucumber's reaction to plant growth regulators displays a remarkable spectrum of floral morphology, including staminate, pistillate, and hermaphrodite flowers that appear in a variety of configurations and provide a variety of sexual expression. Growth regulators have a huge impact on sex expression and blooming in cucumber crops, either suppressing male flowers or increasing the amount of female flowers [5] with no negative consequences for the environment or human health. Plant growth regulators are also used to control cucumber plant vegetative development, which increases plant population per unit area in terms of production [6].

Brassinosteroids are present in a variety of species ranging from lower plants to higher plants. They are steroidal plant hormones that help plants grow and develop. BR may show a keen interest in the role of horticulture crops. It was characterised by a decrease in the quantity of male flowers in the early stages of growth and a promotion of female flower initiation in the main stalk. Another study on cucumber found that BS

is critical during early fruit development [7]. the treatment of tomato plants with BS, at flowering stage led to the enhancement of the number and weight of tomato [8]. The effects of brassinosteroids on both vegetative and reproductive development are investigated and compared to known BR physiological responses such as cell elongation and division, vascular differentiation, blooming, pollen formation, and photomorphogenesis [9]. They also have an impact on cotyledon development, root elongation, leaf creation and growth, and plant biomass. Exogenous BS also enhances the activity of other enzymes such as carbonic anhydrase and nitrate reductase [10,11], rubisco [12]. BS have also been found to affect whole physiology of the plant, starting from seed germination to harvest or seed maturation. Application of BS has been found to enhance the seed germination in chickpea [13], Indian mustard [14] and tobacco [15].

Salicylic acid (SA) is a phytohormone that has recently been added to the family of phytohormones for appropriate plant growth development and induction of tolerance to both biotic and abiotic stresses. Growth-stimulating effects of SA have been reported in soybean [16], wheat [17], maize [18]. SA is a phenolic endogenous growth regulator that controls a variety of physiological processes in agricultural plants such as stomatal closure, ion uptake, ethylene biosynthesis inhibition, and transpiration [19,17]. SA found that 4 μ M SA promotes flower bud formation from tobacco callus [20].

Gibberellic acid is important in seed germination, endosperm mobilisation, stem elongation, leaf expansion, shortening maturation time, and boosting flower and fruit set and composition [21]. GA₃ slows senescence, promotes chloroplast growth and development, and increases photosynthetic efficiency, perhaps leading to enhanced production [22].

2. MATERIALS AND METHODS

The experiment entitled "Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative and flowering parameters of Cucumber (*Cucumis sativus* L) cv. Arpit" was carried out during the summer season of the year 2018-2019 and 2019-20. The experiment was laid out in a Randomized Block Design with 15 treatments and 3 replications and each replicated thrice. Department of Horticulture, Naini Agricultural

Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, provided all of the necessary equipment for the experiment.

It reaching a height of 98 meters above the sea level and is located at 25°.8'N latitude and 81°.50' E longitude. Prayagraj district is situated in Uttar Pradesh's subtropical zone, which has extreme heat summers and mild winters. The area's peak temperature is 46°C–48°C, and it seldom goes below 4°C–5°C. The relative humidity level varies between 20 and 94 percent. The annual average rainfall in this area is 1013.4 mm.

Source of variable were T₀ (Control), T₁ Brassinosteroids 0.05 PPM, T₂ Brassinosteroids 0.10PPM, T₃ Brassinosteroids 0.5 PPM, T₄ Brassinosteroids 1.0 PPM, T₅ Brassinosteroids 2.0PPM, T₆ Salicylic acid 0.05 mM, T₇ Salicylic acid 0.10 mM, T₈ Salicylic acid 0.2 mM, T₉ Salicylic acid 0.5 mM, T₁₀ Salicylic acid 1.0mM, T₁₁ Gibberellic acid 25 PPM, T₁₂ Gibberellic acid 50 PPM, T₁₃ Gibberellic acid 100 PPM, T₁₄ Gibberellic acid 150 PPM. The observations were recorded on vegetative growth and flowering parameters of cucumber as Vine length (cm), No. of leaves /plant, No. of branches /plant, Diameter of main stem (mm), Internode distance (cm), Leaf area (cm²), Days to first flower bud initiation, Days taken to 50% flowering, No. of male flower/plant, No. of female flower/plant were studied in the investigation. Statistical analysis of experiment was laid out in randomized block design with three replications.

3. RESULT AND DISCUSSION

The data on various observations recorded during experimentation were subjected to statistical analysis in Randomized Block Design in order to find out the significance of different treatments by using the analysis of variance. The results have been integrated along with the corresponding tables. In terms of vegetative parameters, it was found that Vine length, number of leaves, number of branches, diameter of main stem(mm), Internodal distance(cm) and Leaf area(cm²) were significantly increased during the both year by different treatments of plant growth regulators at all successive stage of growth.

It is clear from the Table 1 that the maximum vine length at 90 days (143.17 cm and 146.04 cm in 2018-19 and 2019-20 respectively and 144.60 cm as pooled) closely conformity to the

Ramani et al. [23], kumar & kaur [24] and Hirpara et al. [25] Maximum Number of leaves at 90 days (93.74 and 94.74 in 2018-19 and 2019-20 respectively and 94.24 as pooled) closely associated with Lrfan et al. [26] and Vishnu ritti et al. [27]; Number of branches (19.49 and 19.82 in 2018-19 and 2019-20 respectively and 94.24 as pooled) associated with Ghosh et al. [28]; maximum Diameter of main stem (14.70 mm and 16.02 mm in 2018-19 and 2019-20 respectively and 15.36 mm when pooled) closely with Vishnu ritti et al. [27]; and maximum leaf area (277.42 cm² and 280.90 cm² in 2018-19 and 2019-20 respectively and 279.16 cm² when pooled) close conformity to the karuppaiah et al. [29] and khatoon et al. [30] and maximum Internodal distance (7.27 and 6.97 in 2018-19 and 2019-20 respectively and 7.12 when pooled) closely associated with khatoon et al. [31] and Acharya et al. [32] were recorded in T₅ (Brassinosteroids (BR) 2.0 PPM). While minimum Vine length at 90 days (126.62cm, 128.95cm and 127.79cm in 2018-19; 2019-20 and as pooled, respectively), number of leaves (74.53, 75.19 and 74.86 in 2018-19; 2019-20 and as pooled, respectively), Number of branches (12.35, 13.68 and 13.02 in 2018-19, 2019-20 and pooled, respectively), Diameter of main stem (11.05 mm, 11.71mm and 11.38 mm in 2018-19, 2019-20 and pooled, respectively), leaf area (258.07cm², 266.07 cm² and 262.07 cm² in 2018-19, 2019-20 and pooled, respectively), Internodal distance (4.79, 4.39 and 4.59 in 2018-19, 2019-20 and pooled, respectively) were observed with T₀ (Control). According to Papadopoulou and Grumet [33], applying EBL to cucumber plants accelerated and boosted female flower production, and the effects may be mediated, at least in part, by BS-induced ethylene production. Furthermore, external brassinolide treatments were performed on cucumber, melon, and zucchini, and it was discovered that cucumber was more susceptible than zucchini. It was characterised by a decrease in the quantity of male flowers in the early stages of growth and a promotion of female flower initiation in the main stalk. In another study on cucumber, it was demonstrated that BS play an important role during early fruit development [7]. Growth has been suggested to be promoted by increasing the flexibility of the cell wall, followed by hydrolysis of starch to sugars, which reduces the water potential of the cell, resulting in the entrance of water into the cell, inducing elongation. Sargent [34].

In term of Flowering parameters, it was found that Days to first flower bud initiate (DAS), Days taken to 50 % flowering, Number of male flower per plant, Number of female flower per plant were significantly increased by different treatments of plant growth regulator at all successive stages of growth and flowering in both the individual years and when pooled.

Minimum Days to first flower bud initiate (days) (36.66 days and 35.73 days in 2018-19 and 2019-20 respectively and 36.20 days as pooled) closely associated with Abdul et al. [35]. Minimum Days to 50% flowering (days) (42.40 days and 42.73 days in 2018-19 and 2019-20 respectively and 42.57 days as pooled) closely associated with karuppaiah et al. [29] and khatoon et al. [30] and Maximum Number of male flower per plant (91.71 and 92.05 in 2018-19 and 2019-20 respectively and 91.88 as pooled) close conformity with the Khan et al. [36] Kadi et al. [37] Maximum Number of female flower per plant (47.59 and 46.92 in 2018-19 and 2019-20 respectively and 47.26 as pooled) closely associated with khatoon et al. [37], Singh et al. [38] and Pawar et al. [39] were recorded in T₅ (Brassinosteroids (BR) 2.0 PPM). While maximum days to first flower bud initiate (days) (41.40 days and 41.26 days in 2018-19 and 2019-20 respectively and 41.33 days as pooled), maximum days to 50% flowering (days) (46.26 days and 46.53 days in 2018-19 and 2019-20 respectively and 46.40 days as pooled) associated with Ghorbani et al. [40] and minimum Number of male flower per plant (76.41 and 72.07 in 2018-19 and 2019-20 respectively and 74.24 as pooled), minimum Number of female flower per plant (36.32 and 34.65 in 2018-19 and 2019-20 respectively and 35.49 as pooled) were observed with T₀ (Control).

In this study, relatively modest quantities of GA3 delayed the development of the first staminate bloom and increased pistillate flower commencement [41]. In other research, treating tomato plants with BS at the blooming stage increased the number and weight of tomatoes [8]. In field circumstances, the largest crop improvement was obtained when tomato and cucumber plants were treated with EBL twice, first by seed soaking and then by spraying during the flowering stage [42].

Table 1. Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative parameters of Cucumber cv. F₁ hybrid Arpit.

Treatment	Vine length 90 days			Number of leaves per plant 90 days			Number of branches per plant 90 days			Diameter of main stem (mm) 90 days			Leaf area (cm ²) 90 days		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T ₀	126.62	128.95	127.79	74.53	75.19	74.86	12.35	13.68	13.02	11.05	11.71	11.38	258.07	266.07	262.07
T ₁	133.91	136.57	135.24	84.57	85.49	85.03	14.81	15.92	15.37	12.64	13.40	13.02	264.43	272.70	268.57
T ₂	134.57	137.19	135.88	85.28	87.30	86.29	15.24	16.66	15.95	12.99	13.77	13.38	265.60	273.35	269.48
T ₃	139.39	141.31	140.35	90.19	90.89	90.54	17.33	18.53	17.93	13.47	14.28	13.87	270.87	277.93	274.40
T ₄	142.00	144.36	143.18	91.34	93.34	92.34	18.10	18.89	18.50	13.85	15.10	14.47	274.23	279.22	276.73
T ₅	143.17	146.04	144.60	93.74	94.74	94.24	19.49	19.82	19.66	14.70	16.02	15.36	277.42	280.90	279.16
T ₆	129.65	131.99	130.82	79.79	81.45	80.62	13.74	14.46	14.10	11.73	12.44	12.09	260.36	269.68	265.02
T ₇	133.56	135.58	134.57	81.15	83.93	82.54	14.39	15.51	14.95	12.40	13.14	12.77	262.35	270.76	266.56
T ₈	137.08	139.83	138.45	87.65	89.15	88.40	16.08	17.76	16.92	13.15	13.94	13.54	268.35	276.60	272.48
T ₉	138.10	140.74	139.42	89.99	90.67	90.33	16.84	18.47	17.66	13.33	14.13	13.73	269.43	277.75	273.59
T ₁₀	139.96	143.15	141.56	90.33	91.10	90.72	17.58	18.64	18.11	13.56	14.64	14.10	272.73	278.24	275.49
T ₁₁	136.52	139.08	137.80	86.01	88.89	87.45	15.76	17.17	16.47	13.07	13.85	13.46	267.68	274.43	271.06
T ₁₂	137.78	140.15	138.97	89.25	90.33	89.79	16.33	18.33	17.33	13.25	14.05	13.65	269.22	277.38	273.30
T ₁₃	140.31	143.54	141.93	90.60	91.52	91.06	17.75	18.76	18.26	13.67	14.76	14.22	273.38	279.02	276.20
T ₁₄	142.49	145.12	143.80	92.43	94.01	93.22	18.69	19.17	18.93	14.13	15.26	14.70	274.91	280.75	277.83
CD value	6.10	6.22	6.16	3.88	3.94	3.91	0.73	0.78	0.75	0.58	0.62	0.60	8.59	5.50	6.58
S.Ed (±)	2.98	3.03	3.01	1.89	1.92	1.91	0.35	0.38	0.37	0.29	0.30	0.30	4.19	2.68	3.21

Table 2. Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on flowering parameters of Cucumber cv. F₁ hybrid Arpit.

Treatment	Internodal distance (mm)			Days taken to first flower bud initiation (days)			Days taken to 50% flowering (days)			Number of male flower per plant			Number of female flower per plant		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T ₀	4.79	4.39	4.59	41.40	41.26	41.33	46.26	46.53	46.40	76.41	72.07	74.24	36.32	34.65	35.49
T ₁	5.18	4.96	5.07	40.33	40.00	40.17	45.86	45.38	45.62	84.65	84.27	84.46	41.65	40.20	40.93
T ₂	5.29	5.01	5.15	40.26	39.80	40.03	45.72	45.13	45.43	85.08	84.91	85.00	41.75	40.41	41.08
T ₃	6.20	6.11	6.16	38.86	38.53	38.70	44.73	44.63	44.68	87.90	86.18	87.04	43.38	42.38	42.88
T ₄	6.78	6.38	6.58	38.43	38.30	38.36	44.46	43.73	44.10	89.89	88.36	89.13	45.01	42.74	43.88
T ₅	7.27	6.97	7.12	36.66	35.73	36.20	42.40	42.73	42.57	91.71	92.05	91.88	47.59	46.92	47.26
T ₆	5.12	4.78	4.95	41.00	40.33	40.67	46.20	46.20	46.20	81.48	80.48	80.98	40.64	38.64	39.64
T ₇	5.13	4.85	4.99	40.66	40.23	40.45	45.86	46.09	45.98	83.27	83.74	83.51	40.87	40.18	40.53
T ₈	5.38	5.13	5.26	39.93	39.06	39.50	45.06	44.86	44.96	86.18	85.65	85.92	42.75	41.40	42.08
T ₉	6.11	5.78	5.95	38.61	38.68	38.65	44.86	44.73	44.80	87.72	85.85	86.79	43.07	42.30	42.69
T ₁₀	6.50	6.13	6.32	38.43	38.53	38.48	44.63	44.33	44.48	88.85	87.72	88.29	44.63	42.41	43.52
T ₁₁	5.35	5.05	5.20	40.06	39.33	39.70	45.53	44.94	45.24	85.30	85.64	85.47	42.31	40.65	41.48
T ₁₂	5.78	5.53	5.66	39.53	38.93	39.23	44.94	44.75	44.85	86.99	85.66	86.33	42.85	41.65	42.25
T ₁₃	6.71	6.27	6.49	38.42	38.19	38.31	44.53	44.27	44.40	89.69	87.90	88.80	44.97	42.63	43.80
T ₁₄	6.97	6.50	6.74	38.13	37.86	38.00	44.00	43.65	43.83	90.24	90.23	90.24	47.07	43.43	45.25
CD value	0.26	0.25	0.26	1.87	1.75	1.76	2.01	2.00	2.01	3.85	3.80	3.82	1.91	1.84	1.88
S.Ed (±)	0.13	0.12	0.13	0.91	0.85	0.86	0.98	0.98	0.98	1.88	1.86	1.87	0.93	0.90	0.92

4. CONCLUSION

Based on the current experiment, it was established that the treatment T5 (Brassinosteroids (BR) 2.0 PPM) was ideal in terms of vegetative development and flowering characteristics in both the following years 2018-19 and 2019-20 of cucumber. Farmers may be encouraged to utilise those Plant growth regulator combinations after a few more conjoint research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. De Candole A. Origin of Cultivated Plants. Hafnar Publishing Co., New York. 1967; 264.
2. Bailey LH. Manual of Cultivated Plants. Macmillan Company, New York. 1969; 1116.
3. Ahmed M, Hamid A, Akbar Z. Growth and yield performance of six cucumber (*Cucumis sativus* L.) cultivars under agro-climatic conditions of Rawalakot, Azad Jammu and Kashmir. Int. J. Agri. Biol. 2004;6(2):396- 399.
4. Bantoc GB JR. Further studies on sex expression and sex ratio in cucumber as affected by plant regular spray. Indian Journal. Hon. 1964;17: 210- 216.
5. Al-Masoum AA, Al-Masri AA. Effect of ethephon on flowering and yield of monoecious cucumber. Egyptian J. Horti. 1999;26:229–236.
6. Latimer JG. Growth retardants affect landscape performance of Zinnia, Impatiens and Marigold. Hort Sci. 1991;26:557–560.
7. Feng Qing Fu, Wei Hua Mao, Kai Shi, Yan Hong Zhou, Tadao Asami and Jing Quan Yu. A role of brassinosteroids in early fruit development in Cucumber. Journal of Experimental Botany. 2008; 59(9):2299–2308.
8. Balmush GT, Russu MM. Karabdzhak. Effect of epibrassinolide on tomato growth and development. in: Brassinosteroids – biorational ecologically safe regulators of growth and productivity of plants, 4th, Minsk. 1995;22–23.
9. Steven D. Clouse. Brassinosteroids, The Arabidopsis Book. 2011(9);2011.
10. Ali B, Hayat S, Hasan SA, Ahmad A. Effect of root applied 28-homobrassinolide on the performance of *Lycopersicon esculentum*. Sci. Hortic. 2006;110:267–273.
11. Alam MM, Hayat S, Ali B, Ahmad A. Effect of 28-homobrassinolide treatment on nickel toxicity in *Brassica juncea*. Photosynthetica. 2007;45:139–142.
12. Yu JQ, Huang LF, Hu WH, Zhou YH, Mao WH, Ye SF, Nogue S. A role for brassinosteroids in the regulation of photosynthesis in *Cucumis sativus*. J. Exp. Bot. 2004;55:1135–1143.
13. Ali B, Hayat S, Ahmad A. Response of germinating seeds of *Cicer arietinum* to 28-homobrassinolide and/or potassium. Gen. Appl. Plant Physiol. 2005;31:55–63.
14. Sirhindi G. Brassinosteroids: biosynthesis and role in growth, development, and thermotolerance responses. In: Rout, G.R., Das, A.B. (Eds.), Molecular Stress Physiology of Plants. Springer, India. 2013;309–329.
15. Leubner-Metzger G. Brassinosteroids and gibberellins promote tobacco seed germination by distinct pathway. Planta. 2001;213:758–763.
16. Gutiérrez-Coronado MA, Trejo-López C, Larqué-Saavedra A. Effects of salicylic acid on the growth of roots and shoots in soybean, Plant Physiology and Biochemistry. 1998;36:563-565.
17. Shakirova FM, Sakhabutdinova AR, Bezrukova MV, Fatkhutdinova RA, Fatkhutdinova DR. Change in the hormonal status of wheat seedling induced by salicylic acid and salinity. Plant Sci. 2003;164(3):317-22.
18. Gunes A, Inal A, Alpaslan M, Eraslan F, Guneri Bagci E, Cicek N. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. J Plant Physiol. 2007;2007(164): 728:736.
19. Khan W, Prithviraj B, Smith DL. Photosynthetic responses of corn and soybean to foliar application of salicylates. J Plant Physiol. 2003;160(5):485-92.
20. Lee TT, Skoog F. Effect of substituted phenols on bud formation and growth of tobacco tissue culture. Physiol Plant. 1965;18(2):(386-402).

21. Roy R, Nasiruddin KM. Effect of different level of GA3 on growth and yield of cabbage. J Environ Sci Nat Resour. 2011;4:79-82.
22. Yuan L, Xu DQ. Stimulation effect of gibberellic acid short-term treatment on the photosynthesis re-lated to the increase in RuBisCO content in broad bean and soybean. Photosynth Res. 2001;68(1):39-47.
23. Ramani MM, Patel CR, Sharma KM, Amarcholi JJ, Patel AH. Effect of 28-homobrassinolode and Shoot Thinning on Vegetative Growth, Flowering and Yield of Mango cv. Kesar. Adv Life Sci. 2016;5(17). ISSN 2278-3849:7097-9.
24. Kumar S, Kaur G. Effect of pre harvest application of salicylic acid on plant growth parameters and yield of strawberry cv. Chandler. Gurpinder; 2019.
25. Hipara AJ, Vaddoria MA, Jivanni LL, Patel L. JB and Polara AM. Seed yield and quality as influence by plant growth regulators and stages of spray in bitter gourd (*Momordica charantia* L.) An international ejournal. 2014;3(3): 282-7.
26. Irfan M, Alam J, Ahmad I, Ali I, Gul H. Effects of exogenous and foliar applications of brassinosteroid (BRs) and salt stress on the growth, yield and physiological parameters of *Lycopersicon esculentum* (Mill.). Plant Sci Today. 2017;4(3):88-101.
27. Ritti V, Doddagoudar R. vasudevan, basavegowda and mallikarjun kenganal. Influence of foliar spray of boron and plant growth regulators on hybrid seed yield and quality of bottle gourd. Farm J. Sci. 2019;32(3):274-7.
28. Ghosh A, Panda D, Mondal S. Effect of micronutrients and growth regulators on pigment content, nitrate reductase activity and relative leaf water content of chilli (*Capsicum annuum* L.). Plant Cell Biotechnol Mol Biol. 2020;21(13-14): 37-51.
29. Karuppaiah P. Effect of bioregulators and Feso4 on growth, yield and quality of crossandra (*Crossandra infundibuliformis* L.). HIJ. 2019;3(1):1-5.
30. Khatoon F, Kundu M, Mir H, Nandita K, Kumar D. Foliar feeding of brassinosteroid: A potential tool to improve growth, yield and fruit quality of strawberry (*Fragaria x ananassa* Duch.) under nonconventional area. Int J Curr Microbiol Appl Sci. 2020;9(3):733-41. ISSN: 2319-7706.
31. Khatoon F, Kundu M, Mir H, Nahakpam S. Efficacy of foliar feeding of brassinosteroid to improve growth, yield and fruit quality of strawberry (*Fragaria x ananassa* Duch.) grown under subtropical plain. Commun Soil Sci Plant Anal. 2021;52(8):803-14.
32. Acharya SK, Thakar C, Brahmabhatt JH, Joshi N. Effect of plant growth regulators on cucurbits: a review. J Pharmacogn Phytochem. 2020;9(4):540-4.
33. Papadopoulou E, Grumet R. Brassinosteroid-induced femaleness in cucumber and relationship to ethylene production. HortSci. 2005;40(6):1763-7.
34. Sargent JA. The penetration of growth regulators into leaves. Annu Rev Plant Physiol. 1965;16(1):1-12.
35. Baqi A, Manohar RK, Shankar AG. Effect of GA3 and NAA with pruning levels on growth, sex expression and yield attributes of cucumber (*Cucumis sativus* L. Malini F1) under protected condition. Int J Chem Stud. 2018;6(4):1991-6.
36. Khan AS, Chaudhry NY. GA3 improves flower yield in some cucurbits treated with lead and mercury. Afr J Biotechnol. 2006;5(2):149-53.
37. S. Kadi AS, Asati KP, Barche S, Tulasigeri RG. Effect of different plant growth regulators on growth, yield and quality parameters in cucumber (*Cucumis sativus* L.) under polyhouse condition. Int J Curr Microbiol Appl Sci. 2018;7(4): 3339-52.
38. Singh, Pramod Kumar A, Chaturvedi, Varun Kumar A, Bose. Bandana. J Stress Physiol Biochem. Effects of Salicylic Acid On Seedling Growth and Nitrogen Metabolism In Cucumber (*Cucumis Sativus* L.). 2010;6(3):102-13.
39. Pawar V, Meena MK, Jadia M, Basediya SS. Study of influence of plant growth regulators (PGRs) on yield attributing characters in cucumber. Int J Pure App Biosci. 2019;7(3):318-24.
40. Ghorbani P, Eshghi S, Haghi H. Effects of brassinosteroid (24- epibrassinolide) on yield and quality of grape (*Vitis vinifera* L.). Thompson Seedless'. Vitis. 2017;56: 113-7.

41. Wang QM, Zeng GW. Morphological and histochemical study on sex differentiation in *Momordica charantia* Journal of Zhejiang Agriculture University. 1997; 23:149-53.
42. Churikova VV, Derevshchukov SN. Registration trials of the growth regulator Epin on tomato and cucumber. Technical report of Voronezh State University; 1997.

© 2022 Yadav et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/94777>